TOPCARE ALLERGY AND CONGESTION RELIEF- loratadine, pseudoephedrine sulfate tablet, film coated, extended release Topco Associates LLC

Topco Associates LLC. Allergy & Congestion Relief Drug Facts

Active ingredients (in each tablet)

Loratadine 5 mg

Pseudoephedrine sulfate 120 mg

Purpose

Antihistamine Nasal decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- sneezing
- itchy, watery eyes
- runny nose
- itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- thyroid disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland

• liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed.

Taking more than directed may cause drowsiness.

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

• do not divide, crush, chew or dissolve the tablet

adults and children 12 years and over	1 tablet every 12 hours; not more than 2 tablets in
	24 hours
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- each tablet contains: calcium 25 mg
- do not use if blister unit is broken or torn
- store between 20° to 25°C (68° to 77°F)
- keep in a dry place

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate, hypromellose, lactose monohydrate, magnesium stearate, pharmaceutical ink, povidone, titanium dioxide

Questions or comments?

1-888-423-0139

Package/Label Principal Display Panel

COMPARE TO CLARITIN-D[®] 12 HOUR EXTENDED RELEASE TABLETS ACTIVE INGREDIENTS NON-DROWSY* Allergy & Congestion Relief PSEUDOEPHEDRINE SULFATE 120 mg – LORATADINE 5 mg EXTENDED RELEASE TABLETS NASAL DECONGESTANT – ANTIHISTAMINE OUR PHARMACISTS RECOMMEND INDOOR & OUTDOOR ALLERGIES 12 HOUR RELIEF OF: Nasal and Sinus Congestion Due to Colds or Allergies Sneezing – Runny Nose – Itchy, Watery Eyes Itchy Throat or Nose Due to Allergies 30 EXTENDED RELEASE TABLETS *When taken as directed. See Drug Facts Panel. actual size



Product Informat	tion						
Product Type		HUMAN OTC DRUG	Ite m Co	em Code (Source)		NDC:36800-611	
Route of Administra	tion	ORAL					
Active Ingredient		5					-
Ingredient Name					Basis of Strength		Strengt
LORATADINE (UNII: 7AJO3BO7QN) (LORATADINE - UNII:7AJO3BO7QN) PSEUDOEPHEDRINE SULFATE (UNII: Y9DL7QPE6B) (PSEUDOEPHEDRINE UNII:7CUC9DDI9F)				LORATADINE PSEUDOEPHEDRINE SULFATE		DRINE	5 mg 120 mg
Inactive Ingredie	nts						
		Ingredient Name				5	Strength
CROSCARMELLOSE	SODIUM (UNII:	M28OL1HH48)					
ANHYDRO US DIBASI	C CALCIUM PH	OSPHATE (UNII: L11K75P92J)					
HYPROMELLOSES (U	JNII: 3NXW29V3	WO)					
LACTOSE MONOHYI	DRATE (UNII: EV	VQ57Q8I5X)					
MAGNESIUM STEARA	TE (UNII: 7009	7M6I30)					
POVIDONE (UNII: FZ9	89GH94F)						
		Р)					
TITANIUM DIO XIDE (UNII: 15FIX9V2J	Р)					
TITANIUM DIO XIDE (Product Characte Color	UNII: 15FIX9V2J		Score			no scor	'e
TITANIUM DIO XIDE (Product Characte Color	UNII: 15FIX9V2J eristics		Score Size			no scor 12mm	'e
TITANIUM DIO XIDE (Product Characte Color Shape	UNII: 15FIX9V2J eristics WHITE (to off		Size	nt Code			re
TITANIUM DIO XIDE (Product Characte	UNII: 15FIX9V2J eristics WHITE (to off		Size	nt Code		12mm	'e
TITANIUM DIO XIDE (Product Characte Color Shape Flavor Contains	UNII: 15FIX9V2J eristics WHITE (to off		Size			12mm 7U0	
TITANIUM DIO XIDE (Product Characte Color Shap e Flavor Contains Packaging	UNII: 15FIX9V2J eristics WHITE (to off		Size		ng Start Date	12mm 7U0	
TITANIUM DIO XIDE (Product Characte Color Shape Flavor Contains Packaging # Item Code	UNII: 15FIX9V2J	-white) Package Description	Size		-	12mm 7U0	
TITANIUM DIO XIDE (Product Characte Color Shape Flavor Contains Packaging # Item Code 1 NDC:36800-611-65	UNII: 15FIX9V2J eristics WHITE (to off ROUND 30 in 1 CARTON	-white) Package Description	Size Imprin	Marketi	-	12mm 7U0	
TITANIUM DIO XIDE (Product Character Color Shape Flavor Contains Packaging # Item Code 1 NDC:36800-611-65 1	UNII: 15FIX9V2J Pristics WHITE (to off ROUND 30 in 1 CARTON 1 in 1 BLISTER I	-white) Package Description	Size Imprin	Marketi	-	12mm 7U0	
TITANIUM DIO XIDE (Product Characte Color Shape Flavor Contains Packaging # Item Code 1 NDC:36800-611-65	UNII: 15FIX9V2J Pristics WHITE (to off ROUND 30 in 1 CARTON 1 in 1 BLISTER I	-white) Package Description	n Product	Marketi 05/18/20	-	Marketin	

Revised: 7/2019