

**CHILDRENS PLUS MULTI SYMPTOM COLD- acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine hydrochloride suspension**  
**Walgreens**

-----

**Drug Facts**

**Active ingredients (in each 5 mL)**

**Acetaminophen 160 mg**

Chlorpheniramine maleate 1 mg

Dextromethorphan HBr 5 mg

Phenylephrine HCl 2.5 mg

**Purposes**

**Pain reliever/fever reducer**

Antihistamine

Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves the following cold/flu symptoms:
  - minor aches and pains
  - sore throat
  - cough
  - sneezing and runny nose
  - headache
  - nasal congestion
  - stuffy nose
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses (10 mL) in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- to make a child sleepy
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

### **Ask a doctor before use if the child has**

- liver disease
- glaucoma
- thyroid disease
- diabetes
- heart disease
- high blood pressure
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

### **Ask a doctor or pharmacist before use if your child is taking**

- sedatives or tranquilizers
- the blood thinning drug warfarin

### **When using this product**

- **do not exceed recommended dose (see overdose warning)**
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- nervousness, dizziness or sleeplessness occurs
- cough comes back or occurs with rash or headache that lasts. These could be signs

of a serious condition.

## Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

## Directions

- **this product does not contain directions or complete warnings for adult use**
- mL=milliliter
- shake well before using
- do not give more than 5 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- if needed, repeat dose every 4 hours while symptoms last
- find the right dose on chart below. If possible, use weight to dose, otherwise, use age.

Weight (lb)	Age (yr)	Dose (mL)
48-95	6-11	10 mL
36-47	4-5	do not use unless directed by a doctor
under 36	under 4	do not use

## Other Information

- store between 20-25°C (68-77°F). Do not refrigerate

## Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, D&C red #33, FD&C blue #1, FD&C red #40, flavor, glycerin, microcrystalline cellulose, purified water, sodium benzoate, sorbitol, sucrose, xanthan gum

## Questions or comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

## Principal Display Panel

Compare to the active ingredients in Children's Tylenol® Plus Multi-Symptom Cold††  
Multi-Symptom

Children's

**Cold + Flu**

ACETAMINOPHEN /

PAIN RELIEVER / FEVER REDUCER

CHLORPHENIRAMINE MALEATE /

ANTIHISTAMINE

DEXTROMETHORPHAN HBr /

COUGH SUPPRESSANT

PHENYLEPHRINE HCl /

NASAL DECONGESTANT

- Relieves aches, fever, sore throat, sneezing, cough, runny & stuffy nose
- Oral suspension

AGES 6 -11 YEARS

FL OZ (mL)

Grape flavor

DOSAGE CUP PROVIDED

††This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Plus Multi-Symptom Cold.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

DISTRIBUTED BY: WALGREEN CO

200 WILMOT RD., DEERFIELD, IL 60015

walgreens.com

**Package Label**

**Drug Facts (continued)**

- nervousness, dizziness or sleeplessness occurs
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions**

- this product does not contain directions or complete warnings for adult use
- shake well before using
- do not give more than 5 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- if needed, repeat dose every 4 hours while symptoms last
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

Weight (lb)	Age (yr)	Dose (mL)
48-95	6-11	10 mL
36-47	4-5	do not use unless directed by a doctor
under 36	under 4	do not use

**Other information**

- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients:** carbonylmethylcellulose sodium, citric acid, D&C red #33, FD&C blue #1, FD&C red #40, flavor, glycerin, microcrystalline cellulose, purified water, sodium benzoate, sorbitol, sucrose, xanthan gum

**Questions or comments?**  
Call 1-877-753-3836 Monday-Friday 9AM-5PM EST

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

**PARENTS:**  
Learn about drug medicine abuse.  
www.StopMedicineAbuse.org

**DOSAGE CUP PROVIDED**

†Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.  
‡This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Plus Multi-Symptom Cold. PLD-1253C FC008009

NDC 0363-0034-04



Compare to the active ingredients in Children's Tylenol® Plus Multi-Symptom Cold\*

## MULTI-SYMPOM Children's Cold + Flu

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER  
CHLORPHENIRAMINE MALEATE / ANTIHISTAMINE  
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT  
PHENYLEPHRINE HCl / NASAL DECONGESTANT

- Relieves aches, fever, sore throat, sneezing, cough, runny & stuffy nose
- Oral suspension



AGES 6-11 YEARS

4 FL OZ (118 mL) Grape flavor

NDC 0363-0034-04



Compare to the active ingredients in Children's Tylenol® Plus Multi-Symptom Cold\*

## MULTI-SYMPOM Children's Cold + Flu

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER  
CHLORPHENIRAMINE MALEATE / ANTIHISTAMINE  
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT  
PHENYLEPHRINE HCl / NASAL DECONGESTANT

- Relieves aches, fever, sore throat, sneezing, cough, runny & stuffy nose
- Oral suspension



AGES 6-11 YEARS

4 FL OZ (118 mL) Grape flavor

**Drug Facts**

**Active ingredients (in each 5 mL)**

Active ingredients (in each 5 mL)	Purposes
Acetaminophen 160 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 1 mg	Antihistamine
Dextromethorphan HBr 5 mg	Cough suppressant
Phenylephrine HCl 2.5 mg	Nasal decongestant

**Uses**

- temporarily relieves the following cold/flu symptoms
- minor aches and pains
- sore throat
- cough
- sneezing and runny nose
- headache
- nasal congestion
- stuffy nose
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses (10 mL) in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- to make a child sleepy
- with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if your child has**

- liver disease
- glaucoma
- thyroid disease
- diabetes
- heart disease
- high blood pressure
- a breathing problem such as chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

**Ask a doctor or pharmacist before use if your child is taking**

- the blood thinning drug warfarin
- sedatives or tranquilizers

**When using this product**

- do not exceed recommended dose (see Overdose warning)
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com ©2021 Walgreen Co.

ITEM 510122 W10024-0322-L



3 11917 10357 0

Lot No.:  
Exp. Date:

### WALGREENS Multi-Symptom Children's Cold and Flu

## CHILDRENS PLUS MULTI SYMPTOM COLD

acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine hydrochloride suspension

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-0034
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	1 mg in 5 mL

<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>CARBOXYMETHYLCELLULOSE SODIUM</b> (UNII: K679OBS311)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>D&amp;C RED NO. 33</b> (UNII: 9DBA0SBB0L)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	GRAPE	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0034-04	1 in 1 BOX	04/30/2015	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/30/2015	

**Labeler** - Walgreens (008965063)