### CORICIDIN HBP MAXIMUM STRENGTH COLD, COUGH AND FLUacetaminophen, dextromethorphan hydrobromide liquid Bayer HealthCare LLC.

-----

# Coricidin HBP Maximum Strength Cold , Cough & Flu Liquid

### **Drug Facts**

Active ingredients (in each 30 mL)	Purposes		
Acetaminophen 650 mg	Pain reliever/fever reducer		
Dextromethorphan hydrobromide 20 mg	Cough suppressant		

### Uses

- temporarily relieves these symptoms due to a cold or flu:
  - minor aches and pains
  - headache
  - cough
  - sore throat
- temporarily reduces fever

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

 with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen,ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

## Ask a doctor before use if you have

- liver disease
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that is accompanied with excessive phlegm (mucus)
- a sodium-restricted diet

# Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

# Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

# Directions


- do not take more than recommended dose
- adults and children 12 years and over: take 30 mL in the dosing cup provided every 4 hours. Do not exceed 150 mL (5 doses) in 24 hours or as directed by a doctor.
- children under 12 years of age: do not use

# Other information

- each 30 mL contains: sodium 38 mg
- store between 20<sup>o</sup> to 25<sup>o</sup>C (68<sup>o</sup> to 77<sup>o</sup>F)

**Inactive ingredient**s anhydrous citric acid, FD&C blue #1, FD&C red # 40, flavor, glycerin, maltitol solution, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments? 1-800-317-2165 (Mon - Fri 9AM - 5PM EST)



Approved by eoknj on 07-Feb-2022 15:09

#### New Formula!

Coricidin ®

HBP

COLD RELIEF FOR PEOPLE WITH

### **HIGH BLOOD PRESSURE**<sup>†</sup>

MAXIMUM STRENGTH

NIGHTTIME COLD & FLU Relieves Body Aches & Pains Headache, Fever, Cough, Sore Throat, SUGAR FREE Alcohol Free Cherry †Decongestant Free **Acetaminophen -**Pain Reliever/Fever Reducer Dextromethorphan HBr-Cough Suppressant

12 FL OZ (355 mL)

**CORICIDIN HBP MAXIMUM STRENGTH COLD, COUGH AND FLU** acetaminophen, dextromethorphan hydrobromide liquid **Product Information** HUMAN OTC DRUG Item Code (Source) **Product Type** NDC:11523-0093 **Route of Administration** ORAL **Active Ingredient/Active Moiety Ingredient Name Basis of Strength** Strength 650 mg ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D) ACETAMINOPHEN in 30 mL 20 mg DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) DEXTROMETHORPHAN (DEXTROMETHORPHAN - UNII:7355X3ROTS) **HYDROBROMIDE** in 30 mL **Inactive Ingredients Ingredient Name** Strength PROPYLENE GLYCOL (UNII: 6DC9Q167V3) FD&C RED NO. 40 (UNII: WZB9127XOA) WATER (UNII: 059QF0KO0R) SACCHARIN SODIUM (UNII: SB8ZUX40TY)

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3MQ0SDW1A)

SODIUM CITRATE (UNII: 1Q73Q2JULR)

AN	IHYDROUS CITRIC	C ACID (UNII: XF417D3PSL)		
FD	&C BLUE NO. 1	(UNII: H3R47K3TBD)		
GL	YCERIN (UNII: PDC	C6A3C0OX)		
M/	ALTITOL (UNII: D65	5DG142WK)		
Pa	ackaging			
#	ltem Code	Package Description	Marketing Start Date	Marketing End Date
1		355 mL in 1 BOTTLE; Type 0: Not a Combination Product	04/27/2022	
Μ	larketing l	nformation		
M	larketing l Marketing Category	nformation Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

Labeler - Bayer HealthCare LLC. (112117283)

Revised: 12/2024

Bayer HealthCare LLC.