

MINNERVA II- lidocaine, epinephrine gel
SofTap Inc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

DRUG FACTS

ACTIVE INGREDIENTS (per ml) PURPOSE

- Lidocaine USP - 4%

topical anesthetic

- Epinephrine HCl - 0.4%

vasoconstrictor

EXP0326LOT06

Drug Facts

ACTIVE INGREDIENTS (per ml) PURPOSE

- Lidocaine USP- 4%
topical anesthetic
- Epinephrine HCl-.04%
vasoconstrictor

OTHER INGREDIENTS

• Deionized water, glycerin, sodium bisulphate, sodium metabisulfate, dehydro xanthan gum, chlorobutanol, sodium chloride.

USES

- On adults over 18 only
- Temporarily relieves pain due to cuts, scrapes, or abrasions

DO NOT USE

- With occlusion
- If you have a known allergy or sensitivity to any ingredients.
- If pregnant or breastfeeding
- On large areas of the body.
- In large quantities. Apply only 1-2ml to desired area.
- If you have a history of liver disease or impairment.
- If safety seal is broken before first use.



OTHER INGREDIENTS

- Deionized water, glycerin, sodium bisulphate, sodium metabisulfate, dehydroxanthan

gum, chlorobutanol, sodium chloride

DO NOT USE

- With occlusion
- If you have a known allergy or sensitivity to any ingredients.
- If pregnant or breastfeeding
- On large areas of the body
- In large quantities. Apply only 1-2ml to desired area
- If you have a history of liver disease or impairment
- If safety seal is broken before first use

KEEP OUT OF

- The eye and mouth. If accidental contact occurs, you may feel burning or stinging. Wash with water or eyewash immediately. If your eyes have pain, blurry vision, extreme sensitivity to light, or a feeling of sand in the eye, contact an eye care physician immediately.

WHEN USING THIS PRODUCT


- You may not feel pain. Avoid sources of heat or injury.
- You may have temporary rash, redness, swelling, or itching
- Contact your physician promptly if you notice any unusual effects such as dizziness or drowsiness, difficulty breathing, trembling, chest pain, or irregular heartbeat.

Made in the USA for SofTap

550 N Canyons Pkwy

Livermore, CA 94551

www.softaps.com



Drug Facts (cont.)

KEEP OUT OF

• The eye and mouth. If accidental contact occurs, you may feel burning or stinging. Wash with water or eyewash immediately. If your eyes have pain, blurry vision, extreme sensitivity to light, or a feeling of sand in the eye, contact an eye care physician immediately.

WHEN USING THIS PRODUCT

- You may not feel pain. Avoid sources of heat or injury
- You may have temporary rash, redness, swelling, or itching
- Contact your physician promptly if you notice any unusual effects such as dizziness or drowsiness, difficulty breathing, trembling, chest pain, or irregular heartbeat.

Made in the USA for SofTap
550 N. Canyons Pkwy.
Livermore, CA 94551
www.softaps.com

DIRECTIONS

For external use only. Prior to application, a sensitivity test is advised. Before applying,

cleanse the skin with an alcohol-free or acetone-free cleanser. Apply 1-2 cc to the desired area for up to 6 times per day. Discontinue use if allergic reaction occurs. Store in a cool dark place (up to 80 degrees F) or refrigerate.



WARNINGS

KEEP OUT OF REACH OF CHILDREN

If swallowed seek medical help or contact Poison Control Center immediately.

USES

- On adults over 18 only
- Temporarily relieves pain due to cuts, scrapes, or abrasions
-

EXP0324 LOT06

Drug Facts

ACTIVE INGREDIENTS (per ml) PURPOSE

- Lidocaine USP- 4%
topical anesthetic
- Epinephrine HCl-.04%
vasoconstrictor

OTHER INGREDIENTS

- Deionized water, glycerin, sodium bisulphate, sodium metabisulfate, dehydro xanthan gum, chlorobutanol, sodium chloride.

USES

- On adults over 18 only
- Temporarily relieves pain due to cuts, scrapes, or abrasions

DO NOT USE

- With occlusion
- If you have a known allergy or sensitivity to any ingredients.
- If pregnant or breastfeeding
- On large areas of the body.
- In large quantities. Apply only 1-2ml to desired area.
- If you have a history of liver disease or impairment.
- If safety seal is broken before first use.



WARNINGS

KEEP OUT OF REACH OF CHILDREN



ACTIVE INGREDIENTS (per ml) PURPOSE

- Lidocaine USP - 4%

topical anesthetic

- Epinephrine HCl - .04%

vasoconstrictor

Minnerva II



MINNERVA II

lidocaine, epinephrine gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:82718-2210
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
EPINEPHRINE HYDROCHLORIDE (UNII: WBB0470038) (EPINEPHRINE - UNII:YKH834O4BH)	EPINEPHRINE HYDROCHLORIDE	0.004 g in 10 g
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	0.3 g in 10 g

Inactive Ingredients

Ingredient Name	Strength
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
CHLOROBUTANOL (UNII: HM4YQM8WRC)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	
SODIUM BISULFATE (UNII: BU8V88OWMQ)	
DEHYDROXANTHAN GUM (UNII: 63ZP7I1BQO)	

Product Characteristics

Color	white (Clear Gel)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82718-2210-1	10 g in 1 TUBE; Type 0: Not a Combination Product	05/11/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		05/11/2022	

Labeler - SofTap Inc (780568353)

Registrant - SofTap Inc (780568353)

