

**PAIN RELIEF PM- acetaminophen, aspirin, diphenhydramine citrate tablet, film coated**  
**CVS Pharmacy**

-----  
**CVS 44-674**

***Active ingredients (in each caplet)***

Acetaminophen 250 mg

Aspirin 250 mg (NSAID)\*

Diphenhydramine citrate 38 mg

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

Pain reliever

Nighttime sleep-aid

***Uses***

for the temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product

### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- in children under 12 years of age

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have difficulty in urination due to enlargement of the prostate gland
- you have liver disease
- you have a breathing problem such as emphysema or chronic bronchitis
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have glaucoma
- you have asthma

### **Ask a doctor or pharmacist before use if you are taking**

- a prescription drug for:
  - diabetes
  - gout
  - arthritis
- any other drug, or are under a doctor's care for any serious condition
- sedatives or tranquilizers
- any product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

### **When using this product**

- avoid alcoholic beverages
- drowsiness may occur
- do not drive a motor vehicle or operate machinery

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
- pain gets worse or lasts for more than 10 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

- **do not take more than directed**
- adults and children 12 years and over: take 2 caplets at bedtime, with a full glass of water
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- children under 12 years: do not use

### ***Other information***

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

### ***Inactive ingredients***

corn starch, crospovidone, FD&C blue #1 aluminum lake, hypromellose, maltodextrin, medium chain triglycerides, microcrystalline cellulose, polydextrose, povidone, sodium lauryl sulfate, sodium starch glycolate, sodium sulfate, stearic acid, talc, titanium dioxide, yellow iron oxide, zinc stearate

### ***Questions or comments?***

**1-800-426-9391**

## ***Principal Display Panel***

♥CVSHealth®

Compare to the active ingredients  
in Excedrin® PM Headache†

### **Caplets**

#### **Pain Relief PM**

**ACETAMINOPHEN**, 250 mg

**ASPIRIN (NSAID)**, 250 mg

**DIPHENHYDRAMINE CITRATE**, 38 mg

Pain reliever, Nighttime sleep-aid

- Non-habit forming
- For pain with sleeplessness

**100** CAPLETS

Actual Size

Package Contains One Bottle

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by  
GSK Consumer Healthcare SARL, owner of the registered  
trademark Excedrin® PM Headache.

50844 REV1121A67412

**Distributed by: CVS Pharmacy, Inc.**

One CVS Drive, Woonsocket, RI 02895

© 2022 CVS/pharmacy CVS.com® 1-800-SHOP CVS V-11112

BG879L3  
Z FOLD

Actual Size  
Package Contains One Bottle



Package Contains One Bottle



**Stop use and ask a doctor if:**

- you experience any of the following signs of stomach bleeding:
  - an aching or burning course, dark or bloody stools
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- sleepiness persists continuously for more than 2 weeks.
- insomnia may be a symptom of a serious underlying medical illness.

**Using Fodor's (continued)**

**Stop use and ask a doctor if:**

- you experience any of the following signs of stomach bleeding:
  - an aching or burning course, dark or bloody stools
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- sleepiness persists continuously for more than 2 weeks.
- insomnia may be a symptom of a serious underlying medical illness.

**When using this product**

- do not drive a motor vehicle or operate machinery
- avoid alcoholic beverages
- drowsiness may occur

**When using this product**

- gain reverter/reducer
- any product that contains aspirin, acetaminophen, or any other condition
- sedatives or tranquilizers
- any other drug or are under a doctor's care for any condition
- a prescription for the use of sedatives or tranquilizers

**Do not take Fodor's before use if you are taking**

- any other drug or are under a doctor's care for any condition
- a prescription for the use of sedatives or tranquilizers

**Stop use and ask a doctor if:**

- you experience any of the following signs of stomach bleeding:
  - an aching or burning course, dark or bloody stools
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- sleepiness persists continuously for more than 2 weeks.
- insomnia may be a symptom of a serious underlying medical illness.

**KEEP OUTER PACKAGE  
FOR COMPLETE  
PRODUCT INFORMATION**

**Drug Facts**

**Active Ingredients (in each caplet)**

<b>Purpose</b>	Paracetamol Pain reliever
	Acetaminophen (500 mg) Equivalent to 500 mg (USAPD) <sup>1</sup>
	Diphenhydramine (25 mg) Non-sedating anti-inflammatory drug

**Uses**

For the temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

**Warnings**

Play's a game! Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, it changes in behavior with autism and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**May also cause or may cause a severe allergic reaction, which may include:**

- Hives
- Facial swelling
- Shock

**May also cause or may cause severe skin reactions, which may include:**

- Skin peeling
- Rash
- Skin reddening
- A skin reaction occurs; stop use and seek medical help right away.

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

#693397



050428421772

LOT/EXP "NO PRINT"

PAIN RELIEF PM				
acetaminophen, aspirin, diphenhydramine citrate tablet, film coated				
Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59779-894	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	250 mg	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	250 mg	
DIPHENHYDRAMINE CITRATE (UNII: 4OD433S209) (DIPHENHYDRAMINE - UNII:8GTS82S83M)		DIPHENHYDRAMINE CITRATE	38 mg	
Inactive Ingredients				
Ingredient Name			Strength	
STARCH, CORN (UNII: O8232NY3SJ)				
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)				
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
MALTODEXTRIN (UNII: 7CVR7L4A2D)				
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
POLYDEXTROSE (UNII: VH2XOU12IE)				
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)				
SODIUM LAURYL SULFATE (UNII: 368GB5141J)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
SODIUM SULFATE (UNII: OYPR65R21J)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
TALC (UNII: 7SEV7J4R1U)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)				
ZINC STEARATE (UNII: H92E6QA4FV)				
Product Characteristics				
Color	blue	Score	no score	
Shape	OVAL	Size	17mm	
Flavor		Imprint Code	674	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC 59779-894			

<b>1</b>	NDC:59779-894-12	1 in 1 CARTON	08/31/2016	
<b>1</b>		100 in 1 BOTTLE; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M013		08/31/2016	

**Labeler -** CVS Pharmacy (062312574)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(59779-894)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(59779-894)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(59779-894)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(59779-894)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(59779-894)

Revised: 7/2025

CVS Pharmacy