# IDO DISINFECTION ETHANOL- isopropyl alcohol liquid IDO PHARM

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

\_\_\_\_\_

Isopropyl Alcohol

Water

First aid to help prevent the risk of infection in minor cuts, scrapes or burns.

Keep out of reach of children

For the external use only

For external use only

- If taken internally, serious gastric disturbances will result
- flammable, keep away from heat or flame

Ask a doctor before use if you have deep or puncture wounds, animal bites or serious burns

When using this product

- do not use in or near eyes. In case of contact with eyes, rinse eyes thoroughly with water
- do not apply over large areas of the body
- do not use longer than one week unless directed by a doctor

Stop use and ask a doctor if irritation or rash appears

Keep out of reach of children. In case of ingestion, get medical help or contact a Poison Control Center 1-800-222-1222 or dial 911 immediately

Clean affected area

Apply small amount of this product on the affected area 1-3 times daily

May be covered with a sterile bandage. If bandaged, let dry first

children under 6 years of age should be supervised when using this product

# 70% Isopropyl Alcohol



Topical antiseptic & sanitizer

## **Drug Facts**

Active Ingredients Purpose
Isopropyl Alcohol 70% v/v ....... First aid antiseptic

#### Uses

- first aid to help prevent the risk of infection in minor cuts, scrapes or burns
- consumer antiseptic rub to help reduce pathogens that potentially can cause disease

#### Warnings

#### For external use only

- if taken internally, serious gastric disturbances will result
- flammable, keep away from heat or flame.

Ask a doctor before use if you have ■ deep or puncture wounds, animal bites or serious burns

When using this product ■ do not use in or near eyes. In case of contact with eyes, rinse eyes thoroughly with water

- do not apply over large areas of the body
- do not use longer than one week unless directed by a doctor

#### Stop use and ask a doctor

if irritation or rash appears

**Keep out of reach of children.** In case of ingestion, get medical help or contact a Poison Control Center (1-800-222-1222) or dial 911 immediately.

#### Directions

- clean affected area
- apply small amount of this product on the affected area 1-3 times daily
- may be covered with a sterile bandage. If bandaged, let dry first
- children under 6 years of age should be supervised when using this product


#### Other Information

■ store between 15-30°C (59-86°F). Avoid freezing and excessive heat above 40°C (104°F)

Inactive Ingredients purified water

Questions or comments? Call 1-671-649-6978 Monday-Friday 8AM-5PM (CHST)

ADE IN KOREA anufactured by IDO Pharm



α

## 1 Liter (33.81 FL OZ)

### **IDO DISINFECTION ETHANOL**

isopropyl alcohol liquid

#### **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:77039-020

Route of Administration TOPICAL

#### **Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII: ND2M416302)	ISOPROPYL ALCOHOL	83 mL in 100 mL

Inactive Ingredients			
Ing	redient Name	Strength	
WATER (UNII: 059QF0KO0R)			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:77039-020- 01	500 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/01/2022	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	05/01/2022	

## **Labeler -** IDO PHARM (694853523)

## Registrant - IDO PHARM (694853523)

Establishment				
Name	Address	ID/FEI	Business Operations	
IDO PHARM		694853523	manufacture(77039-020)	

Revised: 5/2022 IDO PHARM