

**CD2-FB ACTIVATOR- lactic acid spray, metered
Wausau Chemical**

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Disposal:

Unused teat dip may be diluted with water and flushed down drain. Do not reuse containers. Empty containers should be thoroughly rinsed with water and taken to a recycling center.

Material is not considered hazardous by the OSHA Hazard Communication Standard (29 CFR 1910.1200).

First Aid:

IF SWALLOWED: drink large quantities of water and get medical attention.

IF IN EYES: flush eyes with large quantities of water and get medical attention if irritation occurs.

Handling Instructions:

For external use only. Not for use in sanitizing dairy equipment. Avoid contact with eyes. Avoid contact with food.

Storage & Handling:

Do not mix with any other teat dip or other product. Store at room temperature and out of direct sunlight.

Avoid freezing - If product is exposed to freezing temperatures, components must be mixed thoroughly prior to use.

DIRECTIONS:

Measure equal volumes of CD2 Base and CD2 Activator into a clean dip cup/container and mix until the color is uniform throughout. Do not dilute. Mix only enough product for one milking of the herd. Dip cups should be washed after each milking.

APPLICATION:

Pre-Milking: If teats are visibly dirty, wash and dry teats with a single service towel prior to dipping. Before each cow is milked, dip the teats as far up as possible. Leave CD2 teat dip on teats for at least 15-30 seconds. Wipe teats dry using a single service towel before milking.

Post-Milking: Immediately after milking, dip teats at least 2/3 of their length in CD2 teat dip. Allow to air dry. DO NOT WIPE. CD2 teat dip can be used as a post-dip alone, or as a pre- and post-milking teat dip.

Always use freshly mixed, full strength CD2 teat dip. If product in dip cup becomes visibly dirty, discard contents and fill with fresh CD2 teat dip. CD2 teat dip may be diluted with water and safely flushed down the drain.

Note 1: If teat irritation occurs, discontinue use until irritation subsides. Consult your veterinarian and milking equipment service personnel if irritation persists.

Note 2: CD2 should be used with a compatible pre-dip or udder wash.

Active Ingredient:

Lactic Acid.....2.5%

Inactive Ingredients:

Glycerin.....10.0%

Surfactants, water.....87.5%

Wausau Chemical Corporation

2001 N River Drive

Wausau, WI 54403

800-950-6656

CD2-FB Activator



PRE- & POST-MILKING TEAT DIP FOR USE ONLY WITH CD2 BASE

Active Ingredient: 2.5% Lactic Acid

Contains: 10% Glycerin

Hazards:

Material is not considered hazardous by the OSHA Hazard Communication Standard (29 CFR 1910.1200).

CAUTION:

For external use only. Not for use in sanitizing dairy equipment. Do not mix with any other teat dip or other product. Avoid contact with food. Store at room temperature and out of direct sunlight. Avoid contact with eyes.

FIRST AID:

If swallowed: drink large quantities of water, and get medical attention. If in eyes: flush eyes with large quantities of water, and get medical attention if irritation occurs.

DISPOSAL:

Unused teat dip may be diluted with water and flushed down drain. Do not reuse containers. Empty containers should be thoroughly rinsed with water and taken to a recycling center.

STORAGE:

Avoid Freezing – If product is exposed to freezing temperatures, components must be mixed thoroughly prior to use.

DIRECTIONS FOR USE:

Dispense equal volumes of CD2 Base and CD2-FB Activator through a foaming unit. Do not dilute.

APPLICATION:

Pre-Milking: If teats are visibly dirty, wash and dry teats with a single service towel prior to application. Apply even coating of CD2 foam to each teat. Leave CD2 on teats for at least 15-30 seconds. Wipe teats dry using a single service towel before milking.

Post-Milking: Immediately after milking, Apply even coating of foam to each teat. Allow to air dry. **DO NOT WIPE.** CD2 can be used as a post-dip alone, or as a pre- and post-milking teat treatment.

CD2 may be diluted with water and safely flushed down the drain.

Note 1: If teat irritation occurs, discontinue use until irritation subsides. Consult your veterinarian and milking equipment service personnel if irritation persists.

Note 2: CD2 should be used with a compatible pre-dip or udder wash.

 WAUSAU CHEMICAL CORPORATION 2001 N RIVER DRIVE WAUSAU, WI 54403 800-950-6656	Contents:	Mfg Code: 200106
	<input type="checkbox"/> 15 Gallons (64892-025-17)	Exp. Date: MMDDYY
	<input type="checkbox"/> 55 Gallons (64892-025-21)	
	<input type="checkbox"/> 275 Gallons (64892-025-88)	

CD2-FB ACTIVATOR

lactic acid spray, metered

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:64892-022
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LACTIC ACID (UNII: 33X04XA5AT) (LACTIC ACID - UNII:33X04XA5AT)	LACTIC ACID	25 mg in 1 L

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:64892-022-17	56.78 L in 1 CONTAINER		
2	NDC:64892-022-21	208.2 L in 1 DRUM		
3	NDC:64892-022-88	1040 L in 1 CONTAINER, FLEXIBLE INTERMEDIATE BULK		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		01/06/2020	

Labeler - Wausau Chemical (006136220)

Registrant - Wausau Chemical (006136220)

Establishment

Name	Address	ID/FEI	Business Operations
Wausau Chemical		006136220	manufacture, api manufacture

Revised: 1/2020

Wausau Chemical