

**PRESSURE PLUS PAIN PLUS MUCUS NON-DROWSY- acetaminophen, guaifenesin, phenylephrine hcl tablet, film coated**

**Walgreen Company**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Walgreens 44-527C-PPM**

***Active ingredients (in each caplet)***

Acetaminophen 325 mg  
Guaifenesin 200 mg  
Phenylephrine HCl 5 mg

***Purpose***

Pain reliever/fever reducer  
Expectorant  
Nasal decongestant

***Uses***

- temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold:
  - sinus congestion and pressure
  - minor aches and pains
  - nasal congestion
  - headache
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a

doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

#### **Ask a doctor before use if you have**

- diabetes
- heart disease
- thyroid disease
- liver disease
- high blood pressure
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)

#### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

#### **When using this product**

**do not exceed recommended dosage.**

#### **Stop use and ask a doctor if**

- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

#### **If pregnant or breast-feeding,**

ask a health professional before use.

#### **Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### ***Directions***

- **do not take more than directed**
- adults and children 12 years and over
  - take 2 caplets every 4 hours
  - swallow whole - do not crush, chew, or dissolve
  - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

#### ***Other information***

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15° - 30°C (59°-86°F)
- see end flap for expiration date and lot number

***Inactive ingredients***

corn starch, crospovidone, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

***Questions or comments?***

**1-800-426-9391**

**Principal Display Panel**

***Walgreens***

**Compare to SUDAFED PE<sup>®</sup> Pressure + Pain + Mucus active ingredients<sup>‡‡</sup>**

NDC 0363-4120-08

NON-DROWSY

**Wal-Phed<sup>®</sup> PE**

PRESSURE + PAIN + MUCUS

**ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER**

**GUAIFENESIN / EXPECTORANT / PHENYLEPHRINE HCl / NASAL DECONGESTANT**

MULTY-SYMPTOM

- Relieves sinus pressure/congestion, sinus headache & chest congestion
- Pseudoephedrine free

**24 CAPLETS**                  Actual Size

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

Walgreens Pharmacist Recommended

Walgreens Pharmacist Survey

<sup>‡‡</sup>This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Sudafed PE<sup>®</sup> Pressure + Pain + Mucus.

50844 ORG091752708

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

**100% SATISFACTION GUARANTEED**

walgreens.com    ©2019 Walgreen Co.

ORG00319-F

(Questions or comments? 1-800-426-9391

**Drug Facts (continued)**  
polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

**Inactive ingredients** corn starch, croscopolone, flavor, magnesium stearate, maltoedextrin, microcrystalline cellulose, polyethylene glycol,

■ see end flap for expiration date and lot number

**Drug Facts (continued)**

**Other information** ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN  
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

**Directions** ■ do not take more than directed  
■ adults and children 12 years and over  
■ take 2 caplets every 4 hours  
■ swallow whole - do not crush, chew, or dissolve  
■ do not take more than 10 caplets in 24 hours  
■ children under 12 years: ask a doctor

■ If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

When using this product do not exceed recommended dosage. Stop use and ask a doctor if ■ redness or swelling is present ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**  
■ cough that occurs with too much phlegm (mucus)  
■ Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.  
When using this product do not exceed recommended dosage. Stop use and ask a doctor if ■ redness or swelling is present ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Warnings** ■ This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product ■ Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ rash ■ skin redness ■ blisters ■ If a skin reaction occurs, stop use and seek medical help right away. Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Drug Facts (continued)**  
■ helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive  
**Warnings** ■ This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product ■ Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ rash ■ skin redness ■ blisters ■ If a skin reaction occurs, stop use and seek medical help right away. Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Drug Facts (continued)**  
■ cough that occurs with too much phlegm (mucus)  
■ Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.  
When using this product do not exceed recommended dosage. Stop use and ask a doctor if ■ redness or swelling is present ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**Drug Facts (continued)**  
**Uses** ■ temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold: ■ headache ■ nasal congestion ■ sinus congestion and pressure ■ temporarily reduces fever

**Drug Facts**  
**Active ingredients (in each caplet)**  
Acetaminophen 325 mg... Pain reliever/fever reducer  
Guaifenesin 200 mg... Expectorant  
Phenylephrine HCl 5 mg... Nasal decongestant

**Purpose** Pain reliever/fever reducer, Expectorant, Nasal decongestant

**KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
Walgreens.com ©2019 Walgreen Co.



B-2201-527C-08PPMH  
OR0091752708

**Walgreens**

NON-DROWSY

**Wal-Phed® PE**

PRESSURE + PAIN + MUCUS

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER

GUAIFENESIN / EXPECTORANT / PHENYLEPHRINE HCl / NASAL DECONGESTANT

MULTI-SYMPTOM

- Relieves sinus pressure/congestion, sinus headache & chest congestion
- Pseudoephedrine free

24 CAPLETS



ACTUAL SIZE

Compare to Sudafed PE® Pressure + Pain + Mucus active ingredients!†

NDC 0363-4120-08

NEW

Walgreens 44-527C

**PRESSURE PLUS PAIN PLUS MUCUS NON-DROWSY**

acetaminophen, guaifenesin, phenylephrine hcl tablet, film coated

Walgreens Pharmacist Recommended  
Walgreens Pharmacist Survey  
†This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Sudafed PE® Pressure + Pain + Mucus.  
50844  
ORG0091752708

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

No Print/No Varnish  
Lot and Expiration N

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-4120
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE (UNII: 2S7830E561)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	

## Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	44;527
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-4120-08	2 in 1 CARTON	08/06/2005	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	08/06/2005	

---

**Labeler** - Walgreen Company (008965063)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		832867894	MANUFACTURE(0363-4120)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		832867837	PACK(0363-4120)

Revised: 5/2020

Walgreen Company