

**EQUALINE FIRST AID ANTIBIOTIC- bacitracin, neomycin, polymyxin b,  
pramoxine hcl ointment  
United Natural Foods, Inc. dba UNFI**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**SuperValu, Inc. First Aid Antibiotic Drug Facts**

**Active ingredient (in each gram)**

Bacitracin zinc 500 units

Neomycin sulfate equivalent to 3.5 mg of neomycin base

Polymyxin B 10,000 units as polymyxin B sulfate

Pramoxine HCl 10 mg

**Purpose**

First aid antibiotic

External analgesic

**Uses**

first aid to help prevent infection and for temporary relief of pain or discomfort in minor:

- cuts
- scrapes
- burns

**Warnings**

**For external use only.**

**Do not use**

- if you are allergic to any of the ingredients
- in the eyes
- over large areas of the body

**Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burns

## **Stop use and ask a doctor if**

- you need to use longer than 1 week
- condition persists or gets worse
- symptoms persist for more than 1 week, or clear up and occur again within a few days
- rash or other allergic reaction develops

## **Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## **Directions**

- adults and children 2 years of age or older:
- clean the affected area
- apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage
- children under 2 years of age: ask a doctor

## **Other information**

- store at 20-25°C (68-77°F)

## **Inactive ingredients**

white petrolatum

## **Questions or comments?**

**1-877-932-7948**

## **Principal Display Panel**

compare to Neosporin® active ingredients

helps prevent infection in minor:

cuts

scrapes

burns

first aid antibiotic pain relieving ointment

compare to Neosporin® active ingredients

maximum strength ointment


first aid antibiotic + pain relief


bacitracin zinc - neomycin sulfate - polymyxin B sulfate - pramoxine HCl

first aid antibiotic/pain relieving ointment

NET WT 0.5 OZ (14g)





|  |   |
|--|---|
| <p><b>Drug Facts</b></p> <p><b>Active ingredients (in each gram)</b></p> <p>Bacitracin zinc 500 units..... First aid antibiotic<br/> Neomycin sulfate equivalent to 35 mg of neomycin base.... First aid antibiotic<br/> Polymyxin B 10,000 units as polymyxin B sulfate..... First aid antibiotic<br/> Pramoxine HCl 10 mg..... External analgesic</p> <p><b>Purpose</b></p>  | <p><b>Drug Facts (continued)</b></p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> <li>■ you need to use longer than 1 week</li> <li>■ condition persists or gets worse</li> <li>■ symptoms persist for more than 1 week, or clear up and occur again within a few days</li> <li>■ rash or other allergic reaction develops</li> </ul> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>■ adults and children 2 years of age or older: <ul style="list-style-type: none"> <li>■ clean the affected area</li> <li>■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily</li> <li>■ maybe covered with a sterile bandage</li> </ul> </li> <li>■ children under 2 years of age ask a doctor</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ store at 20-25°C (68-77°F)</li> </ul> <p><b>Inactive ingredient</b> white petrolatum</p> <p><b>Questions or comments?</b> 1-877-932-7948</p> |
| <p><b>Uses</b> First aid to help prevent infection and for temporary relief of pain or discomfort in minor:</p> <ul style="list-style-type: none"> <li>■ cuts</li> <li>■ scrapes</li> <li>■ burns</li> </ul> <p><b>Warnings</b></p> <p><b>For external use only.</b></p> <p><b>Don't use</b></p> <ul style="list-style-type: none"> <li>■ if you are allergic to any of the ingredients</li> <li>■ in the eyes</li> <li>■ over large areas of the body</li> </ul> <p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>■ deep or puncture wounds</li> <li>■ animal bites</li> <li>■ serious burns</li> </ul> | <div style="text-align: center; margin-top: 20px;"> <p><b>Code Area</b></p>  </div>  |

| <b>EQUALINE FIRST AID ANTIBIOTIC</b>  |                         |                           |               |
|---|-------------------------|---------------------------|---------------|
| bacitracin, neomycin, polymyxin b, pramoxine hcl ointment                       |                         |                           |               |
| Product Information   |                         |                           |               |
| <b>Product Type</b>   | HUMAN OTC DRUG          | <b>Item Code (Source)</b> | NDC:41163-040 |
| <b>Route of Administration</b>  | TOPICAL                 |                           |               |
| Active Ingredient/Active Moiety   |                         |                           |               |
| Ingredient Name   | Basis of Strength       | Strength                  |               |
| <b>BACITRACIN ZINC</b> (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)        | BACITRACIN              | 500 [USP'U] in 1 g        |               |
| <b>NEOMYCIN SULFATE</b> (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)         | NEOMYCIN                | 3.5 mg in 1 g             |               |
| <b>POLYMYXIN B SULFATE</b> (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)   | POLYMYXIN B             | 10000 [USP'U] in 1 g      |               |
| <b>PRAMOXINE HYDROCHLORIDE</b> (UNII: 88AYB867L5) (PRAMOXINE - UNII:068X84E056) | PRAMOXINE HYDROCHLORIDE | 10 mg in 1 g              |               |
| Inactive Ingredients  |                         |                           |               |
| Ingredient Name   | Strength                |                           |               |
| <b>PETROLATUM</b> (UNII: 4T6H12BN9U)  |                         |                           |               |

**Product Characteristics**

|                 |                      |                     |  |
|-----------------|----------------------|---------------------|--|
| <b>Color</b>    | YELLOW (Translucent) | <b>Score</b>        |  |
| <b>Shape</b>    |                      | <b>Size</b>         |  |
| <b>Flavor</b>   |                      | <b>Imprint Code</b> |  |
| <b>Contains</b> |                      |                     |  |

**Packaging**

| # | Item Code        | Package Description                               | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:41163-040-64 | 1 in 1 CARTON                                     | 04/02/2015           |                    |
| 1 |                  | 28 g in 1 TUBE; Type 0: Not a Combination Product |                      |                    |
| 2 | NDC:41163-040-58 | 1 in 1 CARTON                                     | 04/02/2015           | 02/28/2021         |
| 2 |                  | 14 g in 1 TUBE; Type 0: Not a Combination Product |                      |                    |

**Marketing Information**

| Marketing Category  | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part333B                                 | 04/02/2015           |                    |

**Labeler** - United Natural Foods, Inc. dba UNFI (943556183)

Revised: 8/2023

United Natural Foods, Inc. dba UNFI