

**EQUALINE FIRST AID ANTIBIOTIC- bacitracin, neomycin, polymyxin b,
pramoxine hcl ointment
United Natural Foods, Inc. dba UNFI**

SuperValu, Inc. First Aid Antibiotic Drug Facts

Active ingredient (in each gram)

Bacitracin zinc 500 units

Neomycin sulfate equivalent to 3.5 mg of neomycin base

Polymyxin B 10,000 units as polymyxin B sulfate

Pramoxine HCl 10 mg

Purpose

First aid antibiotic

External analgesic

Uses

first aid to help prevent infection and for temporary relief of pain or discomfort in minor:

- cuts
- scrapes
- burns

Warnings

For external use only.

Do not use

- if you are allergic to any of the ingredients
- in the eyes
- over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if

- you need to use longer than 1 week

- condition persists or gets worse
- symptoms persist for more than 1 week, or clear up and occur again within a few days
- rash or other allergic reaction develops

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 2 years of age or older:
- clean the affected area
- apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage
- children under 2 years of age: ask a doctor

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

white petrolatum

Questions or comments?

1-877-932-7948

Principal Display Panel

compare to Neosporin® active ingredients

helps prevent infection in minor:

cuts

scrapes

burns

first aid antibiotic pain relieving ointment

compare to Neosporin® active ingredients

maximum strength ointment

first aid antibiotic + pain relief

bacitracin zinc - neomycin sulfate - polymyxin B sulfate - pramoxine HCl

first aid antibiotic/pain relieving ointment

NET WT 0.5 OZ (14g)





<p>Drug Facts</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 50%;">Active ingredients (in each gram)</th> <th style="width: 50%;">Purpose</th> </tr> <tr> <td>Bacitracin zinc 500 units.....</td> <td>First aid antibiotic</td> </tr> <tr> <td>Neomycin sulfate equivalent to 35 mg of neomycin base....</td> <td>First aid antibiotic</td> </tr> <tr> <td>Polymyxin B 10,000 units as polymyxin B sulfate.....</td> <td>First aid antibiotic</td> </tr> <tr> <td>Pramoxine HCl 10 mg.....</td> <td>External analgesic</td> </tr> </table> <p>Uses First aid to help prevent infection and for temporary relief of pain or discomfort in minor:</p> <ul style="list-style-type: none"> ■ cuts ■ scrapes ■ burns <p>Warnings For external use only.</p> <p>Don't use</p> <ul style="list-style-type: none"> ■ if you are allergic to any of the ingredients ■ in the eyes ■ over large areas of the body <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ deep or puncture wounds ■ animal bites ■ serious burns 	Active ingredients (in each gram)	Purpose	Bacitracin zinc 500 units.....	First aid antibiotic	Neomycin sulfate equivalent to 35 mg of neomycin base....	First aid antibiotic	Polymyxin B 10,000 units as polymyxin B sulfate.....	First aid antibiotic	Pramoxine HCl 10 mg.....	External analgesic	<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ you need to use longer than 1 week ■ condition persists or gets worse ■ symptoms persist for more than 1 week, or clear up and occur again within a few days ■ rash or other allergic reaction develops <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p> <p>Directions</p> <ul style="list-style-type: none"> ■ adults and children 2 years of age or older: <ul style="list-style-type: none"> ■ clean the affected area ■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily ■ maybe covered with a sterile bandage ■ children under 2 years of age ask a doctor <p>Other information</p> <ul style="list-style-type: none"> ■ store at 20-25°C (68-77°F) <p>Inactive ingredient white petrolatum</p> <p>Questions or comments? 1-877-932-7948</p>
Active ingredients (in each gram)	Purpose										
Bacitracin zinc 500 units.....	First aid antibiotic										
Neomycin sulfate equivalent to 35 mg of neomycin base....	First aid antibiotic										
Polymyxin B 10,000 units as polymyxin B sulfate.....	First aid antibiotic										
Pramoxine HCl 10 mg.....	External analgesic										

Code Area



EQUALINE FIRST AID ANTIBIOTIC			
bacitracin, neomycin, polymyxin b, pramoxine hcl ointment			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41163-040
Route of Administration	TOPICAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN	500 [USP'U] in 1 g	
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN	3.5 mg in 1 g	
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B	10000 [USP'U] in 1 g	
PRAMOXINE HYDROCHLORIDE (UNII: 88AYB867L5) (PRAMOXINE - UNII:068X84E056)	PRAMOXINE HYDROCHLORIDE	10 mg in 1 g	
Inactive Ingredients			
Ingredient Name	Strength		
PETROLATUM (UNII: 4T6H12BN9U)			

Product Characteristics

Color	YELLOW (Translucent)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41163-040-64	1 in 1 CARTON	04/02/2015	
1		28 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:41163-040-58	1 in 1 CARTON	04/02/2015	02/28/2021
2		14 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M004	04/02/2015	

Labeler - United Natural Foods, Inc. dba UNFI (943556183)

Revised: 10/2024

United Natural Foods, Inc. dba UNFI