

**METRONIDAZOLE- metronidazole cream**  
**Cosette Pharmaceuticals, Inc.**

-----  
**Metronidazole Topical Cream, 0.75%**


**FOR TOPICAL USE ONLY**

**(NOT FOR OPHTHALMIC USE)**

**Rx Only**

**DESCRIPTION**

Metronidazole Topical Cream contains metronidazole, USP, at a concentration of 7.5 mg per gram (0.75%) in an emollient cream consisting of emulsifying wax, sorbitol solution, glycerin, isopropyl palmitate, benzyl alcohol, lactic acid and/or sodium hydroxide to adjust pH, and purified water. Metronidazole is a member of the imidazole class of antibacterial agents and is classified therapeutically as an antiprotozoal and antibacterial agent. Chemically, metronidazole is 2-methyl-5-nitro-1 *H*-imidazole-1-ethanol. The molecular formula is C<sub>6</sub>H<sub>9</sub>N<sub>3</sub>O<sub>3</sub> and molecular weight is 171.16. Metronidazole is represented by the following structural formula:



**CLINICAL PHARMACOLOGY**

The mechanisms by which metronidazole acts in the treatment of rosacea are unknown, but appear to include an anti-inflammatory effect.

**INDICATIONS AND USAGE**

Metronidazole Topical Cream is indicated for topical application in the treatment of inflammatory papules and pustules of rosacea.

**CONTRAINDICATIONS**

Metronidazole Topical Cream is contraindicated in individuals with a history of hypersensitivity to metronidazole, or other ingredients of the formulation.

## **PRECAUTIONS**

### **General:**

Topical metronidazole has been reported to cause tearing of the eyes. Therefore, contact with the eyes should be avoided. If a reaction suggesting local irritation occurs, patients should be directed to use the medication less frequently or discontinue use. Metronidazole is a nitroimidazole and should be used with care in patients with evidence of, or history of blood dyscrasia.

### **Information for patients:**

This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes.

### **Drug Interactions:**

Oral metronidazole has been reported to potentiate the anticoagulant effect of warfarin and coumarin anticoagulants, resulting in a prolongation of prothrombin time. The effect of topical metronidazole on prothrombin time is not known.

### **Carcinogenesis, mutagenesis, impairment of fertility :**

Metronidazole has shown evidence of carcinogenic activity in a number of studies involving chronic, oral administration in mice and rats but not in studies involving hamsters.

Metronidazole has shown evidence of mutagenic activity in several *in vitro* bacterial assay systems. In addition, a dose-response increase in the frequency of micronuclei was observed in mice after intraperitoneal injections and an increase in chromosome aberrations have been reported in patients with Crohn's disease who were treated with 200-1200 mg/day of metronidazole for 1 to 24 months. However, no excess chromosomal aberrations in circulating human lymphocytes have been observed in patients treated for 8 months.

### **Pregnancy:**

*Teratogenic effects: Pregnancy category B*

There are no adequate and well-controlled studies with the use of Metronidazole Topical Cream in pregnant women. Metronidazole crosses the placental barrier and enters the fetal circulation rapidly. No fetotoxicity was observed after oral metronidazole in rats or mice. However, because animal reproduction studies are not always predictive of human response and since oral metronidazole has been shown to be a carcinogen in some rodents, this drug should be used during pregnancy only if clearly needed.

### **Nursing mothers:**

After oral administration, metronidazole is secreted in breast milk in concentrations

similar to those found in the plasma. Even though blood levels are significantly lower with topically applied metronidazole than those achieved after oral administration of metronidazole, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

**Pediatric use:**

Safety and effectiveness in pediatric patients have not been established.

**ADVERSE REACTIONS**

In controlled clinical trials, the total incidence of adverse reactions associated with the use of Metronidazole Topical Cream was approximately 10%. Skin discomfort (burning and stinging) was the most frequently reported event followed by erythema, skin irritation, pruritus and worsening of rosacea. All individual events occurred in less than 3% of patients. The following additional adverse experiences have been reported with the topical use of metronidazole: dryness, transient redness, metallic taste, tingling or numbness of extremities and nausea.

**To report SUSPECTED ADVERSE REACTIONS, contact Cosette Pharmaceuticals, Inc. at 1-800-922-1038 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

**DOSAGE AND ADMINISTRATION**

Apply and rub in a thin layer of Metronidazole Topical Cream twice daily, morning and evening, to entire affected areas after washing.

Areas to be treated should be washed with a mild cleanser before application. Patients may use cosmetics after application of Metronidazole Topical Cream.

**HOW SUPPLIED**

Metronidazole Topical Cream, 0.75% is supplied in a 45 g tube **NDC0713-0633-37**.

**Storage conditions:**

Store at 20°-25°C (68°-77°F) [see USP Controlled Room Temperature]. Protect from Freezing.

**Distributed by:**

Cosette Pharmaceuticals, Inc.  
South Plainfield, NJ 07080

8-0633CPLNC2 VC7649  
Rev. 08/2022

**PRINCIPAL DISPLAY PANEL**

**NDC 0713-0633-37**

**Metronidazole Topical Cream 0.75%**


**45 g**

**Rx only**

**FOR TOPICAL USE ONLY.**

**NOT FOR OPHTHALMIC USE.**

**Cosette Pharmaceuticals, Inc.**



## METRONIDAZOLE

metronidazole cream

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0713-0633
Route of Administration	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
METRONIDAZOLE (UNII: 140QMO216E) (METRONIDAZOLE - UNII:140QMO216E)	METRONIDAZOLE	7.5 mg in 1 g

## Inactive Ingredients

Ingredient Name	Strength
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>ISOPROPYL PALMITATE</b> (UNII: 8CRQ2TH63M)	
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>LACTIC ACID, UNSPECIFIED FORM</b> (UNII: 33X04XA5AT)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>WATER</b> (UNII: 059QF0KO0R)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0713-0633-37	1 in 1 CARTON	03/13/2008	
1		45 g in 1 TUBE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077549	03/13/2008	

**Labeler** - Cosette Pharmaceuticals, Inc. (116918230)

**Registrant** - Cosette Pharmaceuticals, Inc. (116918230)

## Establishment

Name	Address	ID/FEI	Business Operations
Cosette Pharmaceuticals NC Laboratories, LLC		079419931	analysis(0713-0633) , label(0713-0633) , manufacture(0713-0633) , pack(0713-0633)

Revised: 11/2023

Cosette Pharmaceuticals, Inc.