

LIL DRUG STORE ASPIRIN 325- aspirin tablet
Lil Drug Store Products, Inc

Lil' Drug Store ® Aspirin 325

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID) ¹

¹ nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves:

- headache
- muscle pain
- menstrual pain
- toothache
- pain and fever of colds
- minor pain of arthritis

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- stomach bleeding warning applies to you
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - feel faint
 - have bloody or black stools
 - have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- new symptoms occur
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets

every 6 hours, not to exceed 12 tablets in 24 hours

- children under 12 years: ask a doctor

Other information

- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredient

corn starch

Questions or comments?

Call toll-free **1-877-507-6516 (M-F 8AM-4:30PM CST)**

PRINCIPAL DISPLAY PANEL - 325 mg Tablet Bottle Label

Regular Strength

Aspirin 325

325 mg, Pain Reliever/

Fever Reducer (NSAID)

100

Tablets

Lil'

Drug Store ®

Regular Strength
Aspirin 325
 325 mg, Pain Reliever/
 Fever Reducer
 (NSAID)

100
 Tablets

Actual Size

50844 REVO12224912 68406L-US-06-22

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)
 Aspirin 325 mg (NSAID)*
 *nonsteroidal anti-inflammatory drug

Purpose
 Pain reliever/fever reducer

Uses temporarily relieves:

- headache
- muscle pain
- toothache
- menstrual pain
- pain and fever of colds
- minor pain of arthritis

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include:
 ■ hives ■ stomach ■ facial swelling
 ■ asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

3 66715 68406 6

PEEL HERE FOR MORE DRUG FACTS

Drug Facts (continued)

The chance is higher if you

- have had stomach ulcers or bleeding problems
- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have asthma
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - ringing in the ears or a loss of hearing occurs

STOP PEELING

Drug Facts (continued)

- pain gets worse or lasts more than 10 days
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over, take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredient corn starch

Questions or comments?
 Call toll-free 1-877-507-6516
 (M-F 8AM-4:30PM CST)

Product manufactured for: LIL Drug Store Products, Inc.
 6300 Eastern Lane SW, Coon Rapids, IA 52404

LIL DRUG STORE ASPIRIN 325			
aspirin tablet			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-6840
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
Inactive Ingredients			
	Ingredient Name	Strength	

STARCH, CORN (UNII: O8232NY3SJ)

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	ASPIRIN;44249
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-6840-6	100 in 1 BOTTLE; Type 0: Not a Combination Product	05/04/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/04/2016	

Labeler - Lil Drug Store Products, Inc (093103646)

Revised: 12/2023

Lil Drug Store Products, Inc