

DR. GREENGOODS LICE KILLER LICE TREATMENT- lecithin spray
Applied Science Labs, Inc.

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

Dr. Greengoods Lice Killer LICE TREATMENT

Drug Facts

Active ingredient

Lecithin 1x HPUS (1.6%)

Purpose

Pediculicide

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Use treats head lice

Warnings

For external use only

Do not use • near eyes • inside nose, mouth or vagina • on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.

When using this product

- keep eyes tightly closed and protect eyes with a washcloth or towel
- if product gets in eyes, flush with water right away for 15 minutes
- do not rub eyes
- scalp itching and redness may occur.
- do not ingest as may cause gastrointestinal irritation with nausea and diarrhea.
- do not freeze
- do not store at a temperature greater than 140° F

Stop use and ask a doctor if

- eye irritation occurs
- skin or scalp irritation continues or infection occurs

Directions

- **Important: Read warnings before use. Adults and children 2 years and over:**
- **Inspect** • check each household member with a magnifying glass in bright light for lice/nits (eggs) •

look for tiny nits near scalp beginning at back of neck and behind ears • examine small sections of hair at a time • unlike dandruff which moves when touched, nits stick to the hair • if either lice or nits are found, treat with this product.

- while holding the spray bottle in one hand, back stroke your child's hair so as to expose their scalp/skin.
- with these areas exposed, mist the Dr. Greengoods'Lice Killer. No more than a mist is necessary.
- **Treat** • apply thoroughly to dry hair or other affected area. For head lice, first apply behind ears and to back of neck. • allow product to remain for 10 minutes, but no longer • wash area thoroughly with warm water and soap or shampoo • for head lice, towel dry hair and comb out tangles.
- repeat the process until you have treated your child's entire head.
- as you prepare to treat hair toward the front of your child's face, ask them to close their eyes and gently shield the eyes against excess mist with one hand and mist spray with the other.
- hold nozzle 10 to 12 inches from treated area.
- be sure not to spray their eyes directly. When wet use nit comb.
- **Remove lice and their eggs (nits)** • use a fine-tooth or special lice/nit comb. Remove any remaining nits by hand (using a throw-away glove). • hair should remain slightly damp while removing nits • if hair dries during combing, dampen slightly with spray • for head lice, after spraying the entire head, part hair into sections. Do one section at a time starting on top of head. Longer hair may take 1 to 2 hours. • lift a 1-to 2-inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp • pin back each strand of hair after combing • clean comb often. Wipe nits away with tissue and discard in a plastic bag. Seal bag and discard to prevent lice from coming back.
- after combing, thoroughly recheck for lice/nits. Repeat combing if necessary
- check daily for any lice/nits that you missed". • do not soak. Mist only.
- a second treatment must be done in 7 to 10 days to kill any newly hatched lice.
- if infestation continues, see a doctor for other treatments
- **children under 2 years:** ask a doctor

Inactive ingredients

Distilled water and Sodium Benzoate (as preservative)

Other information

Head lice • lay small white eggs (nits) on hair shaft close to scalp • nits are most easily found on back of neck or behind ears • disinfect hats, hair ribbons, scarves, coats, towels, and bed linens by machine washing in hot water, (above 54°C (130°F)), then using hottest dryer cycle for at least 20 minutes • items that cannot be washed (bedspreads, blankets, pillows, stuffed toys, etc.) should be dry-cleaned or sealed in a plastic bag for 4 weeks, then removed outdoors and shaken out very hard before using again • items that cannot be washed, dry-cleaned, or stored may be sprayed with Dr Greengood's Habitat product • soak all combs and brushes in hot water, (above 54°C (130°F)) for at least 10 minutes • vacuum all carpets, mattresses, upholstered furniture, and car seats that may have been used by affected people.

- **Storage:** Store at room temperature. Avoid exposure to heat, freezing and direct sunlight.
- **Disposal:** Place in trash or offer for recycling if available.

Questions?

Call 1-619-825-2121

Mon-Fri 8 AM to 5 PM

100% NON-TOXIC

Kills Lice when used according to label directions

This product is nontoxic to all mammals and does not harm the environment

Caution: Not for food use

Manufactured by:

Applied Science Labs,

Melissa, TX 75454

USA 619-825-2121

Packaging



**Dr Greengood's
Lice Killer**

LICE TREATMENT SPRAY

— HOMEOPATHIC —

Kills Lice when used according
to label directions

This product is nontoxic to all mammals
and does not harm the environment

Caution: Not for food use

**Net Contents:
20 fl. oz. (591.47 mL)**

100% NONTOXIC

Drug Facts		Drug Facts (continued)	
Active ingredient Lecithin 1x HPUS (1.6%)		Purpose Pediculicide	
Use treats head lice			
Warnings For external use only			
Do not use ■ near eyes ■ inside nose, mouth or vagina ■ on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.			
When using this product			
■ keep eyes tightly closed and protect eyes with a washcloth or towel.			
■ if product gets in eyes, flush with water right away for 15 minutes.			
■ do not rub eyes.			
■ scalp itching or redness may occur.			
■ do not ingest as may cause gastrointestinal irritation with nausea and diarrhea.			
■ do not freeze.			
■ do not store at a temperature greater than 140°F.			
Stop use and ask a doctor if			
■ eye irritation occurs ■ skin or scalp irritation continues or infection occurs.			
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.			
Directions			
■ Important: Read warnings before use. Adults and children 2 years and over:			
■ Inspect ■ check each household member with a magnifying glass in bright light for lice/nits (eggs) ■ look for tiny nits near scalp, beginning at back of neck and behind ears ■ examine small sections of hair at a time ■ unlike dandruff which moves when touched, nits stick to the hair ■ if either lice or nits are found, treat with this product.			
■ while holding the spray bottle in one hand, back stroke your child's hair so as to expose their scalp/skin.			
■ with these areas exposed, mist the Dr Greengood's Lice Killer. No more than a mist is necessary.			
■ Treat ■ apply thoroughly to dry hair or other affected area. For head lice, first apply behind ears and to back of neck. ■ allow product to remain for 10 minutes, but no longer ■ wash area thoroughly with warm water and soap or shampoo ■ for head lice, towel dry hair and comb out tangles.			
■ repeat the process until you have treated your child's entire head.			
■ as you prepare to treat hair toward the front of your child's face, ask them to close their eyes and gently shield the eyes against excess mist with one hand and mist spray with the other.			
■ hold nozzle 10 to 12 inches from treated area.			
■ be sure not to spray their eyes directly. When wet use nit comb.			
■ Remove lice and their eggs (nits) ■ use a fine-tooth or special lice/nit comb. Remove any remaining nits by hand (using a throw-away glove). ■ hair should remain slightly damp while removing nits. ■ if hair dries during combing, dampen slightly with spray ■ for head lice, after spraying the entire head, part hair into sections. Do one section at a time starting on top of head. Longer hair may take 1 to 2 hours. ■ lift a 1- to 2-inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp. ■ pin back each strand of hair after combing ■ clean comb often. Wipe nits away with tissue and discard in a plastic bag. Seal bag and discard to prevent lice from coming back.			
■ after combing, thoroughly recheck for lice/nits. Repeat combing if necessary.			
■ check daily for any lice/nits that you missed. ■ do not soak. Mist only.			
■ a second treatment must be done in 7 to 10 days to kill any newly hatched lice.			
■ if infestation continues, see a doctor for other treatments			
■ children under 2 years: ask a doctor.			
Other information		Head lice ■ lay small white eggs (nits) on hair shaft close to scalp ■ nits are most easily found on back of neck or behind ears ■ disinfect hats, hair ribbons, scarves, coats, towels, and bed linens by machine washing in hot water, (above 54°C (130°F)), then using hottest dryer cycle for at least 20 minutes ■ items that cannot be washed (bedspreads, blankets, pillows, stuffed toys, etc.) should be dry-cleaned or sealed in a plastic bag for 4 weeks, then removed outdoors and shaken out very hard before using again ■ items that cannot be washed, dry-cleaned, or stored may be sprayed with Dr Greengood's Habitat product ■ soak all combs and brushes in hot water, (above 54°C (130°F)) for at least 10 minutes ■ vacuum all carpets, mattresses, upholstered furniture, and car seats that may have been used by affected people.	
Inactive ingredients		Distilled water and Sodium Benzoate (as preservative)	
Questions?		Call 1-619-825-2121 Mon-Fri 8 AM to 5 PM	
Manufactured by:		Applied Science Labs, TX 75454 USA 619-825-2121 Lot no.: 031816 D3	

DR. GREENGOODS LICE KILLER LICE TREATMENT

lecithin spray

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70606-107
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
EGG PHOSPHOLIPIDS (UNII: 1Z74184RGV) (EGG PHOSPHOLIPIDS - UNII:1Z74184RGV)	EGG PHOSPHOLIPIDS	1 [hp_X] in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM BENZOATE (UNII: OJ245FE5EU)	
WATER (UNII: 059QF0K00R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70606-107-88	591.47 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/07/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved homeopathic		07/07/2016	

Labeler - Applied Science Labs, Inc. (035483180)**Establishment**

Name	Address	ID/FEI	Business Operations
Applied Science Labs, Inc.		035483180	manufacture(70606-107)

Revised: 4/2016

Applied Science Labs, Inc.