

PROSTIN VR PEDIATRIC- alprostadil injection, solution
Pharmacia & Upjohn Company LLC

Prostin VR Pediatric®
alprostadil injection, USP

500 micrograms per mL

WARNING

Apnea is experienced by about 10 to 12% of neonates with congenital heart defects treated with PROSTIN VR PEDIATRIC Sterile Solution. Apnea is most often seen in neonates weighing less than 2 kg at birth and usually appears during the first hour of drug infusion. Therefore, respiratory status should be monitored throughout treatment, and PROSTIN VR PEDIATRIC should be used where ventilatory assistance is immediately available.


DESCRIPTION

PROSTIN VR PEDIATRIC Sterile Solution for intravascular infusion contains 500 micrograms alprostadil, more commonly known as prostaglandin E₁, in 1.0 mL dehydrated alcohol.

The chemical name for alprostadil is (11 α ,13E,15S)-11,15 dihydroxy-9-oxo-prost-13-en-1-oic acid, and the molecular weight is 354.49.

Alprostadil is a white to off-white crystalline powder with a melting point between 110° and 116°C. Its solubility at 35°C is 8000 micrograms per 100 mL double distilled water.

Structural Formula



CLINICAL PHARMACOLOGY

Alprostadil (prostaglandin E₁) is one of a family of naturally occurring acidic lipids with various pharmacologic effects. Vasodilation, inhibition of platelet aggregation, and stimulation of intestinal and uterine smooth muscle are among the most notable of these effects. Intravenous doses of 1 to 10 micrograms of alprostadil per kilogram of body weight lower the blood pressure in mammals by decreasing peripheral resistance. Reflex increases in cardiac output and rate accompany the reduction in blood pressure.

Smooth muscle of the ductus arteriosus is especially sensitive to alprostadil, and strips

of lamb ductus markedly relax in the presence of the drug. In addition, administration of alprostadil reopened the closing ductus of new-born rats, rabbits, and lambs. These observations led to the investigation of alprostadil in infants who had congenital defects which restricted the pulmonary or systemic blood flow and who depended on a patent ductus arteriosus for adequate blood oxygenation and lower body perfusion.

In infants with restricted pulmonary blood flow, about 50% responded to alprostadil infusion with at least a 10 torr increase in blood pO_2 (mean increase about 14 torr and mean increase in oxygen saturation about 23%). In general, patients who responded best had low pretreatment blood pO_2 and were 4 days old or less.

In infants with restricted systemic blood flow, alprostadil often increased pH in those having acidosis, increased systemic blood pressure, and decreased the ratio of pulmonary artery pressure to aortic pressure.

Alprostadil must be infused continuously because it is very rapidly metabolized. As much as 80% of the circulating alprostadil may be metabolized in one pass through the lungs, primarily by β - and ω - oxidation. The metabolites are excreted primarily by the kidney, and excretion is essentially complete within 24 hours after administration. No unchanged alprostadil has been found in the urine, and there is no evidence of tissue retention of alprostadil or its metabolites.

INDICATIONS AND USAGE

PROSTIN VR PEDIATRIC Sterile Solution is indicated for palliative, not definitive, therapy to temporarily maintain the patency of the ductus arteriosus until corrective or palliative surgery can be performed in neonates who have congenital heart defects and who depend upon the patent ductus for survival. Such congenital heart defects include pulmonary atresia, pulmonary stenosis, tricuspid atresia, tetralogy of Fallot, interruption of the aortic arch, coarctation of the aorta, or transposition of the great vessels with or without other defects.

In infants with restricted pulmonary blood flow, the increase in blood oxygenation is inversely proportional to pretreatment pO_2 values; that is, patients with low pO_2 values respond best, and patients with pO_2 values of 40 torr or more usually have little response.

PROSTIN VR PEDIATRIC should be administered only by trained personnel in facilities that provide pediatric intensive care.

CONTRAINDICATIONS

None.

WARNINGS

See WARNING box.

NOTE: PROSTIN VR PEDIATRIC Sterile Solution must be diluted before it is administered. See dilution instructions in DOSAGE AND ADMINISTRATION section.

The administration of PROSTIN VR PEDIATRIC to neonates may result in gastric outlet

obstruction secondary to antral hyperplasia. This effect appears to be related to duration of therapy and cumulative dose of the drug. Neonates receiving PROSTIN VR PEDIATRIC at recommended doses for more than 120 hours should be closely monitored for evidence of antral hyperplasia and gastric outlet obstruction.

PROSTIN VR PEDIATRIC should be infused for the shortest time and at the lowest dose that will produce the desired effects. The risks of long-term infusion of PROSTIN VR PEDIATRIC should be weighed against the possible benefits that critically ill infants may derive from its administration.

PRECAUTIONS

General Precautions

Cortical proliferation of the long bones, first observed in dogs, has also been observed in infants during long-term infusions of alprostadil. The cortical proliferation in infants regressed after withdrawal of the drug.

In infants treated with PROSTIN VR PEDIATRIC at the usual doses for 10 hours to 12 days and who died of causes unrelated to ductus structural weakness, tissue sections of the ductus and pulmonary arteries have shown intimal lacerations, a decrease in medial muscularity and disruption of the medial and internal elastic lamina. Localized and aneurysmal dilatations and vessel wall edema also were seen compared to a series of pathological specimens from infants not treated with PROSTIN VR PEDIATRIC. The incidence of such structural alterations has not been defined.

Because alprostadil inhibits platelet aggregation, use PROSTIN VR PEDIATRIC cautiously in neonates with bleeding tendencies.

PROSTIN VR PEDIATRIC should not be used in neonates with respiratory distress syndrome. A differential diagnosis should be made between respiratory distress syndrome (hyaline membrane disease) and cyanotic heart disease (restricted pulmonary blood flow). If full diagnostic facilities are not immediately available, cyanosis (pO_2 less than 40 torr) and restricted pulmonary blood flow apparent on an X-ray are appropriate indicators of congenital heart defects.

Necessary Monitoring

In all neonates, arterial pressure should be monitored intermittently by umbilical artery catheter, auscultation, or with a Doppler transducer. *Should arterial pressure fall significantly, decrease the rate of infusion immediately.*

In infants with restricted pulmonary blood flow, measure efficacy of PROSTIN VR PEDIATRIC by monitoring improvement in blood oxygenation. In infants with restricted systemic blood flow, measure efficacy by monitoring improvement of systemic blood pressure and blood pH.

PROSTIN VR PEDIATRIC contains ethanol (79% w/v). Ethanol may cause CNS depressant effects such as somnolence and may alter the effects of other medicines.

Co-administration with medicines containing e.g. propylene glycol or ethanol may lead to accumulation of ethanol and induce adverse effects, particularly in young children with low or immature metabolic capacity.

Drug Interactions

No drug interactions have been reported between PROSTIN VR PEDIATRIC and the therapy standard in neonates with restricted pulmonary or systemic blood flow. Standard therapy includes antibiotics, such as penicillin and gentamicin; vasopressors, such as dopamine and isoproterenol; cardiac glycosides; and diuretics, such as furosemide.

Carcinogenesis, Mutagenesis, and Impairment of Fertility

Long-term carcinogenicity studies and fertility studies have not been done. The Ames and Alkaline Elution assays reveal no potential for mutagenesis.

ADVERSE REACTIONS

Central Nervous System

Apnea has been reported in about 12% of the neonates treated. (See WARNING box.) Other common adverse reactions reported have been fever in about 14% of the patients treated and seizures in about 4%. The following reactions have been reported in less than 1% of the patients: cerebral bleeding, hyperextension of the neck, hyperirritability, hypothermia, jitteriness, lethargy, and stiffness.

Cardiovascular System

The most common adverse reactions reported have been flushing in about 10% of patients (more common after intraarterial dosing), bradycardia in about 7%, hypotension in about 4%, tachycardia in about 3%, cardiac arrest in about 1%, and edema in about 1%. The following reactions have been reported in less than 1% of the patients: congestive heart failure, hyperemia, second degree heart block, shock, spasm of the right ventricle infundibulum, supraventricular tachycardia, and ventricular fibrillation.

Respiratory System

The following reactions have been reported in less than 1% of the patients: bradypnea, bronchial wheezing, hypercapnia, respiratory depression, respiratory distress, and tachypnea.

Gastrointestinal System

See WARNINGS

The most common adverse reaction reported has been diarrhea in about 2% of the patients. The following reactions have been reported in less than 1% of the patients: gastric regurgitation, and hyperbilirubinemia.

Hematologic System

The most common hematologic event reported has been disseminated intravascular coagulation in about 1% of the patients. The following events have been reported in less than 1% of the patients: anemia, bleeding, and thrombocytopenia.

Excretory System

Anuria and hematuria have been reported in less than 1% of the patients.

Skeletal System

Cortical proliferation of the long bones has been reported. See PRECAUTIONS.

Miscellaneous

Sepsis has been reported in about 2% of the patients. Peritonitis has been reported in less than 1% of the patients. Hypokalemia has been reported in about 1%, and hypoglycemia and hyperkalemia have been reported in less than 1% of the patients.

OVERDOSAGE

Apnea, bradycardia, pyrexia, hypotension, and flushing may be signs of drug overdose. If apnea or bradycardia occurs, discontinue the infusion, and provide appropriate medical treatment. Caution should be used in restarting the infusion. If pyrexia or hypotension occurs, reduce the infusion rate until these symptoms subside. Flushing is usually a result of incorrect intraarterial catheter placement, and the catheter should be repositioned.

DOSAGE AND ADMINISTRATION

The preferred route of administration for PROSTIN VR PEDIATRIC Sterile Solution is continuous intravenous infusion into a large vein. Alternatively, PROSTIN VR PEDIATRIC may be administered through an umbilical artery catheter placed at the ductal opening. Increases in blood pO₂ (torr) have been the same in neonates who received the drug by either route of administration.

Begin infusion with 0.05 to 0.1 micrograms alprostadil per kilogram of body weight per minute. A starting dose of 0.1 micrograms per kilogram of body weight per minute is the recommended starting dose based on clinical studies; however, adequate clinical response has been reported using a starting dose of 0.05 micrograms per kilogram of body weight per minute. After a therapeutic response is achieved (increased pO₂ in infants with restricted pulmonary blood flow or increased systemic blood pressure and blood pH in infants with restricted systemic blood flow), reduce the infusion rate to provide the lowest possible dosage that maintains the response. This may be accomplished by reducing the dosage from 0.1 to 0.05 to 0.025 to 0.01 micrograms per kilogram of body weight per minute. If response to 0.05 micrograms per kilogram of body weight per minute is inadequate, dosage can be increased up to 0.4 micrograms per kilogram of body weight per minute although, in general, higher infusion rates do not produce greater effects.

Dilution Instructions

To prepare infusion solutions, dilute 1 mL of PROSTIN VR PEDIATRIC Sterile Solution with Sodium Chloride Injection USP or Dextrose Injection USP. Undiluted PROSTIN VR PEDIATRIC Sterile Solution may interact with the plastic sidewalls of volumetric infusion chambers causing a change in the appearance of the chamber and creating a hazy solution. Should this occur, the solution and the volumetric infusion chamber should be

replaced.

When using a volumetric infusion chamber, the appropriate amount of intravenous infusion solution should be added to the chamber first. The undiluted PROSTIN VR PEDIATRIC Sterile Solution should then be added to the intravenous infusion solution, avoiding direct contact of the undiluted solution with the walls of the volumetric infusion chamber.

Dilute to volumes appropriate for the pump delivery system available. Prepare fresh infusion solutions every 24 hours. *Discard any solution more than 24 hours old.*

Sample Dilutions and Infusion Rates to Provide a Dosage of 0.1 Micrograms per Kilogram of Body Weight per Minute

Add 1 ampoule (500 micrograms) alprostadil to:	Approximate Concentration of resulting solution (micrograms/mL)	Infusion rate (mL/min per kg of body weight)
250 mL	2	0.05
100 mL	5	0.02
50 mL	10	0.01
25 mL	20	0.005

Example: To provide 0.1 micrograms/kilogram of body weight per minute to an infant weighing 2.8 kilograms using a solution of 1 ampoule PROSTIN VR PEDIATRIC in 100 mL of saline or dextrose: $\text{INFUSION RATE} = 0.02 \text{ mL/min per kg} \times 2.8 \text{ kg} = 0.056 \text{ mL/min}$ or 3.36 mL/hr.

HOW SUPPLIED

PROSTIN VR PEDIATRIC Sterile Solution is available in a package of 5 ×1 mL ampoules (NDC 0009-3169-06). Each mL contains 500 micrograms alprostadil in dehydrated alcohol.

Store PROSTIN VR PEDIATRIC Sterile Solution in a refrigerator at 2° to 8°C (36° to 46°F).

Rx only



Distributed by
Pfizer Labs
Division of Pfizer Inc.
New York, NY 10001

LAB-0013-5.0
Revised December 2025

PRINCIPAL DISPLAY PANEL - 1 mL Ampoule Label

1 mL
NDC 0009-3169-01

Prostin VR
Pediatric®
(alprostadil) injection, USP

500 mcg/mL

For Intravenous Use Only.
Dilute Before Use.

Refrigerate at 2° to 8°C (36° to 46°F).

Distributed by Pharmacia & Upjohn Co
Division of Pfizer Inc, NY, NY 10017

Rx only

PAA135258

LOT

EXP

1 mL

NDC 0009-3169-01

Prostin VR Pediatric[®]

(alprostadil) injection, USP

500 mcg/mL

For Intravenous Use Only.
Dilute Before Use.

Refrigerate at 2° to 8°C (36° to 46°F).

Distributed by Pharmacia & Upjohn Co
Division of Pfizer Inc, NY, NY 10017

Rx only

FPO GS1 Data Bar Limited (RSS) - 10 mil

P A A 1 3 5 2 5 8

LOT
EXP

IMPRINT AREA
reads this way
NO VARNISH
NO COPY
NO INK

PRINCIPAL DISPLAY PANEL - 1 mL Ampoule Carton

NDC 0009-3169-06

5—1 mL Ampoules

Prostin VR

Pediatric[®]

(alprostadil) injection, USP

500 mcg/mL

For Intravenous Use Only.

Dilute Before Use.

Pfizer Injectables

Rx only

NDC 0009-3169-06

5—1 mL Ampoules

Prostin VR Pediatric[®] (alprostadil) injection, USP 500 mcg/mL

For Intravenous Use Only.
Dilute Before Use.



Rx only



Distributed by
Pharmacia & Upjohn Co
Division of Pfizer Inc
NY, NY 10017

IMPRINT AREA
READS THIS WAY
NO VARNISH
NO INK
NO COPY

LOT
EXP
NS

PAA135259

GTIN: 003 0009 3169 06 7

Store in a
refrigerator

5—1 mL Ampoules

Prostin VR Pediatric[®]

(alprostadil) injection, USP

500 mcg/mL

For Intravenous Use Only.
Dilute Before Use.



between
2° to 8°C
(36° to 46°F).

Discard any
solution more
than 24 hours old.

**DOSAGE AND
USE:** See
accompanying
prescribing
information.

Each mL contains:
500 micrograms
alprostadil in
dehydrated
alcohol.

PROSTIN VR PEDIATRIC

alprostadil injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0009-3169
Route of Administration	INTRAVENOUS, INTRAVASCULAR		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALPROSTADIL (UNII: F5TD010360) (ALPROSTADIL - UNII:F5TD010360)	ALPROSTADIL	500 ug in 1 mL

Inactive Ingredients

Ingredient Name		Strength		
ALCOHOL (UNII: 3K9958V90M)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0009-3169-06	5 in 1 PACKAGE	10/16/1981	
1	NDC:0009-3169-01	1 mL in 1 AMPULE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
NDA	NDA018484	10/16/1981		

Labeler - Pharmacia & Upjohn Company LLC (618054084)

Establishment			
Name	Address	ID/FEI	Business Operations
Pfizer Manufacturing Belgium NV		370156507	ANALYSIS(0009-3169) , LABEL(0009-3169) , MANUFACTURE(0009-3169) , PACK(0009-3169)

Establishment			
Name	Address	ID/FEI	Business Operations
Pharmacia & Upjohn Company LLC		618054084	ANALYSIS(0009-3169) , API MANUFACTURE(0009-3169) , PACK(0009-3169) , LABEL(0009-3169)

Revised: 2/2026

Pharmacia & Upjohn Company LLC