

**TRANDOLAPRIL- trandolapril tablet**  
**Dispensing Solutions, Inc.**

-----  
**TRANDOLAPRIL TABLETS**

7325

7326

7327


Rx only

**WARNING: FETAL TOXICITY**

- **When pregnancy is detected, discontinue trandolapril as soon as possible.**
- **Drugs that act directly on the renin-angiotensin system can cause injury and death to the developing fetus (see WARNINGS, Fetal Toxicity).**

**DESCRIPTION**

Trandolapril is the ethyl ester prodrug of a nonsulfhydryl angiotensin converting enzyme (ACE) inhibitor, trandolaprilat. Trandolapril is chemically described as (2*S*,3*aR*,7*aS*)-1-[(*S*)-N-[(*S*)-1-carboxy-3-phenylpropyl]alanyl] hexahydro-2-indolinecarboxylic acid, 1-ethyl ester. Its structural formula is:



C<sub>24</sub>H<sub>34</sub>N<sub>2</sub>O<sub>5</sub>      M.W. 430.54

Melting range is 130° to 135°C

Trandolapril is a white or almost white powder that is soluble (> 100 mg/mL) in chloroform, dichloromethane, and methanol. Trandolapril tablets contain 1 mg, 2 mg, or 4 mg of trandolapril for oral administration. Each tablet also contains ferric oxide yellow, lactose anhydrous, povidone, pregelatinized starch, and sodium stearyl fumarate. In addition, 1 mg tablets contain FD&C Blue #2 Lake, FD&C Red #40 Lake, and FD&C Yellow #6 Lake and 4 mg tablets contain ferric oxide red.

**CLINICAL PHARMACOLOGY**

**Mechanism of Action**

Trandolapril is deesterified to the diacid metabolite, trandolaprilat, which is approximately eight times more active as an inhibitor of ACE activity. ACE is a peptidyl dipeptidase that catalyzes the conversion of angiotensin I to the vasoconstrictor, angiotensin II. Angiotensin II is a potent peripheral vasoconstrictor that also stimulates secretion of aldosterone by the adrenal cortex and provides negative feedback for renin secretion. The effect of trandolapril in hypertension appears to result primarily from the inhibition of circulating and tissue ACE activity thereby reducing angiotensin II

formation, decreasing vasoconstriction, decreasing aldosterone secretion, and increasing plasma renin. Decreased aldosterone secretion leads to diuresis, natriuresis, and a small increase of serum potassium. In controlled clinical trials, treatment with trandolapril alone resulted in mean increases in potassium of 0.1 mEq/L. (See **PRECAUTIONS**.)

ACE is identical to kininase II, an enzyme that degrades bradykinin, a potent peptide vasodilator; whether increased levels of bradykinin play a role in the therapeutic effect of trandolapril remains to be elucidated.

While the principal mechanism of antihypertensive effect is thought to be through the renin-angiotensin-aldosterone system, trandolapril exerts antihypertensive actions even in patients with low-renin hypertension. Trandolapril was an effective antihypertensive in all races studied. Both black patients (usually a predominantly low-renin group) and non-black patients responded to 2 to 4 mg of trandolapril.

## **Pharmacokinetics and Metabolism**

### Pharmacokinetics

Trandolapril's ACE-inhibiting activity is primarily due to its diacid metabolite, trandolaprilat. Cleavage of the ester group of trandolapril, primarily in the liver, is responsible for conversion. Absolute bioavailability after oral administration of trandolapril is about 10% as trandolapril and 70% as trandolaprilat. After oral trandolapril under fasting conditions, peak trandolapril levels occur at about one hour and peak trandolaprilat levels occur between 4 and 10 hours. The elimination half-life of trandolapril is about 6 hours. At steady state, the effective half-life of trandolaprilat is 22.5 hours. Like all ACE inhibitors, trandolaprilat also has a prolonged terminal elimination phase, involving a small fraction of administered drug, probably representing binding to plasma and tissue ACE. During multiple dosing of trandolapril, there is no significant accumulation of trandolaprilat. Food slows absorption of trandolapril, but does not affect AUC or  $C_{max}$  of trandolaprilat or  $C_{max}$  of trandolapril.

### Metabolism and Excretion

After oral administration of trandolapril, about 33% of parent drug and metabolites are recovered in urine, mostly as trandolaprilat, with about 66% in feces. The extent of the absorbed dose which is biliary excreted has not been determined. Plasma concentrations ( $C_{max}$  and AUC of trandolapril and  $C_{max}$  of trandolaprilat) are dose proportional over the 1 to 4 mg range, but the AUC of trandolaprilat is somewhat less than dose proportional. In addition to trandolaprilat, at least 7 other metabolites have been found, principally glucuronides or deesterification products.

Serum protein binding of trandolapril is about 80%, and is independent of concentration. Binding of trandolaprilat is concentration-dependent, varying from 65% at 1000 ng/mL to 94% at 0.1 ng/mL, indicating saturation of binding with increasing concentration.

The volume of distribution of trandolapril is about 18 liters. Total plasma clearances of trandolapril and trandolaprilat after approximately 2 mg IV doses are about 52 liters/hour and 7 liters/hour respectively. Renal clearance of trandolaprilat varies from 1 to 4 liters/hour, depending on dose.

## **Special Populations**

### Pediatric

Trandolapril pharmacokinetics have not been evaluated in patients < 18 years of age.

### Geriatric and Gender

Trandolapril pharmacokinetics have been investigated in the elderly (> 65 years) and in both genders. The plasma concentration of trandolapril is increased in elderly hypertensive patients, but the plasma concentration of trandolaprilat and inhibition of ACE activity are similar in elderly and young hypertensive patients. The pharmacokinetics of trandolapril and trandolaprilat and inhibition of ACE

activity are similar in male and female elderly hypertensive patients.

## Race

Pharmacokinetic differences have not been evaluated in different races.

## Renal Insufficiency

Compared to normal subjects, the plasma concentrations of trandolapril and trandolaprilat are approximately 2 fold greater and renal clearance is reduced by about 85% in patients with creatinine clearance below 30 mL/min and in patients on hemodialysis. Dosage adjustment is recommended in renally impaired patients. (See **DOSAGE AND ADMINISTRATION**.)

## Hepatic Insufficiency

Following oral administration in patients with mild to moderate alcoholic cirrhosis, plasma concentrations of trandolapril and trandolaprilat were, respectively, 9 fold and 2 fold greater than in normal subjects, but inhibition of ACE activity was not affected. Lower doses should be considered in patients with hepatic insufficiency. (See **DOSAGE AND ADMINISTRATION**.)

## Drug Interactions

Trandolapril did not affect the plasma concentration (pre-dose and 2 hours post-dose) of oral digoxin (0.25 mg). Coadministration of trandolapril and cimetidine led to an increase of about 44% in  $C_{max}$  for trandolapril, but no difference in the pharmacokinetics of trandolaprilat or in ACE inhibition. Coadministration of trandolapril and furosemide led to an increase of about 25% in the renal clearance of trandolaprilat, but no effect was seen on the pharmacokinetics of furosemide or trandolaprilat or on ACE inhibition.

## Pharmacodynamics and Clinical Effects

A single 2 mg dose of trandolapril produces 70 to 85% inhibition of plasma ACE activity at 4 hours with about 10% decline at 24 hours and about half the effect manifest at 8 days. Maximum ACE inhibition is achieved with a plasma trandolaprilat concentration of 2 ng/mL. ACE inhibition is a function of trandolaprilat concentration, not trandolapril concentration. The effect of trandolapril on exogenous angiotensin I was not measured.

## Hypertension

Four placebo-controlled dose response studies were conducted using once-daily oral dosing of trandolapril in doses from 0.25 to 16 mg per day in 827 black and non-black patients with mild to moderate hypertension. The minimal effective once-daily dose was 1 mg in non-black patients and 2 mg in black patients. Further decreases in trough supine diastolic blood pressure were obtained in non-black patients with higher doses, and no further response was seen with doses above 4 mg (up to 16 mg). The antihypertensive effect diminished somewhat at the end of the dosing interval, but trough/peak ratios are well above 50% for all effective doses. There was a slightly greater effect on the diastolic pressure, but no difference on systolic pressure with b.i.d. dosing. During chronic therapy, the maximum reduction in blood pressure with any dose is achieved within one week. Following 6 weeks of monotherapy in placebo-controlled trials in patients with mild to moderate hypertension, once-daily doses of 2 to 4 mg lowered supine or standing systolic/diastolic blood pressure 24 hours after dosing by an average 7 to 10/4 to 5 mmHg below placebo responses in non-black patients. Once-daily doses of 2 to 4 mg lowered blood pressure 4 to 6/3 to 4 mmHg in black patients. Trough to peak ratios for effective doses ranged from 0.5 to 0.9. There were no differences in response between men and women, but responses were somewhat greater in patients under 60 than in patients over 60 years old. Abrupt withdrawal of trandolapril has not been associated with a rapid increase in blood pressure.

Administration of trandolapril to patients with mild to moderate hypertension results in a reduction of supine, sitting and standing blood pressure to about the same extent without compensatory tachycardia.

Symptomatic hypotension is infrequent, although it can occur in patients who are salt- and/or volume-depleted. (See **WARNINGS**.) Use of trandolapril in combination with thiazide diuretics gives a blood pressure lowering effect greater than that seen with either agent alone, and the additional effect of trandolapril is similar to the effect of monotherapy.

## **INDICATIONS AND USAGE**

### **Hypertension**

Trandolapril Tablets are indicated for the treatment of hypertension. They may be used alone or in combination with other antihypertensive medication such as hydrochlorothiazide.

## **CONTRAINDICATIONS**

Trandolapril tablets are contraindicated in patients who are hypersensitive to this product, in patients with hereditary/idiopathic angioedema, and in patients with a history of angioedema related to previous treatment with an ACE inhibitor.

## **WARNINGS**

### **Anaphylactoid and Possibly Related Reactions**

Presumably because angiotensin converting enzyme inhibitors affect the metabolism of eicosanoids and polypeptides, including endogenous bradykinin, patients receiving ACE inhibitors, including trandolapril, may be subject to a variety of adverse reactions, some of them serious.

Do not coadminister aliskiren with trandolapril in patients with diabetes (see **PRECAUTIONS, Drug Interactions**).

#### **Anaphylactoid Reactions During Desensitization**

Two patients undergoing desensitizing treatment with hymenoptera venom while receiving ACE inhibitors sustained life-threatening anaphylactoid reactions. In the same patients, these reactions did not occur when ACE inhibitors were temporarily withheld, but they reappeared when the ACE inhibitors were inadvertently readministered.

#### **Anaphylactoid Reactions During Membrane Exposure**

Anaphylactoid reactions have been reported in patients dialyzed with high-flux membranes and treated concomitantly with an ACE inhibitor. Anaphylactoid reactions have also been reported in patients undergoing low-density lipoprotein apheresis with dextran sulfate absorption.

### **Head and Neck Angioedema**

In controlled trials ACE inhibitors (for which adequate data are available) cause a higher rate of angioedema in black than in non-black patients. Angioedema of the face, extremities, lips, tongue, glottis, and larynx has been reported in patients treated with ACE inhibitors including trandolapril. Symptoms suggestive of angioedema or facial edema occurred in 0.13% of trandolapril-treated patients. Two of the four cases were life-threatening and resolved without treatment or with medication (corticosteroids). Angioedema associated with laryngeal edema can be fatal. If laryngeal stridor or angioedema of the face, tongue or glottis occurs, treatment with trandolapril should be discontinued immediately, the patient treated in accordance with accepted medical care and carefully observed until the swelling disappears. In instances where swelling is confined to the face and lips, the condition generally resolves without treatment; antihistamines may be useful in relieving symptoms. **Where there is involvement of the tongue, glottis, or larynx, likely to cause airway obstruction, emergency therapy, including but not limited to subcutaneous epinephrine solution 1:1,000 (0.3 to 0.5 mL)**

**should be promptly administered.** (See **PRECAUTIONS, Information for Patients** and **ADVERSERECTIONS.**)

### **Intestinal Angioedema**

Intestinal angioedema has been reported in patients treated with ACE inhibitors. These patients presented with abdominal pain (with or without nausea or vomiting); in some cases there was no prior history of facial angioedema and C-1 esterase levels were normal. The angioedema was diagnosed by procedures including abdominal CT scan or ultrasound, or at surgery, and symptoms resolved after stopping the ACE inhibitor. Intestinal angioedema should be included in the differential diagnosis of patients on ACE inhibitors presenting with abdominal pain.

### **Hypotension**

Trandolapril can cause symptomatic hypotension. Like other ACE inhibitors, trandolapril has only rarely been associated with symptomatic hypotension in uncomplicated hypertensive patients. Symptomatic hypotension is most likely to occur in patients who have been salt- or volume-depleted as a result of prolonged treatment with diuretics, dietary salt restriction, dialysis, diarrhea, or vomiting. Volume and/or salt depletion should be corrected before initiating treatment with trandolapril. (See **PRECAUTIONS, Drug Interactions**, and **ADVERSE REACTIONS.**) In controlled and uncontrolled studies, hypotension was reported as an adverse event in 0.6% of patients and led to discontinuations in 0.1% of patients.

In patients with excessive hypotension, which may be associated with oliguria or azotemia, and rarely, with acute renal failure and death, trandolapril therapy should be started at the recommended dose under close medical supervision. These patients should be followed closely during the first 2 weeks of treatment and, thereafter, whenever the dosage of trandolapril or diuretic is increased. (See **DOSAGE AND ADMINISTRATION.**) Care in avoiding hypotension should also be taken in patients with ischemic heart disease, aortic stenosis, or cerebrovascular disease.

If symptomatic hypotension occurs, the patient should be placed in the supine position and, if necessary, normal saline may be administered intravenously. A transient hypotensive response is not a contraindication to further doses; however, lower doses of trandolapril or reduced concomitant diuretic therapy should be considered.

### **Neutropenia/Agranulocytosis**

Another ACE inhibitor, captopril, has been shown to cause agranulocytosis and bone marrow depression rarely in patients with uncomplicated hypertension, but more frequently in patients with renal impairment, especially if they also have a collagen-vascular disease such as systemic lupus erythematosus or scleroderma. Available data from clinical trials of trandolapril are insufficient to show that trandolapril does not cause agranulocytosis at similar rates. As with other ACE inhibitors, periodic monitoring of white blood cell counts in patients with collagen-vascular disease and/or renal disease should be considered.

### **Hepatic Failure**

ACE inhibitors rarely have been associated with a syndrome of cholestatic jaundice, fulminant hepatic necrosis, and death. The mechanism of this syndrome is not understood. Patients receiving ACE inhibitors who develop jaundice should discontinue the ACE inhibitor and receive appropriate medical follow-up.

### **Fetal Toxicity**

Pregnancy Category D

Use of drugs that act on the renin-angiotensin system during the second and third trimesters of pregnancy reduces fetal renal function and increases fetal and neonatal morbidity and death. Resulting

oligohydramnios can be associated with fetal lung hypoplasia and skeletal deformations. Potential neonatal adverse effects include skull hypoplasia, anuria, hypotension, renal failure, and death. When pregnancy is detected, discontinue trandolapril as soon as possible. These adverse outcomes are usually associated with use of these drugs in the second and third trimester of pregnancy. Most epidemiologic studies examining fetal abnormalities after exposure to antihypertensive use in the first trimester have not distinguished drugs affecting the renin-angiotensin system from other antihypertensive agents. Appropriate management of maternal hypertension during pregnancy is important to optimize outcomes for both mother and fetus.

In the unusual case that there is no appropriate alternative to therapy with drugs affecting the renin-angiotensin system for a particular patient, apprise the mother of the potential risk to the fetus. Perform serial ultrasound examinations to assess the intra-amniotic environment. If oligohydramnios is observed, discontinue trandolapril, unless it is considered lifesaving for the mother. Fetal testing may be appropriate, based on the week of pregnancy. Patients and physicians should be aware, however, that oligohydramnios may not appear until after the fetus has sustained irreversible injury. Closely observe infants with histories of *in utero* exposure to trandolapril for hypotension, oliguria, and hyperkalemia (see **PRECAUTIONS, Pediatric Use**).

Doses of 0.8 mg/kg/day (9.4 mg/m<sup>2</sup>/day) in rabbits, 1000 mg/kg/day (7000 mg/m<sup>2</sup>/day) in rats, and 25 mg/kg/day (295 mg/m<sup>2</sup>/day) in cynomolgus monkeys did not produce teratogenic effects. These doses represent 10 and 3 times (rabbits), 1250 and 2564 times (rats), and 312 and 108 times (monkeys) the maximum projected human dose of 4 mg based on body-weight and body-surface-area, respectively assuming a 50 kg woman.

## PRECAUTIONS

### General

#### Impaired Renal Function

As a consequence of inhibiting the renin-angiotensin-aldosterone system, changes in renal function may be anticipated in susceptible individuals. In patients whose renal function may depend on the activity of the renin-angiotensin-aldosterone system, treatment with ACE inhibitors, including trandolapril, may be associated with oliguria and/or progressive azotemia and rarely with acute renal failure and/or death.

In hypertensive patients with unilateral or bilateral renal artery stenosis, increases in blood urea nitrogen and serum creatinine have been observed in some patients following ACE inhibitor therapy. These increases were almost always reversible upon discontinuation of the ACE inhibitor and/or diuretic therapy. In such patients, renal function should be monitored during the first few weeks of therapy.

Some hypertensive patients with no apparent preexisting renal vascular disease have developed increases in blood urea and serum creatinine, usually minor and transient, especially when ACE inhibitors have been given concomitantly with a diuretic. This is more likely to occur in patients with preexisting renal impairment. Dosage reduction and/or discontinuation of any diuretic and/or the ACE inhibitor may be required. **Evaluation of hypertensive patients should always include assessment of renal function.** (See **DOSAGE AND ADMINISTRATION**.)

#### Hyperkalemia and Potassium-Sparing Diuretics

In clinical trials, hyperkalemia (serum potassium > 6 mEq/L) occurred in approximately 0.4% of hypertensive patients receiving trandolapril. In most cases, elevated serum potassium levels were isolated values, which resolved despite continued therapy. None of these patients were discontinued from the trials because of hyperkalemia. Risk factors for the development of hyperkalemia include renal insufficiency, diabetes mellitus, and the concomitant use of potassium-sparing diuretics, potassium supplements, and/or potassium-containing salt substitutes, which should be used cautiously, if at all,

with trandolapril. (See **PRECAUTIONS, Drug Interactions.**)

### Cough

Presumably due to the inhibition of the degradation of endogenous bradykinin, persistent nonproductive cough has been reported with all ACE inhibitors, always resolving after discontinuation of therapy. ACE inhibitor-induced cough should be considered in the differential diagnosis of cough. In controlled trials of trandolapril, cough was present in 2% of trandolapril patients and 0% of patients given placebo. There was no evidence of a relationship to dose.

### Surgery/Anesthesia

In patients undergoing major surgery or during anesthesia with agents that produce hypotension, trandolapril will block angiotensin II formation secondary to compensatory renin release. If hypotension occurs and is considered to be due to this mechanism, it can be corrected by volume expansion.

## Information for Patients

### Angioedema

Angioedema, including laryngeal edema, may occur at any time during treatment with ACE inhibitors, including trandolapril. Patients should be so advised and told to report immediately any signs or symptoms suggesting angioedema (swelling of face, extremities, eyes, lips, tongue, difficulty in swallowing or breathing) and to stop taking the drug until they have consulted with their physician. (See **WARNINGS** and **ADVERSE REACTIONS.**)

### Symptomatic Hypotension

Patients should be cautioned that light-headedness can occur, especially during the first days of trandolapril therapy, and should be reported to a physician. If actual syncope occurs, patients should be told to stop taking the drug until they have consulted with their physician. (See **WARNINGS.**)

All patients should be cautioned that inadequate fluid intake, excessive perspiration, diarrhea, or vomiting, resulting in reduced fluid volume, may precipitate an excessive fall in blood pressure with the same consequences of light-headedness and possible syncope.

Patients planning to undergo any surgery and/or anesthesia should be told to inform their physician that they are taking an ACE inhibitor that has a long duration of action.

### Hyperkalemia

Patients should be told not to use potassium supplements or salt substitutes containing potassium without consulting their physician. (See **PRECAUTIONS.**)

### Neutropenia

Patients should be told to report promptly any indication of infection (e.g., sore throat, fever) which could be a sign of neutropenia.

### Pregnancy

Female patients of childbearing age should be told about the consequences of exposure to trandolapril during pregnancy. Discuss treatment options with women planning to become pregnant. Patients should be asked to report pregnancies to their physicians as soon as possible.

**NOTE:** As with many other drugs, certain advice to patients being treated with trandolapril is warranted. This information is intended to aid in the safe and effective use of this medication. It is not a disclosure of all possible adverse or intended effects.

## Drug Interactions

## Dual Blockade of the Renin-Angiotensin System (RAS)

Dual blockade of the RAS with angiotensin receptor blockers, ACE inhibitors, or aliskiren is associated with increased risks of hypotension, hyperkalemia, and changes in renal function (including acute renal failure) compared to monotherapy. Closely monitor blood pressure, renal function and electrolytes in patients on trandolapril and other agents that affect the RAS.

Do not coadminister aliskiren with trandolapril in patients with diabetes. Avoid use of aliskiren with trandolapril in patients with renal impairment (GFR < 60 mL/min).

## Concomitant Diuretic Therapy

As with other ACE inhibitors, patients on diuretics, especially those on recently instituted diuretic therapy, may experience an excessive reduction of blood pressure after initiation of therapy with trandolapril. The possibility of exacerbation of hypotensive effects with trandolapril may be minimized by either discontinuing the diuretic or cautiously increasing salt intake prior to initiation of treatment with trandolapril. If it is not possible to discontinue the diuretic, the starting dose of trandolapril should be reduced. (See **DOSAGE AND ADMINISTRATION**.)

## Agents Increasing Serum Potassium

Trandolapril can attenuate potassium loss caused by thiazide diuretics and increase serum potassium when used alone. Use of potassium-sparing diuretics (spironolactone, triamterene, or amiloride), potassium supplements, or potassium-containing salt substitutes concomitantly with ACE inhibitors can increase the risk of hyperkalemia. If concomitant use of such agents is indicated, they should be used with caution and with appropriate monitoring of serum potassium. (See **PRECAUTIONS**.)

## Antidiabetic Agents

Concomitant use of ACE inhibitors and antidiabetic medicines (insulin or oral hypoglycemic agents) may cause an increased blood glucose lowering effect with greater risk of hypoglycemia.

## Lithium

Increased serum lithium levels and symptoms of lithium toxicity have been reported in patients receiving concomitant lithium and ACE inhibitor therapy. These drugs should be coadministered with caution, and frequent monitoring of serum lithium levels is recommended. If a diuretic is also used, the risk of lithium toxicity may be increased.

## Non-Steroidal Anti-Inflammatory Agents Including Selective Cyclooxygenase-2 Inhibitors (COX-2 Inhibitors)

In patients who are elderly, volume-depleted (including those on diuretic therapy), or with compromised renal function, coadministration of NSAIDs, including selective COX-2 inhibitors, with ACE inhibitors, including trandolapril, may result in deterioration of renal function, including possible acute renal failure. These effects are usually reversible. Monitor renal function periodically in patients receiving trandolapril and NSAID therapy.

The antihypertensive effect of ACE inhibitors, including trandolapril may be attenuated by NSAIDs.

## Gold

Nitritoid reactions (symptoms include facial flushing, nausea, vomiting and hypotension) have been reported rarely in patients on therapy with injectable gold (sodium aurothiomalate) and concomitant ACE inhibitor therapy including trandolapril.

## Other

No clinically significant pharmacokinetic interaction has been found between trandolaprilat and food,



cimetidine, digoxin, or furosemide.

The anticoagulant effect of warfarin was not significantly changed by trandolapril.

The hypotensive effect of certain inhalation anesthetics may be enhanced by ACE inhibitors including trandolapril (see **PRECAUTIONS**, *Surgery/Anesthesia*).

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Long-term studies were conducted with oral trandolapril administered by gavage to mice (78 weeks) and rats (104 and 106 weeks). No evidence of carcinogenic potential was seen in mice dosed up to 25 mg/kg/day (85 mg/m<sup>2</sup>/day) or rats dosed up to 8 mg/kg/day (60 mg/m<sup>2</sup>/day). These doses are 313 and 32 times (mice), and 100 and 23 times (rats) the maximum recommended human daily dose (MRHDD) of 4 mg based on body-weight and body-surface-area, respectively assuming a 50 kg individual. The genotoxic potential of trandolapril was evaluated in the microbial mutagenicity (Ames) test, the point mutation and chromosome aberration assays in Chinese hamster V79 cells, and the micronucleus test in mice. There was no evidence of mutagenic or clastogenic potential in these *in vitro* and *in vivo* assays.

Reproduction studies in rats did not show any impairment of fertility at doses up to 100 mg/kg/day (710 mg/m<sup>2</sup>/day) of trandolapril, or 1250 and 260 times the MRHDD on the basis of body-weight and body-surface-area, respectively.

### **Nursing Mothers**

Radiolabeled trandolapril or its metabolites are secreted in rat milk. Trandolapril should not be administered to nursing mothers.

### **Geriatric Use**

In placebo-controlled studies of trandolapril, 31.1% of patients were 60 years and older, 20.1% were 65 years and older, and 2.3% were 75 years and older. No overall differences in effectiveness or safety were observed between these patients and younger patients. (Greater sensitivity of some older individual patients cannot be ruled out.)

### **Pediatric Use**

#### **Neonates With a History of In Utero Exposure to Trandolapril**

If oliguria or hypotension occurs, direct attention toward support of blood pressure and renal perfusion. Exchange transfusions or dialysis may be required as a means of reversing hypotension and/or substituting for disordered renal function.

The safety and effectiveness of trandolapril in pediatric patients have not been established.

## **ADVERSE REACTIONS**

The safety experience in U.S. placebo-controlled trials included 1069 hypertensive patients, of whom 832 received trandolapril. Nearly 200 hypertensive patients received trandolapril for over one year in open-label trials. In controlled trials, withdrawals for adverse events were 2.1% on placebo and 1.4% on trandolapril. Adverse events considered at least possibly related to treatment occurring in 1% of trandolapril-treated patients and more common on trandolapril than placebo, pooled for all doses, are shown below, together with the frequency of discontinuation of treatment because of these events.

### **ADVERSE EVENTS IN PLACEBO-CONTROLLED HYPERTENSION TRIALS**

	<b>Occurring at 1% or Greater</b>	<b>PLACEBO (N = 227)</b>
--	-----------------------------------	------------------------------

	<b>TRANDOLAPRIL (N = 832) % Incidence (% Discontinuation)</b>	<b>(N = 257) % Incidence (% Discontinuation)</b>
Cough	1.9 (0.1)	0.4 (0.4)
Dizziness	1.3 (0.2)	0.4 (0.4)
Diarrhea	1 (0)	0.4 (0)

Headache and fatigue were all seen in more than 1% of trandolapril-treated patients but were more frequently seen on placebo. Adverse events were not usually persistent or difficult to manage.

Clinical adverse experiences possibly or probably related or of uncertain relationship to therapy occurring in 0.3% to 1% (except as noted) of the patients treated with trandolapril (with or without concomitant calcium ion antagonist or diuretic) in controlled or uncontrolled trials (N = 1134) and less frequent, clinically significant events seen in clinical trials or postmarketing experience include (listed by body system):

#### General Body Function

Chest pain.

#### Cardiovascular

AV first degree block, bradycardia, edema, flushing, palpitations.

#### Central Nervous System

Drowsiness, insomnia, paresthesia, vertigo.

#### Dermatologic

Pruritus, rash, pemphigus.

#### Eye, Ear, Nose, Throat

Epistaxis, throat inflammation, upper respiratory tract infection.

#### Emotional, Mental, Sexual States

Anxiety, impotence, decreased libido.

#### Gastrointestinal

Abdominal distention, abdominal pain/cramps, constipation, dyspepsia, diarrhea, vomiting, nausea.

#### Hemopoietic

Decreased leukocytes, decreased neutrophils.

#### Metabolism and Endocrine

Increased liver enzymes including SGPT (ALT).

#### Musculoskeletal System

Extremity pain, muscle cramps, gout.

#### Pulmonary

Dyspnea.

## **Postmarketing**

The following adverse reactions were identified during post approval use of trandolapril. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

### General Body Function

Malaise, fever.

### Cardiovascular

Myocardial infarction, myocardial ischemia, angina pectoris, cardiac failure, ventricular tachycardia, tachycardia, transient ischemic attack, arrhythmia.

### Central Nervous System

Cerebral hemorrhage.

### Dermatologic

Alopecia, sweating, Stevens-Johnson syndrome and toxic epidermal necrolysis.

### Emotional, Mental, Sexual States

Hallucination, depression.

### Gastrointestinal

Dry mouth, pancreatitis, jaundice and hepatitis.

### Hemopoietic

Agranulocytosis, pancytopenia.

### Metabolism and Endocrine

Increased SGOT (AST).

### Pulmonary

Bronchitis.

### Renal and Urinary

Renal failure.

## **Clinical Laboratory Test Findings**

### Hematology

Thrombocytopenia.

### Serum Electrolytes

Hyponatremia.

### Creatinine and Blood Urea Nitrogen

Increases in creatinine levels occurred in 1.1% of patients receiving trandolapril alone and 7.3% of patients treated with trandolapril, a calcium ion antagonist and a diuretic. Increases in blood urea nitrogen levels occurred in 0.6% of patients receiving trandolapril alone and 1.4% of patients receiving trandolapril, a calcium ion antagonist, and a diuretic. None of these increases required discontinuation

of treatment. Increases in these laboratory values are more likely to occur in patients with renal insufficiency or those pretreated with a diuretic and, based on experience with other ACE inhibitors, would be expected to be especially likely in patients with renal artery stenosis. (See **PRECAUTIONS** and **WARNINGS**.)

#### Liver Function Tests

Occasional elevation of transaminases at the rate of 3X upper normals occurred in 0.8% of patients and persistent increase in bilirubin occurred in 0.2% of patients. Discontinuation for elevated liver enzymes occurred in 0.2% of patients.

#### Other

Another potentially important adverse experience, eosinophilic pneumonitis, has been attributed to other ACE inhibitors.

#### OVERDOSAGE

No data are available with respect to overdosage in humans. The oral LD<sub>50</sub> oftrandolapril in mice was 4875 mg/kg in males and 3990 mg/kg in females. In rats, an oral dose of 5000 mg/kg caused low mortality (1 male out of 5; 0 females). In dogs, an oral dose of 1000 mg/kg did not cause mortality and abnormal clinical signs were not observed. In humans the most likely clinical manifestation would be symptoms attributable to severe hypotension. Symptoms also expected with ACE inhibitors are hypotension, hyperkalemia, and renal failure.

Laboratory determinations of serum levels oftrandolapril and its metabolites are not widely available, and such determinations have, in any event, no established role in the management oftrandolapril overdose. No data are available to suggest that physiological maneuvers (e.g., maneuvers to change the pH of the urine) might accelerate elimination oftrandolapril and its metabolites. Trandolaprilat is removed by hemodialysis. Angiotensin II could presumably serve as a specific antagonist antidote in the setting oftrandolapril overdose, but angiotensin II is essentially unavailable outside of scattered research facilities. Because the hypotensive effect oftrandolapril is achieved through vasodilation and effective hypovolemia, it is reasonable to treattrandolapril overdose by infusion of normal saline solution.

#### DOSAGE AND ADMINISTRATION

##### Hypertension

The recommended initial dosage oftrandolapril tablets for patients not receiving a diuretic is 1 mg once daily in non-black patients and 2 mg in black patients. Dosage should be adjusted according to the blood pressure response. Generally, dosage adjustments should be made at intervals of at least 1 week. Most patients have required dosages of 2 to 4 mg once daily. There is little clinical experience with doses above 8 mg.

Patients inadequately treated with once-daily dosing at 4 mg may be treated with twice-daily dosing. If blood pressure is not adequately controlled withtrandolapril tablets monotherapy, a diuretic may be added.

In patients who are currently being treated with a diuretic, symptomatic hypotension occasionally can occur following the initial dose oftrandolapril tablets. To reduce the likelihood of hypotension, the diuretic should, if possible, be discontinued two to three days prior to beginning therapy withtrandolapril tablets. (See **WARNINGS**.) Then, if blood pressure is not controlled withtrandolapril tablets alone, diuretic therapy should be resumed. If the diuretic cannot be discontinued, an initial dose of 0.5 mgtrandolapril tablets should be used with careful medical supervision for several hours until blood pressure has stabilized. The dosage should subsequently be titrated (as described above) to the

optimal response. (See **WARNINGS** and **PRECAUTIONS, Drug Interactions.**)

Concomitant administration of trandolapril tablets with potassium supplements, potassium salt substitutes, or potassium sparing diuretics can lead to increases of serum potassium. (See **PRECAUTIONS.**)

### **Dosage Adjustment in Renal Impairment or Hepatic Cirrhosis**

For patients with a creatinine clearance < 30 mL/min. or with hepatic cirrhosis, the recommended starting dose, based on clinical and pharmacokinetic data, is 0.5 mg daily. Patients should subsequently have their dosage titrated (as described above) to the optimal response.

### **HOW SUPPLIED/STORAGE AND HANDLING**

Trandolapril Tablets are supplied as follows:

1 mg: mottled salmon, capsule-shaped tablet, scored on one side and debossed with “9” on one side of the score and with “3” on the other side. Debossed with “7325” on the other side of the tablet. Available in bottles of 100 tablets.

2 mg: mottled yellow, capsule-shaped tablet, debossed with “93” on one side and with “7326” on the other side of the tablet. Available in bottles of 100 tablets.

4 mg: mottled rose, capsule-shaped tablet, debossed with “93” on one side and with “7327” on the other side of the tablet. Available in bottles of 100 tablets.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense in a tight, light-resistant container as defined in the USP, with a child-resistant closure (as required).

**KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.**

Manufactured In Israel By:

**TEVA PHARMACEUTICAL IND. LTD.**

Jerusalem, 91010, Israel

Manufactured For:

**TEVA PHARMACEUTICALS USA**

Sellersville, PA 18960

And Relabeled By:

**Dispensing Solutions Inc.**

3000 West Warner Ave

Santa Ana, CA 92704

United States

### **PRINCIPAL DISPLAY PANEL**

**BULK SOURCE DATA**  
MFD. FOR: TEVA PHARMACEUTICALS USA  
SELLERSVILLE, PA 18960

**PRODUCT ID:**  
MOTTLED ROSE CAPSULE-SHAPED  
TABLET DEBOSSED 93 / 7327

**BULK SOURCE NDC: 00093-7327-01**  
MFR. LOT: XXXXXX  
PEDIGREE #: 21463462  
DISPENSE IN THIS  
TIGHT/LIGHT RESISTANT CONTAINER



**TAKE \_\_\_ TABLET(S)**  
**ORALLY \_\_\_ TIME(S) A DAY**  
**AS DIRECTED. DO NOT**  
**DISCONTINUE WITHOUT**  
**PHYSICIAN'S ADVICE. MAY**  
**CAUSE HEADACHE, DIZZINESS,**  
**FATIGUE OR COUGHING**



**TRANDOLAPRIL 4 mg**  
**XX TABLETS**  
**NDC 68258-6100-XX**  
**PRODUCT # 6100-X**

Rev. Date: 11/12

**EACH TABLET CONTAINS:**  
**TRANDOLAPRIL . . . . . 4 mg**

**WARNING: USE DURING PREGNANCY CAN CAUSE FETAL HARM.**  
**LOT#SAMPLE EXP: 00-00 Rx # 30158814**

**RX ONLY**

**WARNING: KEEP OUT OF CHILDREN'S REACH**  
**STORE AT 68° - 77° F. SEE USP.**

6100-X NDC 68258-6100-XX  
TRANDOLAPRIL 4 mg  
XX TABLETS  
LOT # SAMPLE EXP: 00-00  
MN 00093-7327-01 RX# 30158814

6100-X NDC 68258-6100-XX  
TRANDOLAPRIL 4 mg  
XX TABLETS  
LOT # SAMPLE EXP: 00-00  
MN 00093-7327-01 RX# 30158814

6100-X NDC 68258-6100-XX  
TRANDOLAPRIL 4 mg  
XX TABLETS  
LOT # SAMPLE EXP: 00-00  
MN 00093-7327-01 RX# 30158814



Packaged Exclusively By:  
**DISPENSING SOLUTIONS<sup>inc</sup>**  
Santa Ana, CA 92704

NDC 68258-6100-XX

NDC 68258-6100-30

## TRANDOLAPRIL

trandolapril tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:68258-6100(NDC:0093-7327)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TRANDOLAPRIL (UNII: 1T0N3G9CRC) (TRANDOLAPRILAT - UNII:RR6866VL0O)	TRANDOLAPRIL	4 mg

### Inactive Ingredients

Ingredient Name	Strength
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
POVIDONE K30 (UNII: U725QWY32X)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UI)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

### Product Characteristics

<b>Color</b>	RED (mottled rose)	<b>Score</b>	no score
<b>Shape</b>	OVAL (capsule-shaped)	<b>Size</b>	11mm
<b>Flavor</b>		<b>Imprint Code</b>	93;7327
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68258-6100-3	30 in 1 BOTTLE, PLASTIC		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077489	02/07/2007	

**Labeler** - Dispensing Solutions, Inc. (066070785)

**Registrant** - PSS World Medical, Inc. (101822682)

## Establishment

Name	Address	ID/FEI	Business Operations
Dispensing Solutions, Inc.		066070785	relabel(68258-6100) , repack(68258-6100)

Revised: 10/2013

Dispensing Solutions, Inc.