

TOPCARE NITE TIME COLD AND FLU RELIEF- acetaminophen, dextromethorphan hbr, doxylamine succinate solution

Topco Associates LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Topco Associates LLC. Nite Time Cold & Flu Relief Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed

by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 30 mL contains: sodium 39 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments?

1-888-423-0139

Package/Label Principal Display Panel

SEE NEW WARNINGS

COMPARE TO VICKS® NYQUIL® ACTIVE INGREDIENTS

MULTI-SYMPTOM RELIEF

Nite Time Cold & Flu Relief

OUR PHARMACISTS RECOMMEND

PAIN RELIEVER-FEVER REDUCER – ACETAMINOPHEN

COUGH SUPPRESSANT – DEXTROMETHORPHAN HBr

ANTIHISTAMINE – DOXYLAMINE SUCCINATE

Headache, Fever, Sore Throat, Minor Aches & Pains

Sneezing, Runny Nose

Cough

ALCOHOL 10%

12 FL OZ (355 mL)

MIXED BERRY FLAVOR

Drug Facts (continued)

- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks

Stop use and ask a doctor if

- redness or swelling is present
- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 30 mL contains: sodium 39 mg
- store at 20-25°C (68-77°F)

Inactive ingredients alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments? 1-888-423-0139

*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® NyQuil®.

DISTRIBUTED BY TOPCO ASSOCIATES LLC, ELK GROVE VILLAGE, IL 60007
©TOPCO PERA1017 QUESTIONS? 1-888-423-0139
topcare@topco.com www.topcarebrand.com

GLUTEN FREE

QUALITY GUARANTEED

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org



0 36800 36738 8

NDC 36800-872-40



SEE NEW WARNINGS COMPARE TO VICKS® NYQUIL® ACTIVE INGREDIENTS*

MULTI-SYMPTOM RELIEF

Nite Time

Cold & Flu Relief



PAIN RELIEVER-FEVER REDUCER - **ACETAMINOPHEN**
COUGH SUPPRESSANT - **DEXTROMETHORPHAN HBr**
ANTHISTAMINE - **DOXYLAMINE SUCCINATE**

- Headache, Fever, Sore Throat, Minor Aches & Pains
- Sneezing, Runny Nose
- Cough

ALCOHOL 10%



12 FL OZ (355 mL) MIXED BERRY FLAVOR

Drug Facts **DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING**

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine

Uses temporarily relieves common cold/flu symptoms:

- fever
- cough due to minor throat and bronchial irritation
- minor aches and pains
- runny nose and sneezing
- sore throat
- headache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen

3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have liver disease, glaucoma, cough that occurs with too much phlegm (mucus), a breathing problem such as emphysema or chronic bronchitis, trouble urinating due to an enlarged prostate gland, persistent or chronic cough as occurs with smoking, asthma, or emphysema, a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers, taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur

: 87240 88 F5

TOPCARE NITE TIME COLD AND FLU RELIEF
acetaminophen, dextromethorphan hbr, doxylamine succinate solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-872
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	

POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

WATER (UNII: 059QF0KO0R)

SACCHARIN SODIUM (UNII: SB8ZUX40TY)

SODIUM CITRATE (UNII: 1Q73Q2JULR)

Product Characteristics

Color	RED (clear dark red)	Score	
Shape		Size	
Flavor	BERRY (berry/fruity)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-872-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/17/2012	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/17/2012	

Labeler - Topco Associates LLC (006935977)

Revised: 12/2019

Topco Associates LLC