


NITROGLYCERIN- nitroglycerin tablet
Golden State Medical Supply, Inc.

Nitroglycerin Sublingual Tablets USP

DESCRIPTION

Nitroglycerin is a stabilized sublingual compressed nitroglycerin tablet that contains 0.3 mg, 0.4 mg, or 0.6 mg nitroglycerin USP; as well as calcium stearate powder, colloidal silicon dioxide, hydrogenated vegetable oil, lactose monohydrate, and pregelatinized starch.

Nitroglycerin, an organic nitrate, is a vasodilating agent. The chemical name for nitroglycerin is 1, 2, 3 propanetriol trinitrate and the chemical structure is:



Molecular weight: 227.09

CLINICAL PHARMACOLOGY

The principal pharmacological action of nitroglycerin is relaxation of vascular smooth muscle. Although venous effects predominate, nitroglycerin produces, in a dose-related manner, dilation of both arterial and venous beds. Dilation of postcapillary vessels, including large veins, promotes peripheral pooling of blood, decreases venous return to the heart, and reduces left ventricular end-diastolic pressure (preload). Nitroglycerin also produces arteriolar relaxation, thereby reducing peripheral vascular resistance and arterial pressure (afterload), and dilates large epicardial coronary arteries; however, the extent to which this latter effect contributes to the relief of exertional angina is unclear.

Therapeutic doses of nitroglycerin may reduce systolic, diastolic, and mean arterial blood pressure. Effective coronary perfusion pressure is usually maintained, but can be compromised if blood pressure falls excessively, or increased heart rate decreases diastolic filling time.

Elevated central venous and pulmonary capillary wedge pressures, and pulmonary and systemic vascular resistance are also reduced by nitroglycerin therapy. Heart rate is usually slightly increased, presumably due to a compensatory response to the fall in blood pressure. Cardiac index may be increased, decreased, or unchanged. Myocardial oxygen consumption or demand (as measured by the pressure-rate product, tension-time index, and stroke-work index) is decreased and a more favorable supply-demand ratio can be achieved. Patients with elevated left ventricular filling pressures and increased systemic vascular resistance in association with a depressed cardiac index are likely to experience an improvement in cardiac index. In contrast, when filling pressures and cardiac index are normal, cardiac index may be slightly reduced following nitroglycerin administration.

Mechanism of Action: Nitroglycerin forms free radical nitric oxide (NO) which activates guanylate

cyclase, resulting in an increase of guanosine 3'5' monophosphate (cyclic GMP) in smooth muscle and other tissues. These events lead to dephosphorylation of myosin light chains, which regulate the contractile state in smooth muscle, and result in vasodilatation.

Pharmacodynamics: Consistent with the symptomatic relief of angina, digital plethysmography indicates that onset of the vasodilatory effect occurs approximately 1 to 3 minutes after sublingual nitroglycerin administration and reaches a maximum by 5 minutes postdose. Effects persist for at least 25 minutes following nitroglycerin administration.

Pharmacokinetics and Drug Metabolism Absorption: Nitroglycerin is rapidly absorbed following sublingual administration of nitroglycerin sublingual tablets. Mean peak nitroglycerin plasma concentrations occur at a mean time of approximately 6 to 7 minutes postdose (Table 1). Maximum plasma nitroglycerin concentrations (C_{max}) and area under the plasma concentration-time curves (AUC) increase dose-proportionally following 0.3 to 0.6 mg nitroglycerin. The absolute bioavailability of nitroglycerin from nitroglycerin sublingual tablets is approximately 40% but tends to be variable due to factors influencing drug absorption, such as sublingual hydration and mucosal metabolism.

Table 1

Parameter	Mean Nitroglycerin (SD) Values	
	2 x 0.3 mg Nitroglycerin Sublingual Tablets	1 x 0.6 mg Nitroglycerin Sublingual Tablets
C_{max} , ng/mL	2.3 (1.7)	2.1 (1.5)
T_{max} , min	6.4 (2.5)	7.2 (3.2)
AUC _(0-∞) , min	14.9 (8.2)	14.9 (11.4)
$t_{1/2}$, min	2.8 (1.1)	2.6 (0.6)

Distribution: The volume of distribution (V_{Area}) of nitroglycerin following intravenous administration is 3.3 L/kg. At plasma concentrations between 50 and 500 ng/mL, the binding of nitroglycerin to plasma proteins is approximately 60%, while that of 1,2- and 1,3-dinitroglycerin is 60% and 30%, respectively.

Metabolism: A liver reductase enzyme is of primary importance in the metabolism of nitroglycerin to glycerol di- and mononitrate metabolites and ultimately to glycerol and organic nitrate. Known sites of extrahepatic metabolism include red blood cells and vascular walls. In addition to nitroglycerin, 2 major metabolites 1,2- and 1,3-dinitroglycerin, are found in plasma. Mean peak 1,2- and 1,3-dinitroglycerin plasma concentrations occur at approximately 15 minutes postdose. The elimination half-life of 1,2- and 1,3-dinitroglycerin is 36 and 32 minutes, respectively. The 1,2- and 1,3-dinitroglycerin metabolites have been reported to possess approximately 2% and 10%, respectively, of the pharmacological activity of nitroglycerin. Higher plasma concentrations of the dinitro metabolites, along with their nearly 10-fold longer elimination half-lives, may contribute significantly to the duration of pharmacologic effect. Glycerol mononitrate metabolites of nitroglycerin are biologically inactive.

Elimination: Nitroglycerin plasma concentrations decrease rapidly, with a mean elimination half-life of 2 to 3 minutes. Half-life values range from 1.5 to 7.5 minutes. Clearance (13.6 L/min) greatly exceeds hepatic blood flow. Metabolism is the primary route of drug elimination.

INDICATIONS AND USAGE

Nitroglycerin sublingual tablets are indicated for the acute relief of an attack or acute prophylaxis of angina pectoris due to coronary artery disease.

CONTRAINDICATIONS

Nitroglycerin sublingual tablets are contraindicated in patients who are allergic to it.

Sublingual nitroglycerin therapy is contraindicated in patients with early myocardial infarction, severe anemia, increased intracranial pressure, and those with a known hypersensitivity to nitroglycerin.

Administration of nitroglycerin sublingual tablets are contraindicated in patients who are using a phosphodiesterase-5 (PDE-5) inhibitor (e.g., sildenafil citrate, tadalafil, vardenafil hydrochloride) since these compounds have been shown to potentiate the hypotensive effects of organic nitrates.

Do not use nitroglycerin sublingual tablets in patients who are taking the soluble guanylate cyclase stimulator riociguat. Concomitant use can cause hypotension.

WARNINGS

The benefits of sublingual nitroglycerin in patients with acute myocardial infarction or congestive heart failure have not been established. If one elects to use nitroglycerin in these conditions, careful clinical or hemodynamic monitoring must be used because of the possibility of hypotension and tachycardia.

PRECAUTIONS

General

Only the smallest dose required for effective relief of the acute anginal attack should be used. Excessive use may lead to the development of tolerance. Nitroglycerin sublingual tablets are intended for sublingual or buccal administration and should not be swallowed.

Severe hypotension, particularly with upright posture, may occur with small doses of nitroglycerin. This drug should therefore be used with caution in patients who may be volume-depleted or who, for whatever reason, are already hypotensive. Hypotension induced by nitroglycerin may be accompanied by paradoxical bradycardia and increased angina pectoris.

Nitrate therapy may aggravate the angina caused by hypertrophic cardiomyopathy.

As tolerance to other forms of nitroglycerin develops, the effects of sublingual nitroglycerin on exercise tolerance, although still observable, is blunted.

In industrial workers who have had long-term exposure to unknown (presumably high) doses of organic nitrates, tolerance rarely occurs. Chest pain, acute myocardial infarction, and even sudden death have occurred during temporary withdrawal of nitrates from these workers, demonstrating the existence of true physical dependence.

Several clinical trials of nitroglycerin patches or infusions in patients with angina pectoris have evaluated regimens that incorporated a 10- to 12-hour nitrate free interval. In some of these trials, an increase in the frequency of anginal attacks during the nitrate free interval was observed in a small number of patients. In one trial, patients had decreased exercise tolerance at the end of the nitrate interval. Hemodynamic rebound has been observed only rarely; on the other hand, few studies were so designed that rebound, if it had occurred, would have been detected.

Nitrate tolerance as a result of sublingual nitroglycerin administration is probably possible, but only in patients who maintain high continuous nitrate levels for more than 10 or 12 hours daily. Such use of sublingual nitroglycerin would entail administration of scores of tablets daily and is not recommended.

The drug should be discontinued if blurring of vision or drying of the mouth occurs. Excessive dosage of nitroglycerin may produce severe headaches.

Information for Patients

Nitroglycerin is a sublingual tablet and should not be chewed, crushed, or swallowed.

If possible, patients should sit down when taking nitroglycerin sublingual tablets and should use caution when returning to a standing position. This eliminates the possibility of falling due to lightheadedness

or dizziness.

One tablet should be dissolved under the tongue or in the buccal pouch at the first sign of an acute anginal attack. The dose may be repeated approximately every 5 minutes until relief is obtained.

If chest pain persists after a total of 3 tablets in a 15-minute period, or if the pain is different than is typically experienced, prompt medical attention is recommended.

Nitroglycerin may be used prophylactically 5 to 10 minutes prior to engaging in activities that might precipitate an acute attack.

Nitroglycerin may produce a burning or tingling sensation when administered sublingually; however, the ability to produce a burning or tingling sensation should not be considered a reliable method for determining the potency of the tablets.

Headaches can sometimes accompany treatment with nitroglycerin. In patients who get these headaches, the headaches may be a marker of the activity of the drug.

Treatment with nitroglycerin may be associated with lightheadedness upon standing, especially just after rising from a recumbent or seated position. This effect may be more frequent in patients who have also consumed alcohol.

Nitroglycerin should be kept in the original glass container and must be tightly capped after each use to prevent loss of tablet potency.

Drug Interactions

Concomitant use of nitroglycerin with soluble guanylate cyclasestimulators is contraindicated (see CONTRAINDICATIONS).

Concomitant use of nitrates and alcohol may cause hypotension.

The vasodilatory and hemodynamic effects of nitroglycerin may be enhanced by concomitant administration of aspirin.

Intravenous administration of nitroglycerin decreases the thrombolytic effect of alteplase. Therefore, caution should be observed in patients receiving sublingual nitroglycerin during alteplase therapy.

Intravenous nitroglycerin reduces the anticoagulant effect of heparin and activated partial thromboplastin times (APTT) should be monitored in patients receiving heparin and intravenous nitroglycerin. It is not known if this effect occurs following single sublingual nitroglycerin doses.

Tricyclic antidepressants (amitriptyline, desipramine, doxepin, others) and anticholinergic drugs may cause dry mouth and diminished salivary secretions. This may make dissolution of sublingual nitroglycerin difficult. Increasing salivation with chewing gum or artificial saliva products may prove useful in aiding dissolution of sublingual nitroglycerin.

Oral administration of nitroglycerin markedly decreases the first-pass metabolism of dihydroergotamine and subsequently increases its oral bioavailability. Ergotamine is known to precipitate angina pectoris. Therefore, patients receiving sublingual nitroglycerin should avoid ergotamine and related drugs or be monitored for symptoms of ergotism if this is not possible.

Administration of nitroglycerin is contraindicated in patients who are using PDE-5 inhibitors (e.g., sildenafil citrate, tadalafil, vardenafil hydrochloride). These compounds have been shown to potentiate the hypotensive effects of organic nitrates.

A decrease in therapeutic effect of sublingual nitroglycerin may result from use of long-acting nitrates.

Drug/Laboratory Test Interactions

Nitrates may interfere with the Zlatkis-Zak color reaction, causing a false report of decreased serum cholesterol.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Animal carcinogenesis studies with sublingually administered nitroglycerin have not been performed.

Carcinogenicity potential of nitroglycerin was evaluated in rats receiving up to 434 mg/kg/day of dietary nitroglycerin for 2 years. Rats developed dose-related fibrotic and neoplastic changes in liver, including carcinomas, and interstitial cell tumors in testes. At high dose, the incidences of hepatocellular carcinomas in males was 48% and in females was 33%, compared to 0% in untreated controls. Incidences of testicular tumors were 52% vs. 8% in controls. Lifetime dietary administration of up to 1058 mg/kg/day of, nitroglycerin was not tumorigenic in mice.

Nitroglycerin was mutagenic in Ames tests performed in 2 different laboratories. Nevertheless, there was no evidence of mutagenicity in an in vivo dominant lethal assay with male rats treated with doses up to about 363 mg/kg/day, PO, or in ex vivo cytogenetic tests in rat and dog cells.

In a 3-generation reproduction study, rats received dietary nitroglycerin at doses up to about 434 mg/kg/day for 6 months prior to mating of the F₀ generation, with treatment continuing through successive F₁ and F₂ generations. The high dose was associated with decreased feed intake and body weight gain in both sexes at all matings. No specific effect on the fertility of the F₀ generation was seen. Infertility noted in subsequent generations, however, was attributed to increased interstitial cell tissue and aspermatogenesis in the high-dose males. In this 3-generation study, there was no clear evidence of teratogenicity.

Pregnancy: Category B.

Animal reproduction and teratogenicity studies have not been conducted with nitroglycerin sublingual tablets. However, teratology studies conducted in rats and rabbits with topically applied nitroglycerin ointment at dosages up to 80 mg/kg/day and 240 mg/kg/day, respectively revealed no toxic effects on dams or fetuses.

There are no adequate and well-controlled studies in pregnant women. Nitroglycerin should be given to a pregnant woman only if clearly needed.

Nursing Mothers

It is not known whether nitroglycerin is excreted in human milk.

Because many drugs are excreted in human milk, caution should be exercised when nitroglycerin is administered to a nursing woman.

Pediatric Use

The safety and effectiveness of nitroglycerin in pediatric patients have not been established.

Geriatric Use

Clinical studies of nitroglycerin did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

ADVERSE REACTIONS

Headache that may be severe and persistent may occur immediately after use. Vertigo, dizziness, weakness, palpitation, and other manifestations of postural hypotension may develop occasionally, particularly in erect, immobile patients. Marked sensitivity to the hypotensive effects of nitrates (manifested by nausea, vomiting, weakness, diaphoresis, pallor, and collapse) may occur at therapeutic

doses. Syncope due to nitrate vasodilatation has been reported. Flushing, drug rash, and exfoliative dermatitis have been reported in patients receiving nitrate therapy.

OVERDOSAGE

Hemodynamic Effects: The effects of nitroglycerin overdose are generally the results of nitroglycerin's capacity to induce vasodilatation, venous pooling, reduced cardiac output, and hypotension. These hemodynamic changes may have protean manifestations, including increased intracranial pressure, with any or all of persistent throbbing headache, confusion, and moderate fever; vertigo; palpitations; tachycardia; visual disturbances; nausea and vomiting (possibly with colic and even bloody diarrhea); syncope (especially in the upright posture); dyspnea, later followed by reduced ventilatory effort; diaphoresis, with the skin either flushed or cold and clammy; heart block and bradycardia; paralysis; coma; seizures; and death.

No specific antagonist to the vasodilator effects of nitroglycerin is known, and no intervention has been subject to controlled study as a therapy of nitroglycerin overdose. Because the hypotension associated with nitroglycerin overdose is the result of venodilatation and arterial hypovolemia, prudent therapy in this situation should be directed toward increase in central fluid volume. Passive elevation of the patient's legs may be sufficient, but intravenous infusion of normal saline or similar fluid may also be necessary.

The use of epinephrine or other arterial vasoconstrictors in this setting is likely to do more harm than good.

In patients with renal disease or congestive heart failure, therapy resulting in central volume expansion is not without hazard. Treatment of nitroglycerin overdose in these patients may be subtle and difficult, and invasive monitoring may be required.

Methemoglobinemia: Methemoglobinemia has been rarely reported in association with organic nitrates. The diagnosis should be suspected in patients who exhibit signs of impaired oxygen delivery despite adequate cardiac output and adequate arterial PO₂. Classically, methemoglobinemic blood is described as chocolate brown, without color change on exposure to air.

If methemoglobinemia is present, intravenous administration of methylene blue, 1 to 2 mg/kg of body weight, may be required.

DOSAGE AND ADMINISTRATION

One tablet should be dissolved under the tongue or in the buccal pouch at the first sign of an acute anginal attack. The dose may be repeated approximately every 5 minutes until relief is obtained. If the pain persists after a total of 3 tablets in a 15-minute period, or if the pain is different than is typically experienced, prompt medical attention is recommended. Nitroglycerin may be used prophylactically 5 to 10 minutes prior to engaging in activities that might precipitate an acute attack.

During administration the patient should rest, preferably in the sitting position.

No dosage adjustment is required in patients with renal failure.

HOW SUPPLIED

Nitroglycerin sublingual tablets are supplied in 3 strengths (0.3 mg, 0.4 mg, and 0.6 mg) in bottles containing 100 tablets each, with color-coded labels, and in color-coded Patient Convenience Packages of 4 bottles of 25 tablets each.

0.3 mg sublingual tablets are white to off-white, modified rectangle shaped tablets debossed with "CL" on one side and "3" on the other side and are supplied in bottles of 100 tablets.

Bottle of 100 NDC 60429-839-01

0.4 mg sublingual tablets are white to off-white, modified rectangle shaped tablets debossed with "CL" on one side and "4" on the other side and are supplied in bottles of 25 and 100 tablets.

Bottle of 100 NDC 60429-840-01

Convenience Package NDC 60429-840-04

0.6 mg sublingual tablets are white to off-white, modified rectangle shaped tablets debossed with "CL" on one side and "6" on the other side and are supplied in bottles of 100 tablets.

Bottle of 100 NDC 60429-841-01

Store at 20°-25°C (68°-77°F) [see USP Controlled Room Temperature].

Rx only

Manufactured by:

Ingenus Pharmaceuticals NJ, LLC

Fairfield, NJ 07004, USA

Manufactured for:

Dr. Reddy's Laboratories Inc.

Princeton, New Jersey 08540, USA

Marketed/Packaged by:

GSMS, Inc.

Camarillo, CA USA 93012

Revised: 0217

PATIENT PACKAGE INSERT

Nitroglycerin Sublingual Tablets, USP

(nahy-truh-glis-er-in)

Read this information carefully before you start **nitroglycerin sublingual tablets** and each time you refill your prescription. There may be new information. This information does not replace talking with your doctor. If you have any questions about **nitroglycerin**, ask your doctor. Your doctor will know if **nitroglycerin sublingual tablet** is right for you.

What is nitroglycerin?

Nitroglycerin is a type of medicine known as an organic nitrate and is a vasodilating agent. It is used to treat a type of chest pain called angina.

What is Angina?

Angina is a pain or discomfort that keeps coming back when part of your heart does not get enough blood. Angina feels like a pressing or squeezing pain, usually in your chest under the breastbone. Sometimes you can feel it in your shoulders, arms, neck, jaws, or back. **Nitroglycerin sublingual tablets** can relieve this pain.

Who should not use nitroglycerin sublingual tablets?

Do not use **nitroglycerin sublingual tablets** if you are allergic to organic nitrates (like the active ingredient in **nitroglycerin sublingual tablets**).

You should not take **nitroglycerin sublingual tablets** if you have the following conditions:

- very recent heart attack

- severe anemia
- increased pressure in the head

Do not take **nitroglycerin sublingual tablets** with drugs for erectile dysfunction, like VIAGRA[®] (sildenafil citrate), CIALIS[®] (tadalafil), or LEVITRA[®] (vardenafil hydrochloride), as this may lead to extreme lowering of your blood pressure.

Do not take **nitroglycerin sublingual tablets** if you take medicines called guanylate cyclase stimulators which include riociguat, a medicine that treats pulmonary arterial hypertension and chronic-thromboembolic pulmonary hypertension.

What should I tell my doctor before taking nitroglycerin sublingual tablets?

Before using **nitroglycerin sublingual tablets**, tell your doctor if:

- You are taking any medicines that are used to treat angina, heart failure, or an irregular heartbeat.
- You are taking any medicines that reduce blood pressure.
- You are taking any diuretics (water pills).
- You are taking medications to treat depression or psychiatric illness.
- You are taking ergotamine or similar drugs for migraine headaches.
- You are taking aspirin.
- You are taking the blood thinner medicine heparin.
- You are taking any medicines for erectile dysfunction.
- You are pregnant or plan to become pregnant.
- You are breastfeeding.

How should I take nitroglycerin sublingual tablets?

- Do not chew, crush, or swallow **nitroglycerin** sublingual tablets.
- You should sit down when taking **nitroglycerin** sublingual tablets and use caution when you stand up. This eliminates the possibility of falling due to lightheadedness or dizziness.
- One tablet should be dissolved under the tongue or in the oral cavity at the first sign of chest pain.
- The dose may be repeated approximately every 5 minutes, until the chest pain is relieved.
- If the pain persists after a total of 3 tablets in a 15-minute period, or is different than you typically experience, call your doctor or seek emergency help.
- **Nitroglycerin sublingual tablets** may be used 5 to 10 minutes prior to activities that might cause chest pain.
- You may feel a burning or tingling sensation in your mouth when you take **nitroglycerin sublingual tablets**.

What should I avoid while taking nitroglycerin sublingual tablets?

- Do not breastfeed. It is not known if **nitroglycerin** will pass through your milk.
- Do not consume alcohol while taking **nitroglycerin sublingual tablets**, as this can lower your blood pressure.
- Do not start any new prescription or non-prescription medicines or supplements, unless you check with your doctor first.

What are the possible side effects of nitroglycerin sublingual tablets?

Nitroglycerin may cause the following side effects:

- headache
- vertigo (a major symptom of balance disorder)
- dizziness
- weakness

- heart palpitations (unusual awareness of the heartbeat)
- low blood pressure upon rising from a seated position
- nausea and vomiting
- sweating
- paleness
- fainting
- flushing (warm or red condition of your skin)
- other skin reactions that may be severe

Nitroglycerin sublingual tablets may cause a false test result of decreased serum cholesterol.

Tell your doctor if you are concerned about any side effects you experience. These are not all the possible side effects of **nitroglycerin**. For a complete list, ask your doctor or pharmacist.

How do I store nitroglycerin sublingual tablets?

Nitroglycerin sublingual tablets should be kept in the original glass container and tightly capped after each use to prevent loss of tablet potency.

Store **nitroglycerin sublingual tablets** at 20°-25°C (68°-77°F) [see USP Controlled Room Temperature].

General advice about nitroglycerin sublingual tablets

Sometimes doctors will prescribe a medicine for a condition that is not included in the patient information leaflets. Only use **nitroglycerin sublingual tablets** the way your doctor told you to. Do not give **nitroglycerin sublingual tablets** to other people, even if they have the same symptoms you have. It may harm them.

You can ask your pharmacist or doctor for information about **nitroglycerin sublingual tablets**, or call Dr. Reddy's Medical Information Department, at 1-888-375-3784.

To reorder additional Patient Package Inserts contact Dr. Reddy's Customer Service at 1-866-733-3952.

Rx Only

Manufactured by:

Ingenus Pharmaceuticals NJ, LLC

Fairfield, NJ 07004, USA

Manufactured for:

Dr. Reddy's Laboratories Inc.

Princeton, New Jersey 08540, USA

Marketed/Packaged by:


GSMS, Inc.

Camarillo, CA USA 93012


Revised: 0217

PACKAGE LABEL PRINCIPAL DISPLAY PANEL SECTION

0.3 mg : Container Label: 100's

<i>GSMS, Incorporated</i>	
60429-839-01	
NITROGLYCERIN	
SUBLINGUAL TABLETS, USP	
0.3 mg	
100 sublingual tablets	
See package insert for dosage information.	
<i>Rx only</i>	
Store at 20°-25°C (68°-77°F) [See USP Controlled Room Temperature]	
	
6042983901	
<ul style="list-style-type: none">• Each tablet contains: 0.3 mg nitroglycerin.• Keep this and all medication out of the reach of children.• Warning - Close tightly immediately after each use to prevent loss of potency.• Do not crush, chew, or swallow Nitroglycerin Sublingual Tablets.• Always Dispense with Patient Package Insert.	<p>Exp.: 02/2021</p> <p>Lot: GS027535</p>
Manufactured by Ingenus Pharmaceuticals N.J., LLC Fairfield, NJ 07004, USA Packaged by GSMS, Incorporated Camranillo, CA 93012, USA	

0.3 mg: Container Carton Label: 100's



0.4 mg: Container Label: 25's

NDC 60429-840-25

**NITROGLYCERIN
SUBLINGUAL TABLETS, USP**

0.4 mg/tablet

**Always Dispense with
Patient Package Insert**

Rx Only

25 Sublingual Tablets



Each tablet contains: 0.4 mg nitroglycerin

USUAL DOSAGE:

See accompanying prescribing information.

Store at 20°-25° C (68°-77°F)

[see USP Controlled Room Temperature]

Keep this and all drugs

out of the reach of children.

Warning - Close tightly immediately

after each use to prevent loss of potency.

Keep these tablets in the original container.

Do not crush, chew, or swallow

Nitroglycerin Sublingual Tablets.

Manufactured by:

Ingenus Pharmaceuticals NJ, LLC

Fairfield, NJ 07004, USA

Marketed by:

GSMS, Incorporated

Camarillo, CA 93012 USA


5081




6042984025

LOT
EXP


0.4 mg : Container Carton Label: 4 x 25's count



0.6 mg : Container Label: 100's

<p><i>GSMS, Incorporated</i></p> <p>60429-841-01</p> <p>NITROGLYCERIN</p> <p>SUBLINGUAL TABLETS, USP</p> <p>0.6 mg</p> <p>100 sublingual tablets</p> <p>See package insert for dosage information.</p> <p><i>Rx only</i></p> <p>Store at 20°-25°C (68°-77°F) [See USP Controlled Room Temperature]</p>		<p>6042984101</p> 	<ul style="list-style-type: none"> • Each tablet contains: 0.6 mg nitroglycerin. • Keep this and all medication out of the reach of children. • Warning - Close tightly immediately after each use to prevent loss of potency. • Do not crush, chew, or split Nitroglycerin Sublingual Tablets. • Always Dispense with Patient Package Insert.
<p>Exp.: 03/2021</p> <p>Lot: GS029010</p>		<p>Manufactured by: Ingenue Pharmaceuticals NJ, LLC Fairfield, NJ 07004, USA Packaged by: GSMS, Incorporated Camden, NJ 08202, USA</p>	

0.6 mg: Container Carton Label: 100's



NITROGLYCERIN

nitroglycerin tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:60 429-8 39(NDC:43598-435)
Route of Administration	SUBLINGUAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
NITROGLYCERIN (UNII: G59M7S0WS3) (NITROGLYCERIN - UNII:G59M7S0WS3)	NITROGLYCERIN	0.3 mg

Inactive Ingredients	
Ingredient Name	Strength
CALCIUM STEARATE (UNII: 776XM7047L)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYDROGENATED COTTONSEED OIL (UNII: Z82Y2C65EA)	

Product Characteristics			
Color	white (off white)	Score	no score
Shape	RECTANGLE	Size	4mm
Flavor		Imprint Code	CL;3
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60429-839-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/12/2016	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA208191	08/26/2016	

NITROGLYCERIN

nitroglycerin tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:60429-840(NDC:43598-436)
Route of Administration	SUBLINGUAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
NITROGLYCERIN (UNII: G59M7S0WS3) (NITROGLYCERIN - UNII:G59M7S0WS3)	NITROGLYCERIN	0.4 mg

Inactive Ingredients	
Ingredient Name	Strength

LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYDROGENATED COTTONSEED OIL (UNII: Z82Y2C65EA)	
CALCIUM STEARATE (UNII: 776XM7047L)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	white (off white)	Score	no score
Shape	RECTANGLE	Size	4mm
Flavor		Imprint Code	CL;4
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60429-840-04	4 in 1 CARTON	09/12/2016	
1		25 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:60429-840-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/12/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA208191	08/26/2016	

NITROGLYCERIN

nitroglycerin tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:60429-841(NDC:43598-437)
Route of Administration	SUBLINGUAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NITROGLYCERIN (UNII: G59M7S0WS3) (NITROGLYCERIN - UNII:G59M7S0WS3)	NITROGLYCERIN	0.6 mg

Inactive Ingredients

Ingredient Name	Strength
CALCIUM STEARATE (UNII: 776XM7047L)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYDROGENATED COTTONSEED OIL (UNII: Z82Y2C65EA)	

Product Characteristics

Color	white (off white)	Score	no score
Shape	RECTANGLE	Size	4mm
Flavor		Imprint Code	CL;6
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60429-841-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/12/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA208191	08/26/2016	

Labeler - Golden State Medical Supply, Inc. (603184490)

Establishment

Name	Address	ID/FEI	Business Operations
Golden State Medical Supply, Inc.		603184490	relabel(60429-839, 60429-840, 60429-841) , repack(60429-839, 60429-840, 60429-841)

Revised: 12/2019

Golden State Medical Supply, Inc.