

**BRONCOCHEM MAXIMUM COUGH NIGHT- dextromethorphan hbr-pyrimilamine maleate syrup**  
**LABORATORIO MAGNACHEM INTERNATIONAL SRL**

-----  
**Broncochem Maximum Cough Night**

**Warnings**

The following side effects may occur, nausea, vomiting, dizziness, headache, or nervousness may occur, if any of these effects last or worsen, contact your doctor or pharmacist immediately. Stop using this medicine and tell your doctor right away if you have dizziness, nervousness, or trouble sleeping.

**OTC-PURPOSE SECTION**

Antitussive

Antihistamine

**OTC - Active Ingredient Section**

Dextromethorphan HBr

Mepyramine maleate

**Uses**

Temporarily relieves these symptoms in all forms of acute, subacute, and chronic cough of any origin, it is very effective in colds, rhinitis, pharyngitis, laryngitis, dry cough, smokers cough, asthma, nervous cough.

**Do Not Use**

Stop using this medicine and contact your doctor right away if any of these unlikely but serious side effects occur, fast and irregular heartbeat, mental mood changes such as confusion, hallucinations, nervousness, seizures, tremor, problems with to pee.

If you are taking any monoamine oxidase inhibitor (MAOI), or two weeks after stopping them (certain medications for depression or psychological conditions may contain them). If you are not sure if your prescription contains an MAOI, consult a professional of health before taking this product.

**Ask a doctor/pharmacist**

Ask a doctor before use if you have

Liver disease, heart disease, high blood pressure, thyroid disease, diabetes, trouble urinating due to enlarged prostate gland, cough that occurs with too much phlegm

(mucus), persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema.

### **Stop use and ask a doctor if**

Symptoms last more than 7 days, symptoms worsen or come back, you develop a headache that won't go away, fever, or skin rash. If pregnant or breast-feeding ask a health professional before use.

### **Keep Out of the Reach of Children**

In case of accidental overdose, seek professional help or go to poison control center immediately.

### **Dosage and Administration**

Do not exceed 4 doses in a 24 hour period.

Adult and children 12 years and older: 10 mL 3-4 times a day.

Children 6 to under 12 years old: 5 mL 3-4 times a day.

### **Inactive Ingredient**

Sugar, Sodium Benzoate, FD&C Yellow #5, Orange flavor, Methyl paraben, Propyl paraben, ethyl alcohol 95%, Sorbitol 70%, Purified water.

### **PACKAGE LABEL PRINCIPAL DISPLAY PANEL**

8.802" 223.58mm

1. Empuje hacia abajo
2. Rota la tapa
3. Utilice la cuchara incluida



2.126" 54mm

**Información del Medicamento**

**Ingredientes Activos (en 5 mL ó cc)**

|                     |          |           |               |
|---------------------|----------|-----------|---------------|
| Mepyrmino maleato   | 50.00 mg | Propósito | Antitussígeno |
| Dextrometorphan HBr | 30.00 mg |           | Antitusivo    |

**Usos**

Está indicado para el tratamiento temporal en todas las formas de los ataques, subagudas y crónicas de cualquier origen, es muy eficaz en el tratamiento, tanto la tos seca, como la tos con flemas, así como, los resaca.

**Advertencias**

Los siguientes efectos adversos se pueden presentar, náusea, vómito, mareo, dolor de cabeza o resaca pueden ocurrir, si alguno de estos efectos es duradero o empeora, comuníquese con su doctor o farmacéutico inmediatamente. Evite el uso de este medicamento y comuníquese a su doctor rápidamente si usted tiene mareo, resaca o problemas para comer.

**No usar**

Si durante el uso de este medicamento y comuníquese con su doctor de inmediato si alguno de estos pocos problemas pero serios efectos adversos ocurren: serios problemas de visión o problemas, cambios de humor mental como confusión, alucinaciones, enrojecimiento, convulsiones, temblor, problemas al caminar.

Si está tomando algún inhibidor de la mono amino oxidasa (MAOI), o dos semanas después de haberlos suspendido (estos medicamentos para la depresión o condiciones psicológicas pueden contenerlos). Si usted no está seguro de que su receta contiene un MAOI consulte a un profesional de la salud antes de tomar este producto.

Pregunte a su médico antes de usar si usted tiene enfermedad hepática, enfermedad cardíaca, presión arterial alta, enfermedad de glándulas endocrinas, problemas para orinar, debido a agrandamiento de la próstata, tos con mucha flema (mucofobia), los asmáticos, asma o enfisema.

**Para el uso de este medicamento y consulte a su médico.**

- Si los síntomas duran más de 7 días
- Si los síntomas empeoran o regresan
- Si usted desarrolla un dolor de cabeza que no se quita, fiebre, o reacciones cutáneas.

Si está embarazada o en período de lactancia, consulte un profesional de la salud antes de usar.

Mantenga fuera del alcance de los niños.

**Advertencia de sobredosis:** En caso de sobredosis accidental, busque ayuda profesional o acuda a un Centro de Intoxicaciones inmediatamente.

**Instrucciones**

Siga las recomendaciones detalladas más abajo. No exceda de 4 dosis en un período de 24 horas.

|  |                           |
|--|---------------------------|
| Adultos y niños de 12 años en adelante | 10 mL, 3 o 4 veces al día |
| Niños entre 6 a 12 años                | 5 mL, 3 o 4 veces al día  |

**Información adicional**

- Contiene a no más de 30% (BPSP)
- No lo use si la tapa de seguridad está alterada o rota.

**Este producto contiene alcohol y azúcar.**

**Ingredientes Inactivos**

Azúcar refinado, Benzato de sodio, Color amarillo #5, Esencia de naranja, Metil paraben, Propil paraben, Alcohol etílico al 95%, Sorbitol al 70%, Agua purificada.

**Preguntas?** 1-661-338-6221 o visitamos [www.broncochem.com](http://www.broncochem.com)

Fabricado en República Dominicana  
Equipamiento para Global Corporation Inc., Boca Raton, FL 33487 USA.

**REGISTRADO EN LA FDA**

**NEW!** NDC-65131-110-35

**Broncochem**  
NIGHT MAXIMUM COUGH

Contains Mepyrmino maleate, Dextrometorphan HBr

**Relieves cough of all origins**  
**Powerful cough suppressant to help you sleep at night**  
**Contains Alcohol & Sugar**

4 FL. OZ. (120 mL or cc)

**Drug Facts**

**Active Ingredients (in each 5 mL)**

|                     |          |         |             |
|---------------------|----------|---------|-------------|
| Mepyrmino maleate   | 50.00 mg | Purpose | Antitussive |
| Dextrometorphan HBr | 30.00 mg |         | Antitussive |

**Uses**

Temporarily relieves these symptoms in all forms of acute, subacute, and chronic cough of any origin. It is very effective in colds, rhinitis, pharyngitis, laryngitis, dry cough, smoker's cough, asthma, nervous cough.

**Warnings**

The following side effects may occur, nausea, vomiting, dizziness, headache, or nervousness may occur, if any of these effects last or worsen, contact your doctor or pharmacist immediately. Stop using this medicine and tell your doctor right away if you have dizziness, nervousness, or trouble sleeping.

**Do not use**

Stop using this medicine and contact your doctor right away if any of these unlikely but serious side effects occur: feel an irregular heartbeat, mental mood changes such as confusion, hallucinations, nervousness, seizures, tremor, problems with eye.

If you are taking any monoamine oxidase inhibitor (MAOI), or two weeks after stopping them (certain medications for depression or psychological conditions may contain them). If you are not sure if your prescription contains an MAOI, consult a professional of health before taking this product.

**Ask a doctor before use if you have**

Liver disease, heart disease, high blood pressure, thyroid disease, diabetes, trouble urinating due to enlarged prostate gland, cough that occurs with too much phlegm (mucus), persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema.

**Stop use and ask a doctor if**

- Symptoms last more than 7 days
- If symptoms worsen or come back
- If you develop a headache that won't go away, fever, or skin rash.

If pregnant or breast-feeding, ask a health professional before use. Keep out of the reach of children.

**Overdose warning:** In case of accidental overdose, seek professional help or go to a Poison Control Center immediately.

**Instructions**

Follow the recommendations detailed below. Do not exceed 4 doses in a 24-hour period.

|  |                           |
|--|---------------------------|
| Adults and children 12 years and older | 10 mL, 3 or 4 times a day |
| Children between 6 to 12 years old     | 5 mL, 3 or 4 times a day  |

**Other information**

- Store at no more than 30°C (86°F)
- Do not use if security cap is altered or broken.

**This product contains alcohol and sugar.**

**Inactive ingredients**

Refined sugar, Sodium benzoate, Yellow #5 color, Orange flavor, Methyl paraben, Propyl paraben, Ethyl alcohol 95%, Sorbitol 70%, Purified water.

**Questions?** 1-661-338-6221 or visit us at [www.broncochem.com](http://www.broncochem.com)

Made in Dominican Republic  
Equipamiento para Global Corporation Inc., Boca Raton, FL 33487 USA.

**FDA REGISTERED**

**NEW!** NDC-65131-110-35

**Broncochem**  
NOCHE MAXIMUM TOS

Contiene Mepyrmino maleato, Dextrometorphan HBr

**Alivia todo tipo de tos**  
**Potente supresor de la tos para ayudarte a dormir mejor**  
**Contiene Alcohol y Azúcar**

4 FL. OZ. (120 mL ó cc)

5.157" 131mm



Batch / Lote:  
Expiration date:



8.802" 223.58mm

1. Empuje hacia abajo
2. Rota la tapa
3. Utilice la cuchara incluida



2.126" 54mm

**Información del Medicamento**

**Ingredientes Activos (en 5 mL ó cc)**

|                     |          |           |               |
|---------------------|----------|-----------|---------------|
| Mepyrmino maleato   | 50.00 mg | Propósito | Antitussígeno |
| Dextrometorphan HBr | 30.00 mg |           | Antitusivo    |

**Usos**

Está indicado para el tratamiento temporal en todas las formas de los ataques, subagudas y crónicas de cualquier origen, es muy eficaz en el tratamiento, irrita la garganta, la nariz, los ojos, los conductos auditivos, la nariz, los nervios.

**Advertencias**

Los siguientes efectos adversos se pueden presentar, náusea, vómito, mareo, dolor de cabeza o resaca pueden ocurrir, si alguno de estos efectos es duradero o empeora, comuníquese con su doctor o farmacéutico inmediatamente. Evite el uso de este medicamento y comuníquese a su doctor rápidamente si usted tiene mareo, nerviosismo o problemas para dormir.

**No usar**

Si tiene el uso de este medicamento y comuníquese con su doctor de inmediato si alguno de estos pocos problemas pero serios efectos adversos ocurren: serios problemas de visión, cambios de humor mental como confusión, alucinaciones, enrojecimiento, convulsiones, temblor, problemas al caminar.

Si está tomando algún inhibidor de la mono amino oxidasa (MAOI), o dos semanas después de haberlos suspendido (estos medicamentos para la depresión o condiciones psicológicas pueden contenerlos). Si usted no está seguro de que su receta contiene un MAOI consulte a un profesional de la salud antes de tomar este producto.

Pregunte a su médico antes de usar si usted tiene enfermedad hepática, enfermedad cardíaca, presión arterial alta, enfermedad de glándulas endocrinas, problemas para orinar, debido a agrandamiento de la próstata, tos con mucha flema (mucofisi), los asmáticos, asma o epilepsia.

**Para el uso de este medicamento y consulte a su médico.**

- Si los síntomas duran más de 7 días
- Si los síntomas empeoran o regresan
- Si usted desarrolla un dolor de cabeza que no se quita, fiebre, o reacciones cutáneas.

Si está embarazada o en período de lactancia, consulte un profesional de la salud antes de usar.

Mantenga fuera del alcance de los niños.

**Advertencia de sobredosis:** En caso de sobredosis accidental, busque ayuda profesional o acuda a un Centro de Intoxicaciones inmediatamente.

**Instrucciones**

Siga las recomendaciones detalladas más abajo. No exceda de 4 dosis en un período de 24 horas.

|  |                           |
|--|---------------------------|
| Adultos y niños de 12 años en adelante | 10 mL, 3 o 4 veces al día |
| Niños entre 6 a 12 años                | 5 mL, 3 o 4 veces al día  |

**Información adicional**

- Contiene a no más de 30% (BPSP)
- No lo use si la tapa de seguridad está alterada o rota.

**Este producto contiene alcohol y azúcar.**

**Ingredientes Inactivos**

Azúcar refinado, Benzato de sodio, Color amarillo #5, Esencia de naranja, Metil paraben, Propil paraben, Alcohol etílico al 95%, Sorbitol al 70%, Agua purificada.

**Preguntas?** 1-661-338-6221 o visitamos [www.broncochem.com](http://www.broncochem.com)

Fabricado en República Dominicana  
Equipamiento para Global Corporation Inc., Boca Raton, FL 33487 USA.

**REGISTRADO EN LA FDA**

**NEW!** NDC-65131-110-35

**Broncochem**  
NIGHT MAXIMUM COUGH

Contains Mepyrmino maleate, Dextrometorphan HBr

**Relieves cough of all origins**  
**Powerful cough suppressant to help you sleep at night**  
**Contains Alcohol & Sugar**

4 FL. OZ. (120 mL or cc)

**Drug Facts**

**Active Ingredients (in each 5 mL)**

|                     |          |         |             |
|---------------------|----------|---------|-------------|
| Mepyrmino maleate   | 50.00 mg | Purpose | Antitussive |
| Dextrometorphan HBr | 30.00 mg |         | Antitussive |

**Uses**

Temporarily relieves these symptoms in all forms of acute, subacute, and chronic cough of any origin. It is very effective in colds, rhinitis, pharyngitis, laryngitis, dry cough, smoker's cough, asthma, nervous cough.

**Warnings**

The following side effects may occur, nausea, vomiting, dizziness, headache, or nervousness may occur, if any of these effects last or worsen, contact your doctor or pharmacist immediately. Stop using this medicine and tell your doctor right away if you have dizziness, nervousness, or trouble sleeping.

**Do not use**

Stop using this medicine and contact your doctor right away if any of these untreatable but serious side effects occur: feel and/or regular heartbeats, mental mood changes such as confusion, hallucinations, nervousness, seizures, tremor, problems with toe.

If you are taking any monoamine oxidase inhibitor (MAOI), or two weeks after stopping them (part from medications for depression or psychological conditions may contain them). If you are not sure if your prescription contains an MAOI, consult a professional of health before taking this product.

**Ask a doctor before use if you have**

Liver disease, heart disease, high blood pressure, thyroid disease, diabetes, trouble urinating due to enlarged prostate gland, cough that occurs with too much phlegm (mucus), persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema.

**Stop use and ask a doctor if**

- Symptoms last more than 7 days
- If symptoms worsen or come back
- If you develop a headache that won't go away, fever, or skin rash.

If pregnant or breast-feeding, ask a health professional before use. Keep out of the reach of children.

**Overdose warning:** In case of accidental overdose, seek professional help or go to a Poison Control Center immediately.

**Instructions**

Follow the recommendations detailed below. Do not exceed 4 doses in a 24-hour period.

|  |                           |
|--|---------------------------|
| Adults and children 12 years and older | 10 mL, 3 or 4 times a day |
| Children between 6 to 12 years old     | 5 mL, 3 or 4 times a day  |

**Other information**

- Store at no more than 30°C (86°F)
- Do not use if security cap is altered or broken.

**This product contains alcohol and sugar.**

**Inactive ingredients**

Refined sugar, Sodium benzoate, Yellow #5 color, Orange flavor, Methyl paraben, Propyl paraben, Ethyl alcohol 95%, Sorbitol 70%, Purified water.

**Questions?** 1-661-338-6221 or visit us at [www.broncochem.com](http://www.broncochem.com)

Made in Dominican Republic  
Equipamiento para Global Corporation Inc., Boca Raton, FL 33487 USA.

**FDA REGISTERED**

**NEW!** NDC-65131-110-35

**Broncochem**  
NOCHE MAXIMUM TOS

Contiene Mepyrmino maleato, Dextrometorphan HBr

**Alivia todo tipo de tos**  
**Potente supresor de la tos para ayudarte a dormir mejor**  
**Contiene Alcohol y Azúcar**

4 FL. OZ. (120 mL ó cc)

5.157" 131mm




Batch / Lote:  
Expiration date:



8.802" 223.58mm

1. Empléjelo hasta albio
2. Rótela la tapa
3. Utilice la cuchara incluída



2.126" 54mm

**Información del Medicamento**

**Ingredientes Activos (en 5 mL ó cc)**

|                     |          |           |               |
|---------------------|----------|-----------|---------------|
| Mepyrmino maleato   | 50.00 mg | Propósito | Antitussígeno |
| Dextrometorphan HBr | 30.00 mg |           | Antitusivo    |

**Usos**

Está indicado para el tratamiento temporal en todas las formas de los ataques, subagudas y crónicas de cualquier origen, es muy eficaz en el tratamiento, irrita la garganta, la nariz, los ojos, los conductos auditivos, así como, los ruidos.

**Advertencias**

Los siguientes efectos adversos se pueden presentar, náusea, vómito, mareo, dolor de cabeza o resaca pueden ocurrir, si alguno de estos efectos es duradero o empeora, comuníquese con su doctor o farmacéutico inmediatamente. Evite el uso de este medicamento y comuníquese a su doctor rápidamente si usted tiene mareo, nerviosismo o problemas para dormir.

**No usar**

Si durante el uso de este medicamento y comuníquese con su doctor de inmediato si alguno de estos pocos problemas pero serios efectos adversos ocurren: serios problemas de visión, cambios de humor mental como confusión, alucinaciones, enrojecimiento, convulsiones, temblor, problemas al caminar.

Si está tomando algún inhibidor de la mono amino oxidasa (MAOI), o dos semanas después de haberlos suspendido (estos medicamentos para la depresión o condiciones psicológicas pueden contenerlos).

Si usted no está seguro de que su receta contiene un MAOI consulte a un profesional de la salud antes de tomar este producto.

Pregunte a su médico antes de usar si usted tiene enfermedad hepática, enfermedad cardíaca, presión arterial alta, enfermedad de glándulas endocrinas, problemas para orinar, debido a agrandamiento de la próstata, tos con mucha flema (mucoosidad), los asmáticos, asma o epilepsia.

**Para el uso de este medicamento y consulte a su médico.**

- Si los síntomas duran más de 7 días
- Si los síntomas empeoran o regresan
- Si usted desarrolla un dolor de cabeza que no se quita, fiebre, o reacciones cutáneas.

Si está embarazada o en período de lactancia, consulte un profesional de la salud antes de usar.

Manténgala fuera del alcance de los niños.

**Advertencia de sobredosis:** En caso de sobredosis accidental, busque ayuda profesional o acuda a un Centro de Intoxicaciones inmediatamente.

**Instrucciones**

Siga las recomendaciones detalladas más abajo. No exceda de 4 dosis en un período de 24 horas.

|  |                           |
|--|---------------------------|
| Adultos y niños de 12 años en adelante | 10 mL, 3 o 4 veces al día |
| Niños entre 6 a 12 años                | 5 mL, 3 o 4 veces al día  |

**Información adicional**

- Contiene a no más de 30% (BPSP)
- No lo use si la tapa de seguridad está alterada o rota.

**Este producto contiene alcohol y azúcar.**

**Ingredientes Inactivos**

Azúcar refinado, Benzato de sodio, Color amarillo #5, Esencia de naranja, Metil paraben, Propil paraben, Alcohol etílico al 95%, Sorbitol al 70%, Agua purificada.

**Preguntas?** 1-661-338-6221 o visitamos [www.broncochem.com](http://www.broncochem.com)

Fabricado en República Dominicana  
Equipamiento para Global Corporation Inc., Boca Raton, FL 33487 USA.

**REGISTRADO EN LA FDA**

**NEW!** NDC-65131-110-35

**Broncochem**  
NIGHT MAXIMUM COUGH

Contains Mepyrmino maleate, Dextrometorphan HBr

**Relieves cough of all origins**  
**Powerful cough suppressant to help you sleep at night**  
**Contains Alcohol & Sugar**

4 FL. OZ. (120 mL or cc)

**Drug Facts**

**Active Ingredients (in each 5 mL)**

|                     |          |         |             |
|---------------------|----------|---------|-------------|
| Mepyrmino maleate   | 50.00 mg | Purpose | Antitussive |
| Dextrometorphan HBr | 30.00 mg |         | Antitussive |

**Uses**

Temporarily relieves these symptoms in all forms of acute, subacute, and chronic cough of any origin. It is very effective in colds, rhinitis, pharyngitis, laryngitis, dry cough, smoker's cough, asthma, nervous cough.

**Warnings**

The following side effects may occur, nausea, vomiting, dizziness, headache, or nervousness may occur, if any of these effects last or worsen, contact your doctor or pharmacist immediately. Stop using this medicine and tell your doctor right away if you have dizziness, nervousness, or trouble sleeping.

**Do not use**

Stop using this medicine and contact your doctor right away if any of these untreatable but serious side effects occur: feel and/or regular heartbeats, mental mood changes such as confusion, hallucinations, nervousness, seizures, tremor, problems with toe pee.

If you are taking any monoamine oxidase inhibitor (MAOI), or two weeks after stopping them (part from medications for depression or psychological conditions may contain them). If you are not sure if your prescription contains an MAOI, consult a professional of health before taking this product.

**Ask a doctor before use if you have**

Liver disease, heart disease, high blood pressure, thyroid disease, diabetes, trouble urinating due to enlarged prostate gland, cough that occurs with too much phlegm (mucus), persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema.

**Stop use and ask a doctor if**

- Symptoms last more than 7 days
- If symptoms worsen or come back
- If you develop a headache that won't go away, fever, or skin rash.

If pregnant or breast-feeding, ask a health professional before use. Keep out of the reach of children.

**Overdose warning:** In case of accidental overdose, seek professional help or go to a Poison Control Center immediately.

**Instructions**

Follow the recommendations detailed below. Do not exceed 4 doses in a 24-hour period.

|  |                           |
|--|---------------------------|
| Adults and children 12 years and older | 10 mL, 3 or 4 times a day |
| Children between 6 to 12 years old     | 5 mL, 3 or 4 times a day  |

**Other information**

- Store at no more than 30°C (86°F)
- Do not use if security cap is altered or broken.

**This product contains alcohol and sugar.**

**Inactive ingredients**

Refined sugar, Sodium benzoate, Yellow #5 color, Orange flavor, Methyl paraben, Propyl paraben, Ethyl alcohol 95%, Sorbitol 70%, Purified water.

**Questions?** 1-661-338-6221 or visit us at [www.broncochem.com](http://www.broncochem.com)

Made in Dominican Republic  
Equipamiento para Global Corporation Inc., Boca Raton, FL 33487 USA.

**FDA REGISTERED**

**NEW!** NDC-65131-110-35

**Broncochem**  
NOCHE MAXIMUM TOS

Contiene Mepyrmino maleato, Dextrometorphan HBr

**Alivia todo tipo de tos**  
**Potente supresor de la tos para ayudarte a dormir mejor**  
**Contiene Alcohol y Azúcar**

4 FL. OZ. (120 mL ó cc)

5.157" 131mm

**Padilla**

OTIMA DE OTIMA DE LA OJEDA

A-0396611

Please recycle

Batch / Lote:

Expiration date:

8 05418 10356 5

8.802" 223.58mm

1. Empuje hacia abajo
2. Rota la tapa
3. Utilice la cuchara incluida



2.126" 54mm

**Información del Medicamento**

**Ingredientes Activos (en 5 mL ó cc)**

|                     |          |           |             |
|---------------------|----------|-----------|-------------|
| Meprobamita maleato | 50.00 mg | Propósito | Antitármico |
| Dextrometorfano HBr | 30.00 mg |           | Antitusivo  |

**Usos**

Está indicado para el tratamiento temporal en todas las formas de los ataques, subagudas y crónicas de cualquier origen, es muy eficaz en el tratamiento de la tos seca, tos de los fumadores, asma, tos nerviosa.

**Advertencias**

Los siguientes efectos adversos se pueden presentar, náusea, vómito, mareo, dolor de cabeza o resaca pueden ocurrir, si alguno de estos efectos es duradero o empeora, comuníquese con su doctor o farmacéutico inmediatamente. Evite el uso de este medicamento y comuníquese a su doctor rápidamente si usted tiene mareo, nerviosismo o problemas para comer.

**No usar**

Si durante el uso de este medicamento y comuníquese con su doctor de inmediato si alguno de estos pocos problemas pero serios efectos adversos ocurren: serios problemas de visión o cambios de visión, cambios de humor mental como confusión, alucinaciones, enrojecimiento, convulsiones, temblor, problemas al caminar.

Si está tomando algún inhibidor de la mono amino oxidasa (IMAO), o dos semanas después de haberlos suspendido (estos medicamentos para la depresión o condiciones psicológicas pueden contenerlos). Si usted no está seguro de que su receta contiene un IMAO consulte a un profesional de la salud antes de tomar este producto.

Pregunte a su médico antes de usar si usted tiene enfermedad hepática, enfermedad cardíaca, presión arterial alta, enfermedad de glándulas endocrinas, problemas para orinar, debido a agrandamiento de la próstata, tos con mucha flema (muco), los asmáticos, asma o epilepsia.

**Para el uso de este medicamento y consulte a su médico.**

- Si los síntomas duran más de 7 días
- Si los síntomas empeoran o regresan
- Si usted desarrolla un dolor de cabeza que no se quita, fiebre, o reacciones cutáneas.

Si está embarazada o en período de lactancia, consulte un profesional de la salud antes de usar.

Mantenga fuera del alcance de los niños.

**Advertencia de sobredosis:** En caso de sobredosis accidental, busque ayuda profesional o acuda a un Centro de Intoxicaciones inmediatamente.

**Instrucciones**

Siga las recomendaciones detalladas más abajo. No exceda de 4 dosis en un período de 24 horas.

|  |                           |
|--|---------------------------|
| Adultos y niños de 12 años en adelante | 10 mL, 3 o 4 veces al día |
| Niños entre 6 a 12 años                | 5 mL, 3 o 4 veces al día  |

**Información adicional**

- Contiene a no más de 30% (BPSP)
- No lo use si la tapa de seguridad está alterada o rota.

**Este producto contiene alcohol y azúcar.**

**Ingredientes Inactivos**

Azúcar refinado, Benzato de sodio, Color amarillo #5, Esencia de naranja, Metil paraben, Propil paraben, Alcohol etílico al 95%, Sorbitol al 70%, Agua purificada.

**Preguntas?** 1-661-338-6221 o visitamos [www.broncochem.com](http://www.broncochem.com)

Fabricado en República Dominicana  
Equipamiento para Global Corporation Inc., Boca Raton, FL 33487 USA.

**REGISTRADO EN LA FDA**

**NEW!** NDC-65131-110-35

**Broncochem**  
NIGHT MAXIMUM COUGH

Contains Meprobamite maleate, Dextrometorphan HBr

**Relieves cough of all origins**  
**Powerful cough suppressant to help you sleep at night**  
**Contains Alcohol & Sugar**

4 FL. OZ. (120 mL or cc)

**Drug Facts**

**Active Ingredients (in each 5 mL)**

|                     |          |         |             |
|---------------------|----------|---------|-------------|
| Meprobamite maleate | 50.00 mg | Purpose | Antitarrmic |
| Dextrometorphan HBr | 30.00 mg |         | Antitussive |

**Uses**

Temporarily relieves these symptoms in all forms of acute, subacute, and chronic cough of any origin. It is very effective in colds, rhinitis, pharyngitis, laryngitis, dry cough, smoker's cough, asthma, nervous cough.

**Warnings**

The following side effects may occur, nausea, vomiting, dizziness, headache, or nervousness may occur, if any of these effects last or worsen, contact your doctor or pharmacist immediately. Stop using this medicine and tell your doctor right away if you have dizziness, nervousness, or trouble sleeping.

**Do not use**

Stop using this medicine and contact your doctor right away if any of these unlikely but serious side effects occur: feel an irregular heartbeat, mental mood changes such as confusion, hallucinations, nervousness, seizures, tremor, problems with eye.

If you are taking any monoamine oxidase inhibitor (MAOI), or two weeks after stopping them (part from medications for depression or psychological conditions may contain them). If you are not sure if your prescription contains an MAOI, consult a professional of health before taking this product.

**Ask a doctor before use if you have**

Liver disease, heart disease, high blood pressure, thyroid disease, diabetes, trouble urinating due to enlarged prostate gland, cough that occurs with too much phlegm (mucus), persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema.

**Stop use and ask a doctor if**

- Symptoms last more than 7 days
- If symptoms worsen or come back
- If you develop a headache that won't go away, fever, or skin rash.

If pregnant or breast-feeding, ask a health professional before use. Keep out of the reach of children.

**Overdose warning:** In case of accidental overdose, seek professional help or go to a Poison Control Center immediately.

**Instructions**

Follow the recommendations detailed below. Do not exceed 4 doses in a 24-hour period.

|  |                           |
|--|---------------------------|
| Adults and children 12 years and older | 10 mL, 3 or 4 times a day |
| Children between 6 to 12 years old     | 5 mL, 3 or 4 times a day  |

**Other information**

- Store at no more than 30°C (86°F)
- Do not use if security cap is altered or broken.

**This product contains alcohol and sugar.**

**Inactive ingredients**

Refined sugar, Sodium benzoate, Yellow #5 color, Orange flavor, Methyl paraben, Propyl paraben, Ethyl alcohol 95%, Sorbitol 70%, Purified water.

**Questions?** 1-661-338-6221 or visit us at [www.broncochem.com](http://www.broncochem.com)

Made in Dominican Republic  
Equipamiento para Global Corporation Inc., Boca Raton, FL 33487 USA.

**FDA REGISTERED**

**NEW!** NDC-65131-110-35

**Broncochem**  
NOCHE MAXIMUM TOS

Contiene Meprobamite maleate, Dextrometorphan HBr

**Alivia todo tipo de tos**  
**Potente supresor de la tos para ayudarte a dormir mejor**  
**Contiene Alcohol y Azúcar**

4 FL. OZ. (120 mL ó cc)

5.157" 131mm




Batch / Lote:  
Expiration date:



8.802" 223.58mm

1. Empléjelo hasta albaño
2. Rótela la tapa
3. Utilice la cuchara incluída



2.126" 54mm

**Información del Medicamento**

**Ingredientes Activos (en 5 mL ó cc)**

|                     |          |           |               |
|---------------------|----------|-----------|---------------|
| Mepyrmino maleato   | 50.00 mg | Propósito | Antitussígeno |
| Dextrometorphan HBr | 30.00 mg |           | Antitusivo    |

**Usos**

Está indicado para el tratamiento temporal en todas las formas de los ataques, subagudas y crónicas de cualquier origen, es muy eficaz en resaca, resaca laríngea, laringitis, tos seca, tos de los fumadores, asma, tos nerviosa.

**Advertencias**

Los siguientes efectos adversos se pueden presentar, náusea, vómito, mareo, dolor de cabeza o resaca pueden ocurrir, si alguno de estos efectos es duradero o empeora, comuníquese con su doctor o farmacéutico rápidamente. Evite el uso de este medicamento y comuníquese a su doctor rápidamente si usted tiene mareo, resaca o problemas para comer.

**No usar**

Si tiene el uso de este medicamento y comuníquese con su doctor de inmediato si alguno de estos pocos problemas pero serios efectos adversos ocurren: serios problemas de visión, cambios de humor mental como confusión, alucinaciones, enrojecimiento, convulsiones, temblor, problemas al caminar.

Si está tomando algún inhibidor de la mono amino oxidasa (MAOI), o dos semanas después de haberlos suspendido (estos medicamentos para la depresión o condiciones psicológicas pueden contenerlos). Si usted no está seguro de que su receta contiene un MAOI consulte a un profesional de la salud antes de tomar este producto.

Pregunte a su médico antes de usar si usted tiene enfermedad hepática, enfermedad cardíaca, presión arterial alta, enfermedad de glándulas endocrinas, problemas para orinar, debido a agrandamiento de la próstata, tos con mucha flema (muco), los asmáticos, asma o enfisema.

**Para el uso de este medicamento y consulte a su médico.**

- Si los síntomas duran más de 7 días
- Si los síntomas empeoran o regresan
- Si usted desarrolla un dolor de cabeza que no se quita, fiebre, o reacciones cutáneas.

Si está embarazada o en período de lactancia, consulte un profesional de la salud antes de usar.

Manténgala fuera del alcance de los niños.

**Advertencia de sobredosis:** En caso de sobredosis accidental, busque ayuda profesional o acuda a un Centro de Intoxicaciones inmediatamente.

**Instrucciones**

Siga las recomendaciones detalladas más abajo. No exceda de 4 dosis en un período de 24 horas.

|  |                           |
|--|---------------------------|
| Adultos y niños de 12 años en adelante | 10 mL, 3 o 4 veces al día |
| Niños entre 6 a 12 años                | 5 mL, 3 o 4 veces al día  |

**Información adicional**

- Contiene a no más de 30% (BPFF)
- No lo use si la tapa de seguridad está alterada o rota.

**Este producto contiene alcohol y azúcar.**

**Ingredientes Inactivos**

Azúcar refinada, Benzato de sodio, Color amarillo #5, Esencia de naranja, Metil paraben, Propil paraben, Alcohol etílico al 95%, Sorbitol al 70%, Agua purificada.

**Preguntas?** 1-661-338-6221 o visitamos [www.broncochem.com](http://www.broncochem.com)

Fabricado en República Dominicana  
Equipamiento para Global Corporation Inc., Boca Raton, FL 33487 USA.

**REGISTRADO EN LA FDA**

**NEW!** NDC-65131-110-35

**Broncochem**  
NIGHT MAXIMUM COUGH

Contains Mepyrmino maleate, Dextrometorphan HBr

**Relieves cough of all origins**  
**Powerful cough suppressant to help you sleep at night**  
**Contains Alcohol & Sugar**

4 FL. OZ. (120 mL or cc)

**Drug Facts**

**Active Ingredients (in each 5 mL)**

|                     |          |         |             |
|---------------------|----------|---------|-------------|
| Mepyrmino maleate   | 50.00 mg | Purpose | Antitussive |
| Dextrometorphan HBr | 30.00 mg |         | Antitussive |

**Uses**

Temporarily relieves these symptoms in all forms of acute, subacute, and chronic cough of any origin. It is very effective in colds, rhinitis, pharyngitis, laryngitis, dry cough, smoker's cough, asthma, nervous cough.

**Warnings**

The following side effects may occur, nausea, vomiting, dizziness, headache, or nervousness may occur, if any of these effects last or worsen, contact your doctor or pharmacist immediately. Stop using this medicine and tell your doctor right away if you have dizziness, nervousness, or trouble sleeping.

**Do not use**

Stop using this medicine and contact your doctor right away if any of these unlikely but serious side effects occur: feel an irregular heartbeat, mental mood changes such as confusion, hallucinations, nervousness, seizures, tremor, problems with eye.

If you are taking any monoamine oxidase inhibitor (MAOI), or two weeks after stopping them (part from medications for depression or psychological conditions may contain them). If you are not sure if your prescription contains an MAOI, consult a professional of health before taking this product.

**Ask a doctor before use if you have**

Liver disease, heart disease, high blood pressure, thyroid disease, diabetes, trouble urinating due to enlarged prostate gland, cough that occurs with too much phlegm (mucus), persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema.

**Stop use and ask a doctor if**

- Symptoms last more than 7 days
- If symptoms worsen or come back
- If you develop a headache that won't go away, fever, or skin rash.

If pregnant or breast-feeding, ask a health professional before use. Keep out of the reach of children.

**Overdose warning:** In case of accidental overdose, seek professional help or go to a Poison Control Center immediately.

**Instructions**

Follow the recommendations detailed below. Do not exceed 4 doses in a 24-hour period.

|  |                           |
|--|---------------------------|
| Adults and children 12 years and older | 10 mL, 3 or 4 times a day |
| Children between 6 to 12 years old     | 5 mL, 3 or 4 times a day  |

**Other information**

- Store at no more than 30°C (86°F)
- Do not use if security cap is altered or broken.

**This product contains alcohol and sugar.**

**Inactive ingredients**

Refined sugar, Sodium benzoate, Yellow #5 color, Orange flavor, Methyl paraben, Propyl paraben, Ethyl alcohol 95%, Sorbitol 70%, Purified water.

**Questions?** 1-661-338-6221 or visit us at [www.broncochem.com](http://www.broncochem.com)

Made in Dominican Republic  
Equipamiento para Global Corporation Inc., Boca Raton, FL 33487 USA.

**FDA REGISTERED**

**NEW!** NDC-65131-110-35


**Broncochem**  
NOCHE MAXIMUM TOS

Contiene Mepyrmino maleato, Dextrometorphan HBr

**Alivia todo tipo de tos**  
**Potente supresor de la tos para ayudarte a dormir mejor**  
**Contiene Alcohol y Azúcar**

4 FL. OZ. (120 mL ó cc)

5.157" 131mm



Batch / Lote:  
Expiration date:



**Warnings**

■ The following side effects may occur, nausea, vomiting, dizziness, headache, or nervousness may occur, if any of these effects last or worsen, contact your doctor or pharmacist immediately. Stop using this medicine and tell your doctor right away if you have dizziness, nervousness, or trouble sleeping.

**Stop use and ask a doctor if**

- Symptoms last more than 7 days
- If symptoms worsen or come back.
- If you develop a headache that won't go away, fever, or skin rash.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of the reach of children.

**Overdose warning:** In case of accidental overdose, seek professional help or go to a Poison Control Center immediately.

**Other information** ■ Store at no more than 30°C (86°F) ■ Do not use if security cap is altered or broken

**This product contains alcohol and sugar.**

**Inactive ingredients** Refined sugar, Sodium benzoate, Yellow #5 color, Orange flavor, Methyl paraben, Propyl paraben, Ethyl alcohol 95%, Sorbitol 70%, Purified water.

NDC-65131-110-35

# Broncochem®

## MAXIMUM COUGH NIGHT

Relieves cough of all origins  
 Powerful cough suppressant to help you sleep at night  
 Contains Alcohol & Sugar

4 FL. OZ. (120mL or so)

Made in Dominican Republic  
Exclusively for: Global Corporation Inc.,  
Boca Raton, FL 33487 U.S.A.

FDA REGISTERED

Lote / Batch:  
Expiration date:

**Active ingredients (in each 10 mL)**

|                      |          |               |
|----------------------|----------|---------------|
| Mepyramine maleate   | 50.00 mg | Antihistamine |
| Dextromethorphan HBr | 30.00 mg | Antitussive   |

**Purpose**

**Uses** Temporarily relieves these symptoms in all forms of acute, subacute, and conical cough of any origin, it is very effective in colds, rhinitis, pharyngitis, laryngitis, dry cough, smokers' cough, asthma, nervous cough.

**Ask a doctor before use if you have**

Liver disease, heart disease, high blood pressure, thyroid disease, diabetes, trouble urinating due to enlarged prostate gland, cough that occurs with too much phlegm (mucus), persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema.

**Directions** Take only as directed. For your convenience, a dosage spoon is provide. **Do not exceed 4 doses in a 24 hour period.**

|  |                          |
|--|--------------------------|
| Adults and children 12 years and older | 10 mL 3 or 4 times a day |
|--|--------------------------|

|                                    |                         |
|------------------------------------|-------------------------|
| Children between 6 to 12 years old | 5 mL 3 or 4 times a day |
|------------------------------------|-------------------------|

See carton for complete information for use.

**Questions?** Please call for medical direction weekdays 9 AM to 5 PM EST. at 1-561-338-5221 or visit us at [www.broncochem.com](http://www.broncochem.com)

**Warnings**

■ The following side effects may occur, nausea, vomiting, dizziness, headache, or nervousness may occur, if any of these effects last or worsen, contact your doctor or pharmacist immediately. Stop using this medicine and tell your doctor right away if you have dizziness, nervousness, or trouble sleeping.

**Stop use and ask a doctor if**

- Symptoms last more than 7 days
- If symptoms worsen or come back.
- If you develop a headache that won't go away, fever, or skin rash.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of the reach of children.

**Overdose warning:** In case of accidental overdose, seek professional help or go to a Poison Control Center immediately.

**Other information** ■ Store at no more than 30°C (86°F) ■ Do not use if security cap is altered or broken

**This product contains alcohol and sugar.**

**Inactive ingredients** Refined sugar, Sodium benzoate, Yellow #5 color, Orange flavor, Methyl paraben, Propyl paraben, Ethyl alcohol 95%, Sorbitol 70%, Purified water.

NDC-65131-110-35

# Broncochem®

## MAXIMUM COUGH NIGHT

Relieves cough of all origins  
 Powerful cough suppressant to help you sleep at night  
 Contains Alcohol & Sugar

4 FL. OZ. (120mL or so)

Made in Dominican Republic  
Exclusively for: Global Corporation Inc.,  
Boca Raton, FL 33487 U.S.A.

FDA REGISTERED

Lote / Batch:  
Expiration date:

**Active ingredients (in each 10 mL)**

|                      |          |               |
|----------------------|----------|---------------|
| Mepyramine maleate   | 50.00 mg | Antihistamine |
| Dextromethorphan HBr | 30.00 mg | Antitussive   |

**Purpose**

**Uses** Temporarily relieves these symptoms in all forms of acute, subacute, and conical cough of any origin, it is very effective in colds, rhinitis, pharyngitis, laryngitis, dry cough, smokers' cough, asthma, nervous cough.

**Ask a doctor before use if you have**

Liver disease, heart disease, high blood pressure, thyroid disease, diabetes, trouble urinating due to enlarged prostate gland, cough that occurs with too much phlegm (mucus), persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema.

**Directions** Take only as directed. For your convenience, a dosage spoon is provide. **Do not exceed 4 doses in a 24 hour period.**

|  |                          |
|--|--------------------------|
| Adults and children 12 years and older | 10 mL 3 or 4 times a day |
|--|--------------------------|

|                                    |                         |
|------------------------------------|-------------------------|
| Children between 6 to 12 years old | 5 mL 3 or 4 times a day |
|------------------------------------|-------------------------|

See carton for complete information for use.

**Questions?** Please call for medical direction weekdays 9 AM to 5 PM EST. at 1-561-338-5221 or visit us at [www.broncochem.com](http://www.broncochem.com)

NDC-65131-110-35

# Broncochem®

## MAXIMUM COUGH NIGHT

**Relieves cough of all origins**  
**Powerful cough suppressant to help you sleep at night**  
**Contains Alcohol & Sugar**

|  |  |   |   |                    |          |               |                      |          |             |  |                          |                                    |                         |
|--|--|---|---|--------------------|----------|---------------|----------------------|----------|-------------|--|--------------------------|------------------------------------|-------------------------|
| <p><b>Warnings</b></p> <ul style="list-style-type: none"> <li>■ The following side effects may occur, nausea, vomiting, dizziness, headache, or nervousness may occur, if any of these effects last or worsen, contact your doctor or pharmacist immediately. Stop using this medicine and tell your doctor right away if you have dizziness, nervousness, or trouble sleeping.</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>■ Symptoms last more than 7 days</li> <li>■ If symptoms worsen or come back.</li> <li>■ If you develop a headache that won't go away, fever, or skin rash.</li> </ul> <p><b>If pregnant or breast-feeding</b>, ask a health professional before use.</p> <p><b>Keep out of the reach of children.</b></p> <p><b>Overdose warning:</b> In case of accidental overdose, seek professional help or go to a Poison Control Center immediately.</p> <p><b>Other information</b> ■ Store at no more than 30°C (86°F) ■ Do not use if security cap is altered or broken</p> <p><b>This product contains alcohol and sugar.</b></p> <p><b>Inactive ingredients</b> Refined sugar, Sodium benzoate, Yellow #5 color, Orange flavor, Methyl paraben, Propyl paraben, Ethyl alcohol 95%, Sorbitol 70%, Purified water.</p> | <p><b>4 FL. OZ. (120 mL or so)</b></p> <p><b>Made in Dominican Republic</b><br/>         Exclusively for: Global Corporation Inc., Boca Raton, FL 33487 U.S.A.</p> | <p><b>FDA REGISTERED</b></p> <p>Lote / Batch:<br/>         Expiration date:</p> | <p><b>Active ingredients (in each 10 mL)</b></p> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 60%;">Mepyramine maleate</td> <td style="width: 20%;">50.00 mg</td> <td style="width: 20%;">Antihistamine</td> </tr> <tr> <td>Dextromethorphan HBr</td> <td>30.00 mg</td> <td>Antitussive</td> </tr> </table> <p><b>Purpose</b></p> <p><b>Uses</b> Temporarily relieves these symptoms in all forms of acute, subacute, and chronic cough of any origin, it is very effective in colds, rhinitis, pharyngitis, laryngitis, dry cough, smokers' cough, asthma, nervous cough.</p> <p><b>Ask a doctor before use if you have</b><br/>         Liver disease, heart disease, high blood pressure, thyroid disease, diabetes, trouble urinating due to enlarged prostate gland, cough that occurs with too much phlegm (mucus), persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema.</p> <p><b>Directions</b> Take only as directed. For your convenience, a dosage spoon is provided. <b>Do not exceed 4 doses in a 24 hour period.</b></p> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 60%;">Adults and children 12 years and older</td> <td style="width: 40%;">10 mL 3 or 4 times a day</td> </tr> <tr> <td>Children between 6 to 12 years old</td> <td>5 mL 3 or 4 times a day</td> </tr> </table> <p>See carton for complete information for use.</p> <p><b>Questions?</b> Please call for medical direction weekdays 9 AM to 5 PM EST. at <b>1-561-338-5221</b> or visit us at <a href="http://www.broncochem.com">www.broncochem.com</a></p> | Mepyramine maleate | 50.00 mg | Antihistamine | Dextromethorphan HBr | 30.00 mg | Antitussive | Adults and children 12 years and older | 10 mL 3 or 4 times a day | Children between 6 to 12 years old | 5 mL 3 or 4 times a day |
| Mepyramine maleate   | 50.00 mg   | Antihistamine   |   |                    |          |               |                      |          |             |  |                          |                                    |                         |
| Dextromethorphan HBr   | 30.00 mg   | Antitussive   |   |                    |          |               |                      |          |             |  |                          |                                    |                         |
| Adults and children 12 years and older   | 10 mL 3 or 4 times a day   |   |   |                    |          |               |                      |          |             |  |                          |                                    |                         |
| Children between 6 to 12 years old   | 5 mL 3 or 4 times a day  |   |   |                    |          |               |                      |          |             |  |                          |                                    |                         |

## BRONCOCHEM MAXIMUM COUGH NIGHT

dextromethorphan hbr-pyrilamine maleate syrup

### Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:65131-110 |
| <b>Route of Administration</b> | ORAL           |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength                | Strength         |
|---|----------------------------------|------------------|
| <b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH)<br>(DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN<br>HYDROBROMIDE | 30 mg<br>in 5 mL |
| <b>PYRILAMINE MALEATE</b> (UNII: R35D29L3ZA) (PYRILAMINE - UNII:HPE317O9TL)                     | PYRILAMINE MALEATE               | 50 mg<br>in 5 mL |

### Inactive Ingredients

| Ingredient Name                                 | Strength         |
|---|------------------|
| <b>SUGARCANE</b> (UNII: 81H2R5AOH3)             | 2.65 mg in 5 mL  |
| <b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)       | 10 mg in 5 mL    |
| <b>FD&amp;C YELLOW NO. 5</b> (UNII: I753WB2F1M) | 0.25 mg in 5 mL  |
| <b>ORANGE OIL</b> (UNII: AKN3KSD11B)            | 0.833 mg in 5 mL |
| <b>METHYL PARABEN SODIUM</b> (UNII: CR6K9C2NHK) | 5 mg in 5 mL     |
| <b>PROPYL PARABEN SODIUM</b> (UNII: 625NNB0G9N) | 1.5 mg in 5 mL   |
| <b>ALCOHOL 95%</b> (UNII: 7528N5H79B)           | 0.25 mL in 5 mL  |
| <b>SORBITOL SOLUTION 70%</b> (UNII: 8KW3E207O2) | 0.083 mL in 5 mL |
| <b>water</b> (UNII: 059QF0K00R)                 | 5 mL in 5 mL     |

### Packaging

| # | Item Code        | Package Description  | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:65131-110-35 | 1 in 1 BOX   | 04/21/2025           |                    |
| 1 |                  | 120 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product |                      |                    |

| Marketing Information |  |                      |                    |
|-----------------------|--|----------------------|--------------------|
| Marketing Category    | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| OTC Monograph Drug    | M  | 04/21/2025           |                    |

**Labeler** - LABORATORIO MAGNACHEM INTERNATIONAL SRL (871446100)

**Registrant** - LABORATORIO MAGNACHEM INTERNATIONAL SRL (871446100)

| Establishment                           |         |           |                        |
|---|---------|-----------|------------------------|
| Name                                    | Address | ID/FEI    | Business Operations    |
| LABORATORIO MAGNACHEM INTERNATIONAL SRL |         | 871446100 | manufacture(65131-110) |

Revised: 3/2025

LABORATORIO MAGNACHEM INTERNATIONAL SRL