

**SATO CLEAR- naphazoline hydrochloride solution/ drops**  
**Sato Pharmaceutical Co., Ltd.**

-----  
**Sato Clear**

**Active ingredient**

Naphazoline hydrochloride 0.012%

**Purpose**

Naphazoline hydrochloride     Vasoconstrictor

**Uses** relieves this symptom due to minor eye irritations

■ redness of the eye

**Warnings**

**For external use only**

**Do not use**

- if solution changes color or becomes cloudy

**Ask a doctor before use if you have**

- narrow angle glaucoma

**When using this product**

- pupils may become enlarged temporarily
- overuse of this product may produce increased redness of the eyes
- to avoid contamination, do not touch tip of container to any surface
- replace cap after using

**Stop use and ask a doctor if you experience**

- eye pain
- changes in vision
- continued redness or irritation of the eye
- the condition worsens or persists for more than 72 hours

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

■ Place 1 or 2 drops in the affected eyes up to 4 times daily.

**Other information**

- use only if imprinted tamper-resistant seal on the bottle cap is intact
- keep container tightly closed

**Inactive ingredients** benzalkonium chloride, boric acid, dl-camphor, sodium borate, sodium chloride, purified water.

# SATO CLEAR

★ Use only if imprinted carton overwrap with "サトウクリア SATO CLEAR SAFETY SEAL" is intact at time of purchase.  
 ★ "サトウクリア SATO CLEAR SAFETY SEAL" と印刷されたオーパレーションフィルムが、完全な場合にのみ使用してください。

## Drug Facts

**Active ingredient** Naphazoline hydrochloride 0.012%.....**Purpose** Vasoconstrictor

**Uses** relieves the symptom due to minor eye irritations  
 ■ redness of the eye

### Warnings

For external use only

#### Do not use

■ if solution changes color or becomes cloudy

Ask a doctor before use if you have

■ narrow angle glaucoma

When using this product

■ pupils may become enlarged temporarily  
 ■ overuse of this product may produce increased redness of the eyes

■ to avoid contamination, do not touch tip of container to any surface

■ replace cap after using

Stop use and ask a doctor if you experience

■ eye pain ■ changes in vision  
 ■ continued redness or irritation of the eye  
 ■ the condition worsens or persists for more than 72 hours

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

### Directions

■ place 1 or 2 drops in the affected eyes up to 4 times daily.

### Other information

■ use only if imprinted tamper-resistant seal on the bottle cap is intact.  
 ■ keep container tightly closed

**Inactive ingredients** benzalkonium chloride, boric acid, dl-camphor, sodium borate, sodium chloride, purified water.

Manufactured by SATO PHARMACEUTICAL CO., LTD.  
 1-5-27 MOTOAKASAKA MINATO-KU TOKYO, JAPAN  
 Reports of serious side effects associated with use of this product can be sent to:  
 SATO PHARMACEUTICAL, INC. 1650 E. Glenn Curiss St., Unit 2, Carson, CA 90746  
 製造元 佐藤製薬株式会社 日本国東京都港区元赤坂1丁目5番27号

医薬品表示(警告)  
 の病気があるときは、重篤に悪化が生じたとき、目の充血や刺激の改善がみられず、場合によっては  
 症状がひどくなる場合は使用を中止し、医師にご相談ください。■ 小児の手のとどかない所に保管して  
 ください。■ 目を洗うときは、目を閉じて洗い、すすぎを繰り返してください。■ 容器のキャップを  
 使用しないでください。■ キヤップに装着されている印刷付き改ざん防止シールが不完全な場合は、使  
 用しないでください。■ キヤップをしっかりと閉めて保管してください。  
 用法・用量 ■ 1回1〜2滴ずつ1日4回を限度として点眼します。  
 その他の情報 ■ 本品は、ベンザルコニウム、ホウ酸、非カチオン性ポリマー、酸化チタン、精製水  
 その他の成分 ■ 塩化ベンザルコニウム、ホウ酸、非カチオン性ポリマー、酸化チタン、精製水

sato NDC49873-044-01

# SATO CLEAR

サトウクリアー  
 充血用点眼薬

Redness Reliever  
 Eye Drops

Naphazoline  
 Hydrochloride  
 ナファゾリン塩酸塩



0.5 FL OZ (15 mL)

配合目的 血管収縮剤  
 有効成分・分量 ナファゾリン 0.012%  
 効能 ■目の炎症による次の症状の緩和:目の充血  
 使用上の注意 外用にのみ使用下さい。■ 薬液が変色したり、混濁したものは使用しないで  
 ください。■ 結膜内障の場合には使用せず、医師にご相談ください。■ 使用により一時的に瞳孔が大き  
 くなる場合があります。■ 過度に使用しますと、かえって充血がひどくなる場合があります。■ 汚染防止  
 のため、容器の先端に絶対に触れないでください。■ 使用後は必ずキャップを閉めてください。■ はげしい目

EXP. DATE



LOT

US-K

## SATO CLEAR

naphazoline hydrochloride solution/ drops

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:49873-044
<b>Route of Administration</b>	OPHTHALMIC		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>NAPHAZOLINE HYDROCHLORIDE</b> (UNII: MZ1131787D) (NAPHAZOLINE - UNII:H231GF11BV)	NAPHAZOLINE HYDROCHLORIDE	12 mg in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>CAMPHOR (SYNTHETIC)</b> (UNII: 5TJD82A1ET)	
<b>BENZALKONIUM CHLORIDE</b> (UNII: F5UM2KM3W7)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>BORIC ACID</b> (UNII: R57ZHV85D4)	
<b>SODIUM BORATE</b> (UNII: 91MBZ8H3QO)	
<b>WATER</b> (UNII: 059QF0KO0R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49873-044-01	1 in 1 CARTON	05/31/1989	
1		15 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M018	05/31/1989	

**Labeler** - Sato Pharmaceutical Co., Ltd. (690575642)

### Establishment

Name	Address	ID/FEI	Business Operations
Sato Pharmaceutical Co., Ltd.		715699133	manufacture(49873-044) , label(49873-044) , pack(49873-044)

Revised: 11/2023

Sato Pharmaceutical Co., Ltd.