

**QUALITY CHOICE STOMACH RELIEF 528 - bismuth subsalicylate liquid**  
**Chain Drug Marketing Association Inc.**

-----  
**QCH Ultra Strength Stomach Relief 528**

**ACTIVE INGREDIENT(in each 15 mL)**

Bismuth subsalicylate 525 mg

**PURPOSE**

Upset stomach reliever and anti-diarrheal

**USE(S)**

relieves:

- diarrhea
- heartburn
- indigestion
- nausea
- upset stomach associated with these symptoms

**WARNINGS**

**Reye's Syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**DO NOT USE IF YOU HAVE**

- an ulcer
- bloody or black stool
- a bleeding problem

**ASK A DOCTOR BEFORE USE IF YOU HAVE**

- fever

- mucus in the stool

## **ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE**

taking any drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

## **WHEN USING THIS PRODUCT**

a temporary, but harmless darkening of the stool and/or tongue may occur

## **STOP USE AND ASK DOCTOR IF**

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

## **IF PREGNANT OR BREAST-FEEDING,**

ask a health professional before use.

## **KEEP OUT OF REACH OF CHILDREN**

In case of overdose, get medical help or contact a Poison Control Center immediately.

## **DIRECTIONS**

- **shake well before use**
- mL = milliliter
- TBSP = tablespoon
- adults and children 12 years and over: 1 dose (2 TBSP or 15 mL) every 1/2 hour as needed
- do not exceed 8 doses (120 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of fluids to help prevent dehydration caused by diarrhea

## **OTHER INFORMATION**

- **each 15 mL or 1 TBSP contains:**
- potassium 15 mg
- salicylate 236 mg
- sodium 5 mg
- protect from freezing

- avoid excessive heat (over 104°F or 40°C)
- dosage cup provided

### **INACTIVE INGREDIENTS**

benzoic acid, D&C red # 22, D&C red # 28, flavor, hydroxyethyl cellulose, potassium hydroxide, purified water, saccharin sodium, salicylic acid, simethicone, xanthan gum.

### **PRINCIPAL DISPLAY PANEL**

NDC 83324-021-08

### **QUALITY CHOICE**

**\*compare to the active ingredient in Pepto-Bismol® Ultra**

**Ultra Strength**

**Stomach Relief**

**Upset Stomach Reliever/Anti-diarrheal  
bismuth subsalicylate 525 mg per 15 mL**

#### **Relieves:**

- Diarrhea
- Heartburn
- Indigestion
- Nausea & Upset Stomach

### **Original Flavor**

8 FL OZ (236 mL)

**Drug Facts** DO NOT USE IF IMPRINTED SHRINKBAND IS MISSING OR BROKEN.

**Active ingredient (per 15 mL)** **Purpose**  
 Bismuth subsalicylate 525 mg .....Upset stomach reliever and anti-diarrheal

**Uses** relieves: ■ diarrhea ■ heartburn ■ indigestion ■ nausea ■ upset stomach associated with these symptoms

**Warnings**  
 Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Contains salicylate. Do not take if you are ■ allergic to salicylates (including aspirin) ■ taking other salicylate products  
**Do not use if you have** ■ an ulcer ■ bloody or black stool ■ a bleeding problem  
**Ask a doctor before use if you have** ■ fever ■ mucus in the stool  
**Ask a doctor or pharmacist before use if you are taking any drug for** ■ anticoagulation (thinning of the blood) ■ diabetes ■ gout ■ arthritis  
**When using this product** a temporary, but harmless darkening of the stool and/or tongue may occur  
**Stop use and ask a doctor if** ■ symptoms get worse ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days  
**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center immediately. ◀ Drug Facts continued



NDC 83324-021-08

\*Compare to the Active Ingredient in Pepto-Bismol® Ultra

## Ultra Strength Stomach Relief

Upset Stomach Reliever/Anti-diarrheal

Bismuth Subsalsicylate | 525 mg Per 15 mL

Relieves:  
 Diarrhea, Heartburn, Indigestion, Nausea & Upset Stomach

Original Flavor

**8 FL OZ (236 mL)**

**Drug Facts (continued)**

**Directions** ■ shake well before use  
 ■ mL = milliliter ■ TBSP = tablespoon ■ adults and children 12 years and over: 1 dose (1 TBSP or 15 mL) every 1/2 hour as needed ■ do not exceed 8 doses (120 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information** ■ each 15 mL or 1 TBSP contains:  
 • potassium 15 mg • salicylate 236 mg • sodium 5 mg  
 ■ protect from freezing. ■ avoid excessive heat (over 104°F or 40°C). ■ dosage cup provided

**Inactive ingredients** benzoic acid, D&C red # 22, D&C red # 28, flavor, hydroxyethyl cellulose, potassium hydroxide, purified water, saccharin sodium, salicylic acid, simethicone, xanthan gum.

\*This product is not manufactured or distributed by Procter & Gamble, Inc., the distributor of Pepto-Bismol® Ultra Strength.

Distributed by CDMA, Inc.  
 Novi, MI 48375  
 www.qualitychoice.com  
 Questions: 800-935-2362



6 35515 98132 4

MF# 53041 REV 0324

# QUALITY CHOICE STOMACH RELIEF 528

bismuth subsalicylate liquid

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:83324-021
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 15 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>BENZOIC ACID</b> (UNII: 8SKN0B0MIM)	
<b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)	
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)	
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ)	

**HYDROXYETHYL CELLULOSE (1500 MPA.S AT 1%)** (UNII: L605B5892V)

**DIMETHICONE** (UNII: 92RU3N3Y1O)

**XANTHAN GUM** (UNII: TTV12P4NEE)

### Product Characteristics

<b>Color</b>	PINK	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83324-021-08	236 mL in 1 BOTTLE; Type 0: Not a Combination Product	04/19/2024	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	04/19/2024	

**Labeler** - Chain Drug Marketing Association Inc. (011920774)

Revised: 1/2025

Chain Drug Marketing Association Inc.