

**ED A-HIST DM- dextromethorphan hydrobromide, chlorpheniramine maleate,  
and phenylephrine hydrochloride liquid  
EDWARDS PHARMACEUTICALS, INC.**

-----  
**ED-EDA-HISTDM 0171**

***Drug Facts***

<b><i>Active Ingredients (in each 5 mL teaspoonful)</i></b>	<b><i>Purpose</i></b>
Chlorpheniramine Maleate 4 mg	Antihistamine
Dextromethorphan HBr 15 mg	Antitussive
Phenylephrine HCl 10 mg	Nasal Decongestant

**Uses**

temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- cough due to minor throat and bronchial irritation
- nasal congestion
- reduces swelling of nasal passages

**Warnings**

**Do not exceed recommended dosage.**

**Do not use this product**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Do not use this product, unless directed by a doctor, if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- a persistent or chronic cough that occurs with too much phlegm (mucus)
- heart disease
- high blood pressure
- thyroid disease
- diabetes mellitus
- difficulty in urination due to enlargement of the prostate gland

**Do not take this product if you are taking sedatives or tranquilizers, without**

## **first consulting your doctor.**

### **When using this product**

- excitability may occur, especially in children
- may cause marked drowsiness
- sedatives and tranquilizers may increase drowsiness effect
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

### **Stop use and ask doctor if**

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. These could be signs of a serious condition.
- new symptoms occur

**If pregnant or breastfeeding,** ask a health professional before use.

### **Keep out of reach of children.**

In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

## **Directions**

### **Do not exceed recommended dosage**

Adults and children 12 years of age and over:	1 teaspoonful (5 mL) every 4 hours, not to exceed 6 teaspoonfuls in 24 hours, or as directed by a doctor.
Children 6 to under 12 years of age:	1/2 teaspoonful (2.5 mL) every 4 hours, not to exceed 3 teaspoonfuls in 24 hours, or as directed by a doctor.
Children under 6 years of age:	Consult a doctor.

## **Other information**

Store at 59° - 86° F (15° - 30° C) [see USP for Controlled Room Temperature].

## **Inactive ingredients**

Artificial and Natural Banana Flavor, Citric Acid, FD&C Red #40, FD&C Yellow #6, Methyl Paraben, Propyl Paraben, Propylene Glycol, Purified Water, Sodium Citrate, Sucralose.

**Question? Comments?**

Call 1-800-543-9560

**PRINCIPAL DISPLAY PANEL - 473 mL Bottle Label**

**E**

NDC 0485-0171-16

**ED A-HIST DM**

**Antihistamine • Antitussive  
• Nasal Decongestant**

**Sugar Free • Gluten Free**

**Each teaspoonful (5 mL)  
for oral administration contains:**

Chlorpheniramine Maleate 4 mg

Dextromethorphan HBr 15 mg

Phenylephrine HCl 10 mg

**Banana Flavor**

Tamper evident by foil seal under cap.

Do not use if foil seal is broken or  
missing.

**Manufactured for:**

***EDWARDS***

**Pharmaceuticals, Inc.**

**Ripley, MS 38663**

**16oz. (473 mL)**



NDC 0485-0171-16

### ED A-HIST DM

Antihistamine • Antitussive  
• Nasal Decongestant

Sugar Free • Gluten Free

Each teaspoonful (5 mL)

for oral administration contains:

Chlorpheniramine Maleate .....4 mg

Dextromethorphan HBr.....15 mg

Phenylephrine HCl.....10 mg

Banana Flavor

Tamper evident by foil seal under cap.  
Do not use if foil seal is broken or missing.

Manufactured for:  
**EDWARDS**  
Pharmaceuticals, Inc.  
Berwyn, PA 19312

16 oz (473 mL)

ED A-HIST DM

#### Drug Facts

##### Active Ingredients (in each 5 mL teaspoonful)

Chlorpheniramine Maleate 4 mg ..... Antihistamine  
Dextromethorphan HBr 15 mg ..... Antitussive  
Phenylephrine HCl 10 mg ..... Nasal Decongestant

**Uses** temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other respiratory allergies: ■ runny nose ■ sneezing  
■ itching of the nose or throat ■ itchy, watery eyes  
■ cough due to minor throat and bronchial irritation  
■ nasal congestion ■ reduces swelling of nasal passages

##### Warnings

**Do not exceed recommended dosage.**

**Do not use this product**

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Do not use this product, unless directed by a doctor, if you have**

■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ a persistent or chronic cough that occurs with too much phlegm (mucus) ■ heart disease  
■ high blood pressure ■ thyroid disease ■ diabetes mellitus  
■ difficulty in urination due to enlargement of the prostate gland

**Do not take this product if you are taking sedatives or tranquilizers, without first consulting your doctor.**

**When using this product**

■ excitability may occur, especially in children  
■ may cause marked drowsiness  
■ sedatives and tranquilizers may increase drowsiness effect  
■ avoid alcoholic beverages  
■ use caution when driving a motor vehicle or operating machinery

##### Purpose

#### Drug Facts (continued)

**Stop use and ask doctor if**

■ nervousness, dizziness, or sleeplessness occur  
■ cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. These could be signs of a serious condition. ■ new symptoms occur

**If pregnant or breastfeeding**, ask a health professional before use.

**Keep out of reach of children.**

In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

#### Directions

**Do not exceed recommended dosage**

Adults and children 12 years of age and over: 1 teaspoonful (5 mL) every 4 hours, not to exceed 6 teaspoonfuls in 24 hours, or as directed by a doctor.

Children 6 to under 12 years of age: 1/2 teaspoonful (2.5 mL) every 4 hours, not to exceed 3 teaspoonfuls in 24 hours, or as directed by a doctor.

Children under 6 years of age: Consult a doctor.

#### Other information

Store at 59° - 86° F (15° - 30° C) [see USP for Controlled Room Temperature].

#### Inactive ingredients

Artificial and Natural Banana Flavor, Citric Acid, FD&C Red #40, FD&C Yellow #6, Methylparaben, Propylene Glycol, Propylparaben, Purified Water, Sodium Citrate, Sucralose.

**Question? Comments?** Call 1-800-543-9560

Rev. 06/22 L-201



Lot#  
Exp. Date:

3 04850 17116 4

## ED A-HIST DM

dextromethorphan hydrobromide, chlorpheniramine maleate, and phenylephrine hydrochloride liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0485-0171
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg in 5 mL
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	4 mg in 5 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	

SUCRALOSE (UNII: 96K6UQ3ZD4)

CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)

### Product Characteristics

Color	yellow	Score	
Shape		Size	
Flavor	BANANA	Imprint Code	
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0485-0171-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/11/2012	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/11/2012	

**Labeler** - EDWARDS PHARMACEUTICALS, INC. (195118880)

Revised: 12/2025

EDWARDS PHARMACEUTICALS, INC.