

**ANADENT KANKA GEL- benzocaine gel**  
**BELMORA LLC**

-----  
**AnaDent Kanka Gel**

**Drug Facts**

**Active Ingredient**

Benzocaine 20%

**Purpose**

oral pain reliever

**Uses**

- For the temporary relief of pain due to toothache
- To help protect against infection of minor oral irritation.

**Warnings**

**Methemoglobinemia warning:** use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy alert:** Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics

**Do not use**

- more than directed
- for more than 7 days unless told to do so by a dentist or doctor
- for teething
- in children under 2 years of age

**Stop use and ask a dentist or doctor if**

- swelling, rash or fever develops
- irritation, pain or redness persists or worsens
- symptoms do not improve in 7 days
- allergic reaction occurs

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- Cut open tip of tube on score mark

- Adults and children 2 years of age and older: apply a small amount of product to the cavity and around the gum surrounding the teeth. Use up to 4 times daily or as directed by a dentist or doctor
- Children under 12 years of age should be supervised in the use of this product
- Children under 2 years of age: do not use

### ***Other Information***

- Do not use if tip is cut prior to opening
- This preparation is intended for use in cases of toothache, only as a temporary expedient until a dentist can be consulted
- Do not use continuously
- This formula will stay in place for extended duration of relief
- Avoid using toothpaste or drinking soft drinks or fruit juices for at least one hour after applying

### ***Inactive Ingredients:***

Ammonium Glycyrrhizate, Flavor, PEG-400, PEG-3350, Sodium Saccharin Dihydrate, Sorbic Acid

20% BENZOCAINE TO RELIEVE ORAL PAIN

COOLS GUM & RELIEVES IRRITATION

**For toothaches**

**Toothache and Gum pain Relief Gel**

**TAMPER-EVIDENT:**

Do not use if tube is cut prior to use.  
Retain carton for complete product information.

Distributed by Belmora LLC, 3033 Wilson Blvd. Suite 700 Arlington, VA 22201 |  
[www.anadentusa.com](http://www.anadentusa.com)

Questions/Comments please call: 1 888 779 2877 Monday to Friday 9AM - 5PM EST

**Packaging**



**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:27854-177-01	1 in 1 CARTON	04/15/2022	
1		7 g in 1 TUBE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	04/15/2022	

**Labeler** - BELMORA LLC (112753244)

Revised: 10/2024

BELMORA LLC