

**MARKET BASKET EFFERVESCENT ANTACID AND PAIN RELIEF- aspirin, citric acid, sodium bicarbonate tablet, effervescent**  
**Demoulas Super Markets, Inc.**

-----  
**Market Basket Effervescent Antacid & Pain Relief**

**Active ingredients**

**(in each effervescent tablet)**

Anhydrous citric acid 1000 mg

Aspirin 325 mg **(NSAID)\***

Sodium bicarbonate (heat-treated) 1916 mg

\*Nonsteroidal anti-inflammatory drug

**Purpose**

Anhydrous citric acid 1000 mg ..... Antacid

Aspirin 325 mg **(NSAID)\***.....Pain reliever

Sodium bicarbonate (heat-treated) 1916 mg .....Antacid

\*Nonsteroidal anti-inflammatory drug

**Uses**

**Uses** for the temporary relief of

- heartburn, acid indigestion and sour stomach when accompanied by headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- headache, body aches, and pain alone

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid

drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

### **Do not use**

**Do not use** ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients.

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic ■ you have asthma ■ you have a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.
- taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

### **When using this product**

**When using this product** do not take more than 8 tablets (adults) or 4 tablets (60 years and over) in a 24-hour period or use the maximum dosage for more than 2 weeks, expect under the advise and supervision of a physician.

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding
- feel faint ■ vomit blood ■ have bloody or black stools
- have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- pain lasts for more than 10 days or gets worse
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs

### **If pregnant or breast-feeding**

**If pregnant or breast-feeding**, ask a health professional before use. **It is especially**

**important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.**

**Keep out of reach of children.**

### **In case of overdose**

In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

■ do not exceed recommended dosage

■ to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking  
adults and children 12 years and over (up to 60 years of age): 2 tablets every 4 hours - do not exceed 8 tablets in 24 hours

60 years and over: 2 tablets every 4 hours - do not exceed 4 tablets in 24 hours

Children under 12 years: consult a doctor

Take this product any time – morning, noon or night – when you need relief from heartburn, acid indigestion, sour stomach with headache or body aches and pains.

Place two (2) tablets in 4 oz. of water. Dissolve tablets completely.

You do not have to drink any of the residue that may be at the bottom of the glass. The medicines are in the water.

### **Other information**

■ **each tablet contains:** sodium 524 mg

■ store at room temperature (59°-86°F)

### **Inactive ingredients**

■ none

### **Questions or comments?**

**1-888-228-6937**

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**



# MARKET BASKET EFFERVESCENT ANTACID AND PAIN RELIEF

aspirin, citric acid, sodium bicarbonate tablet, effervescent

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:53942-421
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO) (BICARBONATE ION - UNII:HN1Z RA3Q20)	SODIUM BICARBONATE	1916 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL) (ANHYDROUS CITRIC ACID - UNII:XF417D3PSL)	ANHYDROUS CITRIC ACID	1000 mg

**Product Characteristics**

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	25mm
<b>Flavor</b>		<b>Imprint Code</b>	L421
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53942-421-36	18 in 1 CARTON	10/22/2021	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	10/22/2021	12/31/2026

**Labeler** - Demoulas Super Markets, Inc. (007869647)

Revised: 12/2024

Demoulas Super Markets, Inc.