

**VIRO FLU PM RELIEF- acetaminophen, dextromethorphan hydrobromide,  
doxylamine succinate capsule, liquid filled  
OPMX LLC**

-----  
**Viro-Flu PM Relief**

**Drug Facts**

<b>Active ingredients (in each Softgels)</b>	<b>Purpose</b>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 15 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine

**Uses**

Temporarily relieves common cold/flu symptoms:

- Cough due to minor throat & bronchial irritation
- Sore throat
- Headache
- Minor aches & pains
- Fever
- Runny nose & sneezing

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy Alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening • blisters • rash • If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning**

If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). • If you are not sure whether a drug contains acetaminophen, ask a doctor or

pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

**When using this product**

- Excitability may occur, especially in children
- Marked drowsiness may occur
- Avoid Alcoholic drinks
- Be careful when driving a motor vehicle or operating machinery
- Alcohol, sedatives, & tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- Pain or cough gets worse or lasts more than 7 days
- Fever gets worse or lasts more than 3 days
- Redness or swelling is present
- New symptoms occur
- Cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

**Directions**

- Take only as directed
- Do not exceed 4 doses per 24 hrs

Adults & children 12 yrs & over	2 softgels with water every 6 hrs
Children 4 to under 12 yrs	Ask a doctor
Children under 4 yrs	Do not use

**Other information**

- Store at room temperature

**Inactive ingredients**

D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol special, titanium dioxide

**Questions?**

Call toll free 619-600-5632 Monday through Friday 9AM - 5PM EST

**Made for:**

**RRX**

**Exclusively distributed by:**

**OPMX**

**Chula Vista, CA 91910**

**PRINCIPAL DISPLAY PANEL**

NDC 69729-057-10

Viro-Flu PM

- Cough/ Tos
- Cold/Resfriado
- Nasal Congestion & Runny Nose/ Flujo y congestión nasal
- Sore Throat/ Ardor de Garganta
- Headache and Body Ache/ Dolor de Cabeza y Cuerpo
- Cough/Tos
- Flu/ Gripe

10 Softgels

**Drug Facts**

**Active ingredients (in each Softgel)**  
 Acetaminophen 325 mg, Pain reliever/fever reducer  
 Dextromethorphan Hbr 15mg, Cough suppressant  
 Doxylamine succinate 6.25 mg, Antihistamine

**Uses**  
 Temporarily relieves common cold/flu symptoms: • Cough due to minor throat and bronchial irritation • Sore throat • Headache • Fever • Runny nose and sneezing

**Warnings**  
 Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take: • More than 4 doses in 24 hr, which is the maximum daily amount for this product. • With other drugs containing acetaminophen. • 3 or more alcoholic drinks daily while using this product.  
**Every alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • Skin redness • Blisters • Rash  
 • If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.  
**Do not use:** • With any other drug containing acetaminophen (prescription or nonprescription). • If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
**Ask a doctor before use if you have:** • Liver disease • Glaucoma • Cough that occurs with too much phlegm (mucus). • A breathing problem or chronic cough that lasts or as occurs with smoking. • Asthma, chronic bronchitis or emphysema.

**Drug Facts (continued)**

**Purpose**  
 • Trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are:** • Taking sedatives or tranquilizers • Taking the blood thinning drug warfarin.

**When using this product:** • Crisitalily may occur, especially in children. • Marked drowsiness may occur. • Avoid alcoholic drinks.

**Stop use and ask a doctor if:** • Pain or cough gets worse or lasts more than 7 days. • Fever gets worse or lasts more than 3 days. • Redness or swelling is present. • Cough comes back or occurs with rash or headache that lasts. • New symptoms occur. These could be signs of a serious condition.

**Keep out of reach of children.**

**If pregnant or breast-feeding,** ask a health professional before use. Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center for advice. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
 • Take only as directed. • Do not exceed 4 doses per 24 hours. Adults & children 12 yrs & over: 2 softgels with water every 6 hrs. Children 4 to under 12 yrs: Ask a doctor. Children under 4 yrs: Do not use.

**Other information** • Store at room temperature.

**Inactive ingredients:** D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, titanium dioxide.

**Questions?**  
 Call toll free 619-600-5632 Monday through Friday 9AM - 5PM EST

READ & KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

TAMPER EVIDENT: Do not open or if blister unit is torn, broken or shows any sign of tampering

Exclusively distributed by:

OPMX  
 Chula Vista, CA 91910  
 Phone: 619-400-5632

Made for:  
 PRX  
 San Diego, CA

Nighttime Formula  
**Viro-Flu**  
 p.m. GelCaps

NDC 69729-057-10

**Viro-Flu**  
 p.m. GelCaps  
 10 Soft Gels

**New Formula**

**Fever/Fiebre  
 Cold / Resfriado  
 Nasal Congestion & Runny Nose /  
 Flujo y congestión Nasal  
 Sore Throat / Ardor de Garganta  
 Headache and Body Ache /  
 Dolor de Cabeza y Cuerpo  
 Cough / Tos  
 Flu / Gripe**

Nighttime Formula  
**Viro-Flu**  
 p.m. GelCaps

Lot No.:  
 Exp. date:

QR Code

8 10053 18012 9  
 REV. 06.2022 ITEM # RCV1085

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69729-057
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	AP02
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:69729-057-10	10 in 1 CARTON	06/10/2022	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
---------------------------	---	-----------------------------	---------------------------

OTC Monograph Drug	M012	06/10/2022	
--------------------	------	------------	--

**Labeler** - OPMX LLC (029918743)

**Establishment**

Name	Address	ID/FEI	Business Operations
SOFTECH PHARMA PRIVATE LIMITED		677111277	label(69729-057) , manufacture(69729-057) , pack(69729-057)

Revised: 4/2024

OPMX LLC