

LICEL - piperonyl butoxide, pyrethrum extract oil

Sujanil Chemo Industries

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Licel Lice Treatment Oil

Drug Facts

Active ingredients

Piperonyl butoxide 4%

Pyrethrum extract (equivalent to 0.33% pyrethrins)

Purpose

Lice Treatment

Uses

treats head, pubic (crab), and body lice

Warnings

For external use only

Do not use

- near eyes
- inside nose, mouth, or vagina
- on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.

Ask a doctor before use if you are:

allergic to ragweed. May cause breathing difficulty or an asthmatic attack.

When using this product

- keep eyes tightly closed and protect eyes with a washcloth or towel.
- if product gets into eyes, flush with water right away.
- Scalp itching or redness may occur

Stop use and ask a doctor if

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Shake well before using

**Important: Read warnings before use
adults and children 2 years and over:**

Inspect

- check each household member with a magnifying glass in bright light for lice/nits (eggs)
- look for tiny nits near scalp, beginning at back of neck and behind ears
- examine small sections of hair at a time
- unlike dandruff which moves when touched, nits stick to the hair
- if either lice or nits are found, treat with this product

Treat

- apply thoroughly to dry hair or other affected area. For head lice, first apply behind ears and to back of neck.
- allow product to remain for a minimum 10 minutes.
- wash area thoroughly with warm water and soap or shampoo.
- for head lice, towel dry hair and comb out tangles.

Remove lice and their eggs (nits)

- use a fine-tooth or special lice/nit comb. Remove any remaining nits by hand (using a throw-away glove).
- hair should remain slightly damp while removing nits
- if hair dries during combing, dampen slightly with water
- for head lice, part hair into sections. Do one section at a time.
- starting on top of the head. Longer hair may take 1 to 2 hours.
- lift a 1- to 2- inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp.
- pin back each strand of hair after combing.
- clean comb often. Wipe nits away with tissue and discard in a plastic bag. Seal bag and discard to prevent lice from coming back.
- after combing, thoroughly recheck for lice/nits. Repeat combing if necessary.
- check daily for any lice/nits that you missed.
- a second treatment must be done in 7 to 10 days to kill any newly hatched lice.
- if infestation continues, see a doctor for other treatments.
- **children under 2 years: ask a doctor**

Other Information

- store in cool and dry place
- keep carton for important product information
- see package insert for additional information.

Inactive ingredients

vegetables oil

Questions or comments?

855-KIL-LICE - 855-545-5423

www.sujanil.com or e-mail: sales@sujanil.com

Package Insert

DAILY CHECKING

To Prevent Reinfestation:

- Carefully inspect infested family member daily for at least 2 weeks. If you see eggs or nits, repeat step 2.
- Check other family members for Head Lice. If anyone in the family is infested, follow Step 1 and Step 2. Repeat Step 3 with any articles or items with which they have come in contact.

Second Application

- Repeat the same procedures within 7 - 10 days for

Step 1: Treat and Step 2: Remove

There may be some eggs that were not combed out after the first treatment. If infestation continues, see a doctor for other treatments. These eggs will hatch and need to be killed with the second treatment.

Did you know?

Head Lice

- Head lice lay small white eggs (nits) on hair shaft close to the scalp
- Nits are most easily found on back of neck or behind ears
- Disinfect hats, ribbons, scarves, coats, towels and bed linens by machine washing in hot water (above 54°C (130°F), then using hottest dryer cycle for at least 20 minutes
- Items that cannot be washed (bedspreads, blankets, pillows, stuffed toys, etc.) should be dry-cleaned or sealed in a plastic bag for 4 weeks, then removed outdoors and shaken out very hard before using again
- Items that cannot be washed, dry-cleaned, or stored may be sprayed with a product designed for this purpose
- Soak all combs and brushes in hot water (above 54°C (130°F) for at least 10 minutes
- Vacuum all carpets, mattresses, upholstered furniture, and car seats that may have been used by affected people.

Pubic (Crab) Lice

- Pubic Lice may be transmitted by sexual contact. Sexual partners should be treated simultaneously to avoid reinfestation.
- Pubic Lice are very small and look like brown or gray dots on skin.
- Pubic Lice usually cause intense itching and lay small white eggs (nits) on the hair shaft generally close to the skin surface.
- Pubic Lice may be presented on the short hairs of groin, thighs, trunk and underarms and occasionally on the beard and mustache.
- Pubic Lice infested underwear can be disinfected by machine washing in hot water (above 54°C (130°F), then using hottest dryer cycle for at least 20 minutes.

Body Lice

- Body Lice and their eggs (nits) are generally found in the seams of clothing particularly in waistline and armpit area.
- Body Lice feed on skin then return to clothing to lay their eggs.
- Body Lice infested clothing can be disinfected by machine washing in hot water (above 54°C (130°F), then using the hottest dryer cycle for at least 20 minutes.

- Body Lice nits can remain dormant for up to 30 days and therefore clothing should not be sealed in a plastic bag
- Save this package insert for future reference.

STEP 1 TREAT

Licel Lice Treatment Oil

Steps to Success
with 1-2-3 Lice Control System

Complete Lice Control for your family and home.

SHAKE WELL BEFORE USING

DIRECTIONS: IMPORTANT: READ WARNINGS BEFORE USING. A SECOND TREATMENT MUST BE DONE IN 7-10 DAYS TO KILL ANY NEWLY HATCHED LICE.

Warnings

For external use only

Do not use:

- near eyes,
- inside nose, mouth or vagina,
- on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.

Ask a doctor before use if you are

- allergic to ragweed. May cause breathing difficulty or an asthmatic attack

When using this product

- keep eyes tightly closed and protect eyes with a washcloth or towel
- if product gets in eyes, flush with water right away
- scalp itching or redness may occur

Stop use and ask a doctor if

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs
- Keep out of reach of children. if swallowed, get medical help or contact a Poison Control Center right away

See Usage Guidelines below to estimate how much product you will need.

Inspect

- check each household member with a magnifying glass in bright light for lice/nits (eggs)
- look for tiny nits near scalp, beginning at back of neck and behind ears
- examine small sections of hair at a time

- unlike dandruff which moves when touched, nits stick to the hair
- If either lice or nits are found, treat with this product

Note:

- Use towels to protect child's eyes from treatment. Ask the child to keep eyes tightly closed.
- Apply Licel Lice Treatment oil thoroughly to **DRY HAIR** or other affected area
- Wetting the hair dilutes the treatment making it less effective.
- Also, lice can hold their breath when immersed in water making it difficult for the active ingredient to penetrate the lice.
- For head lice, thoroughly massage oil into scalp, behind ears and to the back of the neck. Lice can crawl up and down the shaft of the hair very quickly so it is important to apply the treatment from the roots to the ends of the hair.
- Allow product to remain for 10 minutes, but no longer (or other affected area) for at least an hour.
- Wash area thoroughly with warm water and soap or shampoo
- For head lice, towel dry hair and comb out tangles.
- Proceed to Step 2.
- A second treatment must be done in 7 to 10 days to kill any newly hatched lice.

Usage Guidelines

<u>Hair Length</u>	<u>Approximate Amount for 1 Adult/Child:</u>
Short (ear length or shorter)	First Application : 1 oz – 2 oz
	<u>Second Application : 1 oz – 2 oz</u>
	Total : 2 oz – 4 oz
Medium (shoulder length)	First Application : 2 oz – 3 oz
	<u>Second Application : 2 oz – 3 oz</u>
	Total : 2 oz – 4 oz
Long (past shoulder length)	First Application : 3 oz – 4 oz
	<u>Second Application : 3 oz – 4 oz</u>
	Total : 6 oz – 8 oz

The Licel comb is proven 100% effective as demonstrated in laboratory studies performed by trained testers, individual results may vary.

STEP 2 REMOVE

NOTE:

- A fine - toothed comb or special lice / nit removing comb (included) must be used to remove lice, eggs, and nits from hair.
- Before starting have on hand:
 - extra towels
 - bobby pins or hair clips
 - facial tissues or paper towels

- regular comb for parting and detangling hair
- throw-away gloves
- Find a distraction for your child (e.g. books, TV, game)
- Comb out hair with a regular wide-tooth comb to take out snarls and tangles.
- When combing, use the fine-toothed lice/nit comb provided and make sure the teeth are facing you.
- Part hair into section. Do one section at a time. Starting on top of the head. Longer hair may take 1 to 2 hours.
- Lift a 1- to 2- inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp.
- Always comb away from the head.
- Pin back each strand of hair after combing.
- Wipe eggs and nits from the teeth of the comb with a tissue or paper towel often.
- Remove any remaining nits by hand (using a throw-away glove)
- Discard all the used tissue or cloths into bag, seal the bag and dispose of immediately to prevent lice from coming back.
- After combing, thoroughly recheck for lice/nits. Repeat combing if necessary
- After you have combed and pinned the entire head, remove all bobby pins or hair clips and thoroughly recheck for lice/nits. Repeat combing if necessary.
- Rinse hair thoroughly with warm water, If desired, you may also wash with regular shampoo.
- Check daily for any lice/nits that you missed.
- A second treatment must be done in 7 to 10 days to kill any newly hatched lice.
- If infestation continues, see a doctor for other treatments.
- Children under 2 years: Ask a doctor

Step 3 Control

- Head lice lay small white eggs (nits) on the hair shaft close to the scalp
- nits are most easily found on the back of the neck or behind the ears
- Disinfect hats, hair ribbons, scarves, coats, towels and bed linens by machine washing in hot water (above 54°C (130°F)), then using the hottest dryer cycle for at least 20 minutes.
- Items that cannot be washed (bedspreads, blankets, pillows, stuffed toys, etc.) should be dry-cleaned or sealed in a plastic bag for 4 weeks, then removed outdoors and shaken out very hard before using again
- Items that cannot be washed, dry-cleaned, or stored may be sprayed with a product designed for this purpose
- Soak all combs and brushes in water (above 54°C (130°F)) for at least 10 minutes
- Vacuum all carpets, mattresses, upholstered furniture, and car seats that may have been used by affected people.

CHECKLIST

- Home and auto upholstery
- Pillows, pillowcases, mattresses
- Bed Linen
- Towels
- Stuffed animals toys
- Hats
- Headphones

- Headbands
- Bike/sports helmets
- Coats, gloves, scarves, earmuffs
- Play mats
- Clothing
- Barettes/Hair ornaments
- Hair rollers
- Combs and brushes
- Baseball caps

Principal Display Panel

NDC No.: 76441-902-97

Comb with Magnifier Included

India's leading Lice Killer

Sujanil

Maximum Strength

Licel

Lice Treatment Oil

Proven and tested in India

for more than 4 Decades now in USA

Manufactured By:

Sujanil Chemo Industries

P.O. Box No. 1451, Marketyard P.O.,

Pune - 411037 (M.S.) India

Carton Label

Drug Facts

Active ingredients	Purpose
Piperonyl butoxide 4%.....	Lice treatment
Pyrethrum extract	Lice treatment (equivalent to 0.33% pyrethrins)

Uses treats head, pubic (crab), and body lice

Warnings

For external use only

- Do not use:**
- near eyes,
 - inside nose, mouth or vagina,
 - on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.

Ask a doctor before use if you are

- allergic to ragweed. May cause breathing difficulty or an asthmatic attack

When using this product

- keep eyes tightly closed and protect eyes with a washcloth or towel
- If product gets in eyes, flush with water right away
- scalp itching or redness may occur

Stop use and ask a doctor if

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away

Directions

Shake well before using
Important: Read warnings before use
adults and children 2 years and over:

- Inspect**
- check each household member with a magnifying glass in bright light for lice/nits (eggs)
 - look for tiny nits near scalp, beginning at back of neck and behind ears
 - examine small sections of hair at a time
 - unlike dandruff which moves when touched, nits stick to the hair
 - If either lice or nits are found, treat with this product
- Treat**
- apply thoroughly to dry hair or other affected area. For head lice, first apply behind ears and to back of neck.
 - allow product to remain for minimum 10 minutes
 - wash area thoroughly with warm water and soap or shampoo
 - for head lice, towel dry hair and comb out tangles
- Remove lice and their eggs (nits)**
- use a fine-tooth or special lice nit comb. Remove any remaining nits by hand (using a throw-away glove)
 - hair should remain slightly damp while removing nits
 - if hair dries during combing, dampen slightly with water

Drug Facts (continued)

- for head lice, part hair into sections. Do one section at a time starting on top of head. Longer hair may take 1 to 2 hours.
- lift a 1-to 2-inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp
- pin back each strand of hair after combing
- clean comb often. Wipe nits away with tissue and discard in a plastic bag. Seal bag and discard to prevent lice from coming back
- after combing, thoroughly recheck for lice/nits. Repeat combing if necessary
- check daily for any lice/nits that you missed
- a second treatment must be done in 7 to 10 days to kill any newly hatched lice
- if infestation continues, see a doctor for other treatments
- children under 2 years: ask a doctor

Other information

- store in cool and dry place
- keep carton for important product information
- see package insert for additional information

Inactive ingredients

vegetables oil

Questions or comments?

855-KIL-LICE - 855-545-5423
www.sujanil.com or e-mail: sales@sujanil.com

'Licel' Lice Treatment Oil will kill lice completely without leaving a chemical residue. Amount of lice treatment oil needed will vary by hair length. See below for guidelines:

Hair Length	Approximate Amount for 1 Adult/Child:
Short (ear-length or shorter)	First Application : 1 oz - 2 oz Second Application : 1 oz - 2 oz Total : 2 oz - 4 oz
Medium (shoulder-length)	First Application : 2 oz - 3 oz Second Application : 2 oz - 3 oz Total : 4 oz - 6 oz
Long (past shoulder-length)	First Application : 3 oz - 4 oz Second Application : 3 oz - 4 oz Total : 6 oz - 8 oz

The Licel comb is proven 100% effective as demonstrated in laboratory studies performed by trained testers; individual results may vary.

LICE TREATMENT OIL



NDC No. : 76441-902-97



LICE TREATMENT OIL



A PRODUCT OF INDIA

Proven & tested in India
for more than 4 Decades
Now in USA

NET WT : 4 FL OZ (118 mL)



'Licel' Lice Treatment Oil

Lice Treatment

- Nothing proven more effective
- Leaves no chemical residue behind
- Easy to use oil

Licel 1-2-3 LICE ELIMINATION SYSTEM

Step 1 **Head Lice**

- Head lice lay small white eggs (nits) on hair shaft close to the scalp • Nits are most easily found on back of neck or behind ears • Disinfect hats, hair ribbons, scarves, coats, towels and bed linens by machine washing in hot water (above 54°C (130°F)), then using hottest dryer cycle for at least 20 minutes
- Items that cannot be washed (bedspreads, blankets, pillows, stuffed toys, etc.) should be dry-cleaned or sealed in a plastic bag for 4 weeks, then removed outdoors and shaken out very hard before using again • Items that cannot be washed, dry-cleaned, or stored may be sprayed with a product designed for this purpose • Soak all combs and brushes in hot water (above 54°C (130°F)) for at least 10 minutes • Vacuum all carpets, mattresses, upholstered furniture, and car-seats that may have been used by affected people.

Step 2 **Pubic (Crab) Lice**

- Pubic lice may be transmitted by sexual contact
- Sexual partners should be treated simultaneously to avoid reinfection • Lice are very small and look like brown or grey dots on skin • Pubic lice usually cause intense itching and lay small white eggs (nits) on the hair shaft generally close to the skin surface
- Pubic lice may be present on the short hairs of the groin, thighs, trunk and underarms, and occasionally on the beard and mustache • Disinfect underwear by machine washing in hot water (above 54°C (130°F)), then using hottest dryer cycle for at least 20 minutes.

Step 3 **Body Lice**

- Body lice and their eggs (nits) are generally found in the seams of clothing, particularly in the waistline and armpit area • Body lice feed on skin when return to clothing to lay their eggs • Disinfect clothing by machine washing in hot water (above 54°C (130°F)), then using hottest dryer cycle for at least 20 minutes
- Do not seal clothing in a plastic bag because nits can remain dormant for up to 30 days

Manufactured By :

sujanil chem industries
P.O. Box No. 1451, Marketyard P.O.,
Pune - 411037 (M.S.) India

Imported and Distributed by :

Global Tradelinks
5938, Benjamin Road, TAMPA, FL 33634, USA
Tel. : 727-388-1173, Fax is 813-886-3881
E-mail : gaj@sujanil.com

Batch No. :

Mfg. Date :

Expiry Date :



Bottle Label

Active ingredients **Purpose**
 Piperonyl butoxide 4% Lice treatment
 Pyrethrum extract
 (equivalent to 0.33% pyrethrins) ... Lice treatment

Uses treats head, pubic (crab), and body lice

Inactive ingredients :Vegetables Oil

Warnings : For external use only

Do not use :

- near eyes,
- inside nose, mouth or vagina,
- on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.

Ask a doctor before use if you are

- allergic to ragweed. May cause breathing difficulty or an asthmatic attack

When using this product

- keep eyes tightly closed and protect eyes with a wash cloth or towel
- If product gets in eyes, flush with water right away
- scalp itching or redness may occur

Stop use and ask a doctor if

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away

Other information : ■ store in cool and dry place ■ keep carton for important product information ■ see package insert for additional information


Questions or comments?
 855-KIL-LICE - 855-545-5423

www.sujanil.com or e-mail : sales@sujanil.com

NDC No. : 76441-902-97



LICE TREATMENT OIL



A PRODUCT OF INDIA

Proven & tested in India for more than 4 decades

Now in USA

NET WT : 4 FL OZ (118 mL)



Directions:

Shake well before using

Important: Read warnings before use adults and children 2 years and over:

Inspect : ■ check each household member with a magnifying glass in bright light for lice/nits (eggs) ■ look for tiny nits near scalp, beginning at back of neck and behind ears ■ examine small sections of hair at a time ■ unlike dandruff which moves when touched, nits stick to the hair ■ if either lice or nits are found, treat with this product.

Treat : ■ apply thoroughly to dry hair or rather affected area. For head lice, first apply behind ears and to back of neck ■ allow product to remain for minimum 10 minutes ■ wash area thoroughly with warm water and soap or shampoo ■ for head lice, towel dry hair and comb out tangles.

Remove lice and their eggs (nits) : ■ use a fine tooth or special lice/nit comb. Remove any remaining nits by hand (using a throw away glove) ■ hair should remain slightly damp while removing nits ■ if hair dries during combing, dampen slightly with water ■ for head lice, part hair into sections. Do one section at a time starting on top of head. Longer hair may take 1 to 2 hours ■ lift a 1 to 2 inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp ■ pin back each strand of hair after combing ■ clean comb often. Wipe nits away with tissue and discard in a plastic bag. Seal bag and discard to prevent lice from coming back ■ after combing, thoroughly recheck for lice/nits. Repeat combing, if necessary ■ check daily for any lice/nits that you missed ■ a second treatment must be done in 7 to 10 days to kill any newly hatched lice ■ if infestation continues, see a doctor for other treatments ■ **children under 2 years: ask a doctor**

Batch No. :

Mfg. Date :

Expiry Date :

Manufactured By :

sujanil chem industries

P.O. Box No. 1451, Marketyard P.O.,
 Pune - 411037 (M.S.) India

Imported and Distributed by :

Global Tradelinks

5936, Benjamin Road, TAMPA, FL 33634, USA

Tel. : 727-359-1173, Fax is 813-886-3881

E-mail : jay@sujanil.com

DAILY CHECKING



TO PREVENT REINFESTATION:

- Carefully inspect infested family member **daily** for at least 2 weeks. If you see eggs or nits, repeat step 2.
- Check other family members for Head Lice. If anyone in the family is infested, follow Step 1 and Step 2. Repeat Step 3 with any articles of items with which they have come into contact.

Second Application

- Repeat the same procedures within 7 - 10 days for

Step 1: Treat and Step 2: Remove

There may be some eggs that were not combed out after the first treatment. If infestation continues, see a doctor for other treatments. These eggs will hatch and need to be killed with the second treatment.

Did you know?

Head Lice

- Head lice lay small white eggs (nits) on hair shaft close to the scalp.
- Nits are most easily found on back of neck or behind ears.
- Disinfect hats, hair ribbons, scarves, coats, towels and bed linens by machine washing in hot water (above 54°C (130°F)), then using hottest dryer cycle for at least 20 minutes.
- Items that cannot be washed (bedspreads, blankets, pillows, stuffed toys, etc.) should be dry-cleaned or sealed in a plastic bag for 4

weeks, then removed outdoors and shaken out very hard before using again.

- Items that cannot be washed, dry-cleaned, or stored may be sprayed with a product designed for this purpose.
- Soak all combs and brushes in hot water (above 54°C (130°F)) for at least 10 minutes.
- Vacuum all carpets, mattresses, upholstered furniture, and car seats that may have been used by affected people.

Pubic (Crab) Lice

- Pubic Lice may be transmitted by sexual contact. Sexual partners should be treated simultaneously to avoid reinfestation.
- Pubic Lice are very small and look like brown or grey dots on skin.
- Pubic Lice usually cause intense itching and lay small white eggs (nits) on the hair shaft generally close to the skin surface.
- Pubic Lice may be present on the short hairs of groin, thighs, trunk and underarms and occasionally on the beard and mustache.
- Pubic Lice infested underwear can be disinfected by machine washing in hot water (above 54°C (130°F)), then using hottest dryer cycle for at least 20 minutes.

Body Lice

- Body Lice and their eggs (nits) are generally found in the seams of clothing particularly in waistline and armpit area.
- Body Lice feed on skin then return to clothing to lay their eggs.
- Body Lice infested clothing can be disinfected by machine washing in hot water (above 54°C (130°F)), then using hottest dryer cycle for at least 20 minutes.
- Body Lice nits can remain dormant for up to 30 days and therefore clothing should not be sealed in a plastic bag.
- Save this package insert for future reference.



LICE TREATMENT OIL

Questions or comments?
855-KIL-LICE • 855-545-5423

www.sujanil.com or e-mail : sales@sujanil.com

Manufactured By

sujanil chem industries

P.O. Box No. 1451, Marketyard (P.O.)
Pune 411037 (M.S.) India

Imported and Distributed by :

Global Tradelinks

5938, Benjamin Road, TAMPA, FL 33634, USA
Tel : 727-359-1173, Fax Is 813-886-3881
E-mail : jay@sujanil.com



NDC No. : 76441-902-97

Sujanil's
Maximum Strength
Licel
LICE TREATMENT OIL



A PRODUCT OF INDIA

Proven & tested in India
for more than 4 Decades
Now in USA



STEP 1 TREAT



LICE TREATMENT OIL

Steps to Success

with 1-2-3 Lice Control System

Complete Lice Control for your family and home.

SHAKE WELL BEFORE USING

DIRECTIONS - IMPORTANT - READ WARNINGS BEFORE USING. A SECOND TREATMENT MUST BE DONE IN 7-10 DAYS TO KILL ANY NEWLY HATCHED LICE.

Warnings

For external use only

Do not use:

- near eyes,
- inside nose, mouth or vagina,
- on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.

Ask a doctor before use if you are

- allergic to pyrethrin. May cause breathing difficulty or an asthmatic attack.

When using this product

- keep eyes tightly closed and protect eyes with a washcloth or towel
- if product gets in eyes, flush with water right away

scalp itching or redness may occur

Stop use and ask a doctor if

- breathing difficulty occurs
- eye irritation occurs



- skin or scalp irritation continues or infection occurs
- Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

See Usage Guidelines below to estimate how much product you will need.

Inspect

- check each household member with a magnifying glass in bright light for lice/nits (eggs)
- look for tiny nits near scalp, beginning at back of neck and behind ears
- examine small sections of hair at a time
- unlike dandruff which moves when touched, nits stick to the hair
- if either lice or nits are found, treat with this product

NOTE:

- Use towels to protect child's eyes from treatment. Ask the child to keep eyes tightly closed.
- Apply Licel Lice Treatment Oil thoroughly to **DRY HAIR** or other affected area.
- Wetting the hair dilutes the treatment making it less effective.
- Also, lice can hold their breath when immersed in water making it difficult for the active ingredient to penetrate the lice.
- For head lice, thoroughly massage oil into scalp, behind the ears and to the back of the neck. Lice can crawl up and down the shaft of the hair very quickly so it is important to apply

the treatment from the roots to the ends of the hair.

- Allow product to remain for minimum 10 minutes. For best results leave overnight.
- Wash area thoroughly with warm water and soap or shampoo.

- For head lice, towel dry hair and comb out tangles.
- Proceed to Step 2.
- A second treatment must be done in 7 to 10 days to kill any newly hatched lice.

Usage Guidelines

Hair Length	Approximate Amount for 1 Adult/Child
Short (ear length or shorter)	First Application : 1 oz - 2 oz
	Second Application : 1 oz - 2 oz
	Total : 2 oz - 4 oz
Medium (shoulder length)	First Application : 2 oz - 3 oz
	Second Application : 2 oz - 3 oz
	Total : 4 oz - 6 oz
Long (past shoulder length)	First Application : 3 oz - 4 oz
	Second Application : 3 oz - 4 oz
	Total : 6 oz - 8 oz

The Licel comb is proven 100% effective as demonstrated in laboratory studies performed by trained testers. Individual results may vary.

STEP 2 REMOVE

NOTE:

- A fine - toothed comb or a special lice / nit removing comb (included) must be used to remove lice, eggs, and nits from hair.
- Before starting have on hand:
 - extra towels
 - bobby pins or hair clips
 - facial tissues or paper towels
 - regular comb for parting and detangling hair
 - throw-away gloves
- Find a distraction for your child (e.g. books, TV, game)
- Comb out hair with a regular wide-tooth comb to take out snarls and tangles.



- When combing, use the fine-toothed lice/nit comb provided and make sure the teeth are facing you.
- Part hair into section. Do one section at a time. Starting on top of the head. Longer hair may take 1 to 2 hours.
- Lift a 1 to 2 inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp.

- Always comb away from the head.
- Pin back each strand of hair after combing.
- Wipe eggs and nits from the teeth of the comb with a tissue or paper towel often.
- Remove any remaining nits by hand (using a throw-away glove).
- Discard all the used tissues or cloths into bag, seal the bag and dispose of immediately to prevent lice from coming back.
- After combing, thoroughly recheck for lice/nits. Repeat combing if necessary.
- After you have combed and pinned the entire head, remove all bobby pins or hair clips and thoroughly recheck for lice/nits. Repeat combing if necessary.

STEP 3 CONTROL



- Head lice by small white eggs (nits) on the hair shaft close to the scalp
- Nits are most easily found on the back of the neck or behind the ears
- Disinfect hats, hair ribbons, scarves, coats, towels and bed linens by machine washing in hot water (above 54°C (130°F)), then using the hottest dryer cycle for at least 20 minutes
- Items that cannot be washed (bedspreads, blankets, pillows, stuffed toys, etc.) should be dry-cleaned or sealed in a plastic bag for 4 weeks, then removed outdoors and shaken out very hard before using again
- Items that cannot be washed, dry-cleaned, or stored may be sprayed with a product designed for this purpose
- Soak all combs and brushes in water (above 54°C (130°F)) for at least 10 minutes
- Vacuum all carpets, mattresses, upholstered furniture, and car seats that may have been used by affected people.

CHECKLIST

- Home and auto upholstery
- Pillows, pillowcases, mattresses
- Bed linen
- Towels
- Stuffed animals toys
- Hats
- Headphones
- Headbands
- Bikes / sports helmets
- Coats, gloves, scarves, earmuffs
- Play mats
- Clothing
- Barettes / Hair ornaments
- Hair rollers
- Combs and brushes
- Baseball caps

LICEL

piperoonly butoxide, pyrethrum extract oil

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:76441-902

Route of Administration

TOPICAL

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
PIPERONYL BUTOXIDE (UNII: LWK91TU9AH) (PIPERONYL BUTOXIDE - UNII:LWK91TU9AH)		PIPERONYL BUTOXIDE	4 mL in 100 mL	
PYRETHRUM EXTRACT (UNII: ZUM06L90GV) (PYRETHRUM EXTRACT - UNII:ZUM06L90GV)		PYRETHRUM EXTRACT	0.33 mL in 100 mL	
Inactive Ingredients				
Ingredient Name			Strength	
SESAME OIL (UNII: QX10HYY4QV)				
COCONUT OIL (UNII: Q9L0O73W7L)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76441-902-97	1 in 1 BOX		
1		118 mL in 1 BOTTLE		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC monograph final	part358G		06/20/2012	

Labeler - Sujanil Chemo Industries (650280816)

Establishment

Name	Address	ID/FEI	Business Operations
Sujanil Chemo Industries		915896189	manufacture(76441-902)

Revised: 9/2012

Sujanil Chemo Industries