SENNA LAXATIVE- sennosides tablet Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each tablet)

Sennosides 8.6 mg

Purpose

Laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 6 12 hours

Warnings

Do not use

• laxative products for longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- stomach pain
- nausea
- vomiting
- noticed a sudden change in bowel habits that continues over a period of 2 weeks

Stop use and ask a doctor if

you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

• take preferably at bedtime or as directed by a doctor

a ge	starting dosage	maximum dosage
adults and children 12 years	2 tablets once a	4 tablets twice a day
of age or older	day	
children 6 to under 12 years	1 tablet once a day	2 tablets twice a day
children 2 to under 6 years	1/2 tablet once a	1 tablet twice a day
•	day	-
children under 2 years	ask a doctor	ask a doctor

Other information

- each tablet contains: calcium 25 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, hypromellose*, liquid paraffin*, magnesium stearate, maltodextrin*, microcrystalline cellulose, mineral oil*, polyethylene glycol*, polyvinyl alcohol*, silicon dioxide*, stearic acid*, sodium lauryl sulfate*, talc*.

*contains one or more of these ingredients

Questions or comments?

Call **1-888-309-9030**

Principal Display Panel

COMPARE TO THE ACTIVE INGREDIENT IN SENOKOT®**

Senna Laxative

SENNOSIDES 8.6 mg

- Natural Vegetable Laxative Ingredient
- Gentle overnight relief

TABLETS

**This product is not manufactured or distributed by Purdue Products L.P., owner of the registered trademark Senokot®.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Package Label



PLD-E22W

Exp. Date:

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION,

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

- %OOF --Satisfaction Guaranteed! 0509-90-9030

DISTRIBUTED BY DOLGENCORP, LLC 300 MISSION RIDGE GOODLETTSVILLE, TN 37072

**This product is not manufactured or distributed by Purdue Products L.P., distributor of Senokot®.

Guestions or comments? Call 1-888-309-9030

*contains one or more of these ingredients

dioxide*, sodium lauryl sulfate*, stearic acid*, talc*

Inactive ingredients croscarmellose sodium, dibasic calcium phosphate dihydrate, hypromellose", liquid paraffinr, microcrystalline cellulose, mineral oil", polyethylene glycol", polyvinyl alcohol", silicon

- store between 25°C (77°F); excursions permitted between 15-30°C (59-86°F)
 - each tablet contains: calcium 25 mg

Other information

Drug Facts (continued)

_		<u> </u>
sak a doctor	sask a doctor	children under 2 years
1 tablet twice a day	1/2 tablet once a day	children 2 to under 6 years
2 tablets twice a day	1 tablet once a day	children 6 to under 12 years
4 tablets twice a day	S tablets once a day	adults and children 12 years of age and over
agesob mumixsm	starting dosage	ege
Directions Iske preferably at bedfime or as directed by a doctor		
If pregnant or breast-feeding, sak a health professional before use. Keep out of reach of children, In case of overdose, get medical help or contact a Polson Control Center right away.		
Stop use and sak a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition.		
Ask a doctor before use if you have stomach pain mausea womiting moticed a sudden change in bowel habits that continues over a period of 2 weeks		
Warnings Do not use laxative products for longer than 1 week unless directed by a doctor		
USes ■ relieves occasional constitation (irregularity) ■ generally produces a bowel movement in 6 -1 2 hours		
Active ingredient (in each tablet) Purpose Sennosides 8.6 mg		
		Drug Facts

Sennosides 8.6 mg laxative tablets

sennosides tablet

Product Information

Product Type HUMAN OTC DRUG NDC:55910-486 **Item Code (Source)**

Route of Administration ORAL

Active Ingredient/Active Moiety Basis of Ingredient Name Strength Strength SENNOSIDES A AND B (UNII: 1B5FPI42EN) (SENNOSIDES A AND B -

UNII:1B5FPI42EN)

SENNOSIDES A AND B 8.6 mg

Inactive Ingredients	
Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
PARAFFIN (UNII: 1900E3H2ZE)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
MINERAL OIL (UNII: T5L8T28FGP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	

Product Characteristics			
Color	brown	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	0813;AV;TCL;080;S8
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-486- 01	1 in 1 BOX	12/09/2015	04/26/2024
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	12/09/2015	04/26/2024

Labeler - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)

Registrant - P & L Development, LLC (079765031)

Revised: 10/2021 Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)