

BLOAT GUARD- poloxalene liquid
Phibro Animal Health

Bloat Guard®
(poloxalene)
Liquid Type A Medicated Article

For use in the manufacture of liquid feed supplements

For control of legume (alfalfa, clover) and wheat pasture bloat in cattle

Labeling

Active Drug Ingredient

Poloxalene 99.5%

Inert Ingredients:

Ethoxyquin (a preservative), 0.12%;

butylated hydroxytoluene (a preservative), 0.38%

Follow Directions For Use

Directions for Use in Liquid Feeds

Bloat Guard Liquid Type A Medicated Article must be thoroughly blended and evenly distributed into a liquid feed supplement and offered to cattle in a covered liquid feed supplement feeder with lick wheels. The formula for the liquid feed supplement is as follows:

For control of legume (alfalfa, clover, and wheat pasture bloat in cattle...7.5 g/lb	% w/w	For control of legume (alfalfa, clover) bloat in cattle grazing pre-bloom legumes...10.0 g/lb	% w/w
Poloxalene Type A Medicated Article (99.5%)	1.65	Poloxalene Type A Medicated Article (99.5%)	2.2
Ammonium Polyphosphate	2.66	Ammonium Polyphosphate	2.66
Phosphoric Acid	3.37	Phosphoric Acid	3.37
Sulfuric Acid	1.00	Sulfuric Acid	1.00
Water	10.00	Water	10.00
Molasses, liquid – sufficient to make	100.00	Molasses, liquid – sufficient to make	100.00
Vitamins A&D and/or trace minerals may be added.		Vitamins A&D and/or trace minerals may be added.	

The concentration of Bloat Guard per lb of liquid feed supplement is as follows:

Indications for Use	Required Level of Bloat Guard
For control of legume (alfalfa, clover) and wheat pasture bloat in cattle	7.5 g/lb (1.65% w/w)
For control of legume (alfalfa, clover) bloat in cattle grazing pre-bloom legumes	10.0 g/lb (2.2% w/w)

Important:

To maintain proper distribution of Bloat Guard in liquid feed supplements, the viscosity of the medicated liquid supplement must be no less than 275 cps. During storage the medicated liquid may separate out of liquid supplements within 5 days if not agitated. It is essential that all of the stored mixture be thoroughly remixed (for example, by using a recirculating pump) before being transferred to feeders with free-turning lick wheels. Movement of the lick wheels is sufficient to prevent separation in the feeders.

Feeding and Management Directions for Bloat Guard

These medicated supplements are effective for the control of legume (alfalfa, clover) and wheat pasture bloat in cattle, when consumed as recommended.

- Use a covered liquid feeder with 1 free-turning lick wheel per 25 head at a height so that all cattle can conveniently lick the wheel.
- The location of the liquid feeder is extremely important for adequate consumption. Place feeder where cattle congregate (for example, near drinking water), thus providing ready access to the medicated liquid feed supplement. In large pastures, additional feeders should be placed so that cattle are never more than 500 feet from a feeder while grazing.
- For adequate protection, it is essential that each animal consumes the total recommended dosage of medication daily.
- If grazing on pre-bloom legumes, use a liquid feed supplement containing 10 g of Bloat Guard per lb (2.2% w/w). Each animal must consume the medicated liquid feed supplement at a rate of 0.15 lb per 100 lb of body weight per day for adequate protection. Example: 1.5 lb of 2.2% medicated supplement for a 1000-lb animal daily. If consumption exceeds 0.2 lb per 100 lb of body weight per day, cattle should be changed to a liquid feed supplement containing 7.5 g of Bloat Guard per lb (1.65% w/w).
- If grazing legumes or wheat pasture, use a liquid feed supplement containing 7.5 g of Bloat Guard per lb (1.65% w/w). Each animal must consume the medicated liquid feed supplement at a rate of 0.2 lb per 100 lb of body weight per day for adequate protection. Example: 2 lb of the 1.65% medicated supplement for a 1000-lb animal daily.
- Introduce the feeding of the medicated liquid feed supplement at least 2-5 days before pasturing on legume or wheat to accustom the cattle to the medicated liquid supplement and to lick wheel feeding. If the medicated liquid feed supplement feeding is interrupted, this 2-5 day introductory feeding should be repeated.

Cold Weather Note:

At temperatures below 50°F, this product will thicken and may appear cloudy, but is not damaged.

Keep Container Closed When Not in Use

Store At Or Below 25°C (77°F), Excursions Permitted Up To 40°C (104°F)

Not For Human Use

Restricted Drug (California) – Use Only As Directed

Net Weight 450 lb (204.12 kg)

Bloat Guard is a registered trademark of Phibro Animal Health Corporation for poloxalene

Phibro Animal Health, Inc., Teaneck, NJ 07666

Approved by FDA under NADA #038-281

8816000



Bloat Guard[®] (poloxalene)

Liquid Type A Medicated Article

For use in the manufacture of liquid medicated feeds

For control of legume (alfalfa, clover) and wheat pasture bloat in cattle

Active Drug Ingredient
Poloxalene 99.5%
Inert Ingredients:
Ethoxyquin (a preservative), 0.12%;
butylated hydroxytoluene (a preservative), 0.38%

FOLLOW DIRECTIONS FOR USE

Bloat Guard is a registered trademark of Phibro Animal Health Corporation for poloxalene
 Phibro Animal Health Corporation, Teaneck, NJ 07666

Approved by FDA under NADA # 038-281

Net Weight 450 lb (204.12 kg)



Directions for Use in Liquid Feeds

Bloat Guard Liquid Type A Medicated Article must be thoroughly blended and evenly distributed into a liquid feed supplement and offered to cattle in a covered liquid feed supplement feeder with lick wheels. The formula for the liquid feed supplement is as follows:

For control of legume (alfalfa, clover) and wheat pasture bloat in cattle . . . 7.5 g/lb	% w/w	For control of legume (alfalfa, clover) bloat in cattle grazing pre-bloom legumes . . . 10.0 g/lb	% w/w
Poloxalene Type A Medicated Article (99.5%)	1.65	Poloxalene Type A Medicated Article (99.5%)	2.2
Ammonium Polyphosphate	2.66	Ammonium Polyphosphate	2.66
Phosphoric Acid (75%)	3.37	Phosphoric Acid (75%)	3.37
Sulfuric Acid	1.00	Sulfuric Acid	1.00
Water	10.00	Water	10.00
Molasses, liquid – sufficient to make	100.00	Molasses, liquid – sufficient to make	100.00
Vitamins A & D and/or trace minerals may be added.		Vitamins A & D and/or trace minerals may be added.	

The concentration of Bloat Guard per lb of liquid feed supplement is as follows:

Indications for Use	Required Level of Bloat Guard
For control of legume (alfalfa, clover) and wheat pasture bloat in cattle	7.5 g/lb (1.65% w/w)
For control of legume (alfalfa, clover) bloat in cattle grazing pre-bloom legumes	10.0 g/lb (2.2% w/w)

Important: To maintain proper distribution of Bloat Guard in liquid feed supplements, the viscosity of the medicated liquid supplement must be no less than 275 cps. During storage the medicated liquid may separate out of liquid supplements within 5 days if not agitated. It is essential that all of the stored mixture be thoroughly remixed (for example, by using a recirculating pump) before being transferred to feeders with free-turning lick wheels. Movement of the lick wheels is sufficient to prevent separation in the feeders.

Feeding and Management Directions for Bloat Guard

These medicated feeds are effective for the control of legume (alfalfa, clover) and wheat pasture bloat in cattle, when consumed as recommended.

- Use a covered liquid feeder with 1 free-turning lick wheel per 25 head at a height so that all cattle can conveniently lick the wheel.
- The location of the liquid feeder is extremely important for adequate consumption. Place feeder where cattle congregate (for example, near drinking water), thus providing ready access to the medicated liquid feed. In large pastures, additional feeders should be placed so that cattle are never more than 500 feet from a feeder while grazing.
- For adequate protection, it is essential that each animal consumes the total recommended amount of medication daily.
- If grazing on pre-bloom legumes, use a liquid feed containing 10 g of Bloat Guard per lb (2.2% w/w). Each animal must consume the medicated liquid feed at a rate of 0.15 lb per 100 lb of body weight per day for adequate protection.
 Example: 1.5 lb of 2.2% medicated feed for a 1000-lb animal daily. If consumption exceeds 0.2 lb per 100 lb of body weight per day, cattle should be changed to a liquid feed containing 7.5 g of Bloat Guard per lb (1.65% w/w).
- If grazing legumes or wheat pasture, use a liquid feed containing 7.5 g of Bloat Guard per lb (1.65% w/w). Each animal must consume the medicated liquid feed at a rate of 0.2 lb per 100 lb of body weight per day for adequate protection.
 Example: 2 lb of the 1.65% medicated feed for a 1000-lb animal daily.
- Introduce the feeding of the medicated liquid feed at least 2-5 days before pasturing on legume or wheat to acustom the cattle to the medicated liquid and to lick wheel feeding. If the medicated liquid feed feeding is interrupted, this 2-5 day introductory feeding should be repeated.

Cold Weather Note: At temperatures below 50°F, this product will thicken and may appear cloudy, but is not damaged.

Keep Container Closed When Not in Use

Store At Or Below 25°C (77°F), Excursions Permitted Up To 40°C (104°F)

Not For Human Use

Restricted Drug (California) – Use Only as Directed

8816000
101-8031-07



Lot

Exp

(01) 1 03 66104 88160 8

Distributed by:
Phibro Animal Health Corporation
 Teaneck, NJ 07666

BLOAT GUARD

poloxalene liquid

Product Information

Product Type	OTC TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:66104-8816
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POLOXALENE (UNII: V8B3K56SW0) (POLOXALENE - UNII:V8B3K56SW0)	POLOXALENE	240.4 g in 0.45 kg

Inactive Ingredients

Ingredient Name	Strength
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ETHOXYQUIN (UNII: 9T1410R4OR)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66104-8816-0	204.12 kg in 1 CONTAINER		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA038281	08/14/1963	

Labeler - Phibro Animal Health (006989008)**Registrant** - Phibro Animal Health (006989008)

Revised: 7/2019

Phibro Animal Health