

**MAXIMUM STRENGTH COLD FLU AND SORE THROAT- acetaminophen,
dextromethorphan hbr, guaifenesin, phenylephrine hcl solution
KROGER COMPANY**

Kroger Maximum Strength Cold, Flu & Sore Throat

Active ingredients (in each 20 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCl 10 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pain
 - sore throat
 - headache
 - stuffy nose
 - sinus congestion and pressure
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks everyday while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product,

do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided.
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter
- **adults and children 12 years of age and older:** 20 mL in dosing cup provided every 4 hours
- **children under 12 years of age:** Do not use

Other information

- **each 20 mL contains:** sodium 8 mg
- store at room temperature
- do not refrigerate.

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C Blue No. 1, FD&C Red No. 40, flavor, potassium citrate, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

Questions or comments?

1-866-467-2748

Principal Display Panel

Compare to Maximum Strength †Mucinex[®] Fast-Max[®] Cold, Flu & Sore Throat* Active Ingredients

NDC# 30142-731-06

Maximum Strength †

FAST

Mucus relief

Cold, Flu & Sore Throat

Acetaminophen -Pain Reliever/Fever Reducer

Dextromethorphan HBr- Cough Suppressant

Guaifenesin -Expectorant

Phenylephrine HCl- Nasal Decongestant

RELIVES

Headache, Fever & Sore Throat

Nasal & Chest Congestion

Controls Cough

Thins & Loosen Mucus

For Ages 12+

6 FL OZ (180 mL)

Tamper evident: Do not use if printed seal under cap is broken or missing.

***Per 4-hour dose.**


AGES 12+ YEARS

Distributed by:

***This product is not manufactured or distributed by Reckitt Benckiser, the distributor of Maximum Strength Mucinex® Fast-Max™ Cold, Flu & Sore Throat.**

PEEL CORNER TO READ COMPLETE DRUG FACTS AND INFORMATION

MULTI-SYMPTOM



MAXIMUM STRENGTH COLD FLU AND SORE THROAT

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:30142-731
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYL GALLATE (UNII: 8D45NN7V92)	
WATER (UNII: 059QF0K00R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	blue	Score	
Shape		Size	
Flavor	FRUIT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30142-731-06	180 mL in 1 BOTTLE, PLASTIC; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)	09/03/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/03/2020	

Labeler - KROGER COMPANY (006999528)

Revised: 7/2024

KROGER COMPANY