

**NIGHTTIME COLD AND FLU RELIEF- acetaminophen, doxylamine succinate, dextromethorphan hbr capsule, liquid filled**  
**Bi-Mart**

-----

**Night-Time Cold & Flu Relief**

***Active ingredients (in each softgels)***

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

***Purpose***

Pain reliever/fever reducer

Cough suppressant

Antihistamine

***Uses***

Temporarily relieves common cold and flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or non-prescription).
- if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drink
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**ask a health professional before use.

**Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-122) right away. Quick medical attention is critical for adults and for children even if you do not notice any signs or symptoms.

**Direction**

- take only as directed

- do not exceed 4 doses per 24 hours

---

<b>adults and children 12 years and over</b>	take 2 softgels with water every 6 hours
<b>children 4 to under 12 years</b>	ask a doctor
<b>children under 4 years</b>	do not use

---

***Other information***

- store at room temperature.
- **DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED**

***Inactive ingredients***

D&C yellow# 10, FD&C Blue #1, Gelatin, Glycerin, Polyethylene Glycol 400, Povidone K30, Propylene Glycol, Purified Water, Sorbitol Sorbitan Solution

***Questions?***

call toll-free **1-844-912-4012**

**Distributed by:**

**BI-MART**

Eugene, OR 97402

**PRINCIPAL DISPLAY PANEL-24 Count**

**NDC 37835-522-24**

Compare to the active ingredients in Vicks <sup>®</sup>NyQuil <sup>®</sup>Cold & Flu LiquiCaps <sup>®</sup>\*

**NIGHTTIME COLD AND FLU RELIEF**

Acetaminophen,

Doxylamine Succinate,

Dextromethorphan HBr

24 SoftGels

\*This product is not manufactured or distributed by the owner of the registered trademark Vicks <sup>®</sup>NyQuil <sup>®</sup>Cold & Flu LiquiCaps <sup>®</sup>\*

701167551707  
 0  
 7155791107  
 3

**BI-MART** NDC 37835-522-24

Compare to the active ingredients in Vicks<sup>®</sup> NyQuil<sup>®</sup> Cold & Flu LiquiCaps<sup>®\*</sup>

# NIGHTTIME

COLD & FLU RELIEF

  
**Acetaminophen**  
 Doxylamine Succinate  
 Dextromethorphan HBr

**24**  
 SOFTGELS

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

**Drug Facts**

**Active Ingredients (in each softgel)**

Acetaminophen 325 mg.....Pain reliever/fever reducer  
 Dextromethorphan HBr 15 mg.....Cough suppressant  
 Doxylamine succinate 6.25 mg.....Antihistamine

**Uses** Temporarily relieves common cold and flu symptoms: ■ cough due to minor throat and bronchial irritation ■ sore throat ■ headache ■ minor aches and pains ■ fever ■ runny nose and sneezing

**Warnings**

**Liver warnings:** This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4 doses in 24 hours, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash ■ if a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other products containing acetaminophen (prescription or non-prescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts or as occurs with smoking.

PEEL FOR DIRECTIONS 522-24-BM 02/25

LIFT HERE

**Drug Facts (continued)**

asthma, chronic bronchitis, or emphysema ■ trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are** ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

**When using this product** ■ do not exceed recommended dosage ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if** ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults and for children even if you do not notice any signs or symptoms.

**Directions**

■ take only as directed  
 ■ do not exceed 4 doses per 24 hours

adults and children 12 years and over	take 2 softgels with water every 6 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

**Other information**

■ store at room temperature

**Inactive ingredients**

D&C Yellow #10, FD&C Blue #1, gelatin, glycerin, polyethylene glycol 400, povidone K30, propylene glycol, purified water, sorbitol sorbitan solution.

**Questions?** call toll-free 1-844-912-4012

\*This product is not manufactured or distributed by the owner of the registered trademark Vicks<sup>®</sup> NyQuil<sup>®</sup> Cold & Flu LiquiCaps<sup>®</sup>

Distributed by: **BI-MART** Eugene, OR 97402

BI-MART UNCONDITIONAL MONEY BACK GUARANTEE  
 If you are not completely satisfied with your BI-MART QUALITY PRODUCT, regardless of the reason, return the unused portion and your entire purchase price will be cheerfully refunded.

**PRINCIPAL DISPLAY PANEL-96 Count**  
**NDC 37835-522-96**

Compare to the active ingredients in Vicks<sup>®</sup> NyQuil<sup>®</sup> Cold & Flu LiquiCaps<sup>®\*</sup>

**NIGHTTIME COLD AND FLU RELIEF**

Acetaminophen,  
 Doxylamine Succinate,  
 Dextromethorphan HBr  
 96 SoftGels

\*This product is not manufactured or distributed by the owner of the registered trademark Vicks<sup>®</sup> NyQuil<sup>®</sup> Cold & Flu LiquiCaps<sup>®\*</sup>



**BI-MART**

NDC 37835-522-96

Compare to the active ingredients in Vicks® NyQuil® Cold & Flu LiquiCaps®\*

# NIGHTTIME

**COLD & FLU RELIEF**

**Acetaminophen**  
Doxylamine Succinate  
Dextromethorphan HBr

Value Size!

**96**  
SOFTGELS

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

**Drug Facts**

**Active ingredients (in each softgel)**

Acetaminophen 325 mg ..... Pain reliever/fever reducer  
Dextromethorphan HBr 15 mg ..... Cough suppressant  
Doxylamine succinate 6.25 mg ..... Antihistamine

**Uses** Temporarily relieves common cold and flu symptoms: ■ cough due to minor throat and bronchial irritation ■ sore throat ■ headache ■ minor aches and pains ■ fever ■ runny nose and sneezing

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4 doses in 24 hours, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other products containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts or as occurs with smoking.

522-96-BM 02/25  
PEEL FOR DIRECTIONS

LIFT HERE

**Drug Facts (continued)**

asthma, chronic bronchitis, or emphysema ■ trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are** ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

**When using this product** ■ do not exceed recommended dosage ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if** ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults and for children even if you do not notice any signs or symptoms.

**Directions**

■ take only as directed  
■ do not exceed 4 doses per 24 hours

adults and children 12 years and over	take 2 softgels with water every 6 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

**Other information**

■ store at room temperature

**Inactive ingredients**

D&C Yellow #10, FD&C Blue #1, gelatin, glycerin, polyethylene glycol 400, povidone K30, propylene glycol, purified water, sorbitol sorbitan solution.

**Questions?** call toll-free 1-844-912-4012

\*This product is not manufactured or distributed by the owner of the registered trademark Vicks® NyQuil® Cold & Flu LiquiCaps®

Distributed by: **BI-MART** Eugene, OR 97402

**BI-MART UNCONDITIONAL MONEY BACK GUARANTEE**  
If you are not completely satisfied with your BI-MART QUALITY PRODUCT, regardless of the reason, return the unused portion and your entire purchase price will be cheerfully refunded.

## NIGHTTIME COLD AND FLU RELIEF

acetaminophen, doxylamine succinate, dextromethorphan hbr capsule, liquid filled

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:37835-522
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	

### Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	21mm
<b>Flavor</b>		<b>Imprint Code</b>	CF02
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37835-522-24	24 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2025	
2	NDC:37835-522-96	96 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2025	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/01/2025	

**Labeler** - Bi-Mart (027630078)

**Registrant** - Bi-Mart (027630078)

### Establishment

Name	Address	ID/FEI	Business Operations
OCEAN HEALTHCARE PVT. LTD.		873673519	manufacture(37835-522)