

SCOT-TUSSIN DIABETES COUGH FORMULA WITH DM- dextromethorphan hydrobromide liquid

SCOT-TUSSIN Pharmacal Co., Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Scot-Tussin Diabetes Cough Formula with DM

Active Ingredient

(in each 5 mL. tsp. teaspoon)

Dextromethorphan HBr. USP 10mg

Purpose

Cough Suppressant

Uses

- Temporarily quiets and calms a dry cough due to minor throat and bronchial irritation.

Warnings

Do not use

- more than the recommended dosage.
- If drowsiness occurs, do not drive or operate machinery.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for two weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacists before taking this product.

Ask a doctor before use is you have

- a chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema.
- difficulty in urination due to enlargement of the prostate gland.
- Glaucoma.
- Thyroid disease.
- a cough that occurs with too much phlegm (mucus).

Stop use and ask a doctor if

- Cough lasts more than seven days, returns or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding,

Do Not Use.

Keep out of reach of children.

In case of accidental overdose get medical help or contact a poison control center right away.

Directions

- Follow Dosage Chart.
- Do not take more than 6 doses in any 24 hour period.
- This product is not intended for use in children under 12 years of age.

age	dose
Adults and children over 12 years old	2 teaspoons (2 tsp. 10 ml) every 4 hours
Children Under 12 years old	DO NOT USE

Other Information

- Store at 20° – 25° C (68° – 77° F).
- Do not refrigerate.
- Dosage cup provided.

Inactive Ingredients

Ammonium Glycyrrhizate, Citric Acid, Clear Cherry-Strawberry Flavor, Glycerin, Hydroxyethylcellulose, Menthol, Methyl Paraben, Potassium Benzoate, Propyl Paraben, Propylene Glycol, Purified Water.

Principal Display Panel

NDC 0372-0043-04

SCOT-TUSSIN

DIABETES

COUGH SUPPRESASANT Formula with DM

4 FL OZ (118 ml)

SUGAR-FREE	SUGAR-FREE	SUGAR-FREE	SUGAR-FREE										
<h1>SCOT-TUSSIN®</h1>	<h1>SCOT-TUSSIN®</h1>	ADULT <small>NDC 0372-0043-04</small> For Diabetics since 1956 <h1>SCOT-TUSSIN®</h1>	<h1>SCOT-TUSSIN®</h1>										
<p>Drug Facts (Continued)</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> cough lasts more than seven days, returns or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition. <p>If pregnant or breast-feeding. Do Not Use. Keep out of reach of children. In case of accidental overdose get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> follow Dosage Chart. do not take more than 6 doses in any 24 hour period. this product is not intended for use in children under 12 years of age. <table border="1"> <thead> <tr> <th>age</th> <th>dose</th> </tr> </thead> <tbody> <tr> <td>Adults and children over 12 years old</td> <td>2 teaspoons (2 tsp, 10 mL) every 4 hours</td> </tr> <tr> <td>Children under 12 years old</td> <td>DO NOT USE</td> </tr> </tbody> </table> <p>Other information</p> <ul style="list-style-type: none"> store at 20°-25° C (68°-77° F). do not refrigerate. dosage cup provided. <p>Inactive ingredients</p> <p>ammonium glycyrrhizate, citric acid, clear cherry-strawberry flavor, glycerin, hydroxyethylcellulose, menthol, methyl-paraben, potassium benzoate, propyl-paraben, polyethylene glycol, purified water.</p>	age	dose	Adults and children over 12 years old	2 teaspoons (2 tsp, 10 mL) every 4 hours	Children under 12 years old	DO NOT USE	<p>SCOT-TUSSIN QUALITY ASSURANCE</p> <p>Decades of experience since 1956 in providing you with quality medicines goes into every product. Scot-Tussin has been an innovator in the formulation of sugar-free and alcohol-free medicines for the diabetic consumer and has been fortunate to have received recommendations from generations of doctors. Scot-Tussin has been and still is committed to the often over-looked special needs patient. Our medicines are safe for the typical consumer as well as those with diabetes, celiac sprue, high blood pressure, and heart disease.</p> <p>All Scot-Tussin Brand Medicines are Satisfaction Guaranteed or Your Money Back</p> <p>Distributed by: Scot-Tussin Pharmacal Co., Inc. Cranston, RI 02920 U.S.A. Toll Free # 1-800-638-7268 (Scot) 1-401-942-8555 © Copyright 1983 © TM Reg. US Patent Office #657-674</p>	<p>DIABETES</p> <p>COUGH SUPPRESSANT Formula with DM</p> <p>DEXTROMETHORPHAN HBr (Cough Suppressant)</p> <p>FAST RELIEF OF: DRY HACKING COUGH</p> <ul style="list-style-type: none"> FIRST SUGAR & ALCOHOL-FREE HIGH BLOOD PRESSURE SAFE LACTOSE & GLUTEN-FREE SODIUM-FREE DYE-FREE <p>LIQUID 4 FL OZ (118 ml)</p>	<p>IF TAMPER-EVIDENT SEAL AROUND CAP IS BROKEN OR MISSING, DO NOT USE.</p> <p>Drug Facts</p> <table border="1"> <thead> <tr> <th>Active ingredients</th> <th>Purpose</th> </tr> </thead> <tbody> <tr> <td>(in each 5 mL, tsp, teaspoon) Dextromethorphan HBr, USP 10 mg</td> <td>Cough Suppressant</td> </tr> </tbody> </table> <p>Uses</p> <ul style="list-style-type: none"> temporarily quiets and calms a dry cough due to minor throat and bronchial irritation. <p>Warnings</p> <p>Do not use</p> <ul style="list-style-type: none"> more than the recommended dosage. if drowsiness occurs, do not drive or operate machinery. if you are taking a prescription monoamine oxidase inhibitor (MAOI); (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> a chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema. difficulty in urination due to enlargement of the prostate gland. glaucoma. thyroid disease. a cough that occurs with too much phlegm (mucus). 	Active ingredients	Purpose	(in each 5 mL, tsp, teaspoon) Dextromethorphan HBr, USP 10 mg	Cough Suppressant
age	dose												
Adults and children over 12 years old	2 teaspoons (2 tsp, 10 mL) every 4 hours												
Children under 12 years old	DO NOT USE												
Active ingredients	Purpose												
(in each 5 mL, tsp, teaspoon) Dextromethorphan HBr, USP 10 mg	Cough Suppressant												

SCOT-TUSSIN DIABETES COUGH FORMULA WITH DM

dextromethorphan hydrobromide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0372-0043
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
AMMONIUM GLYCYRRHIZATE (UNII: 3VRD35U26C)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYDROXYETHYL CELLULOSE (2000 MPAS AT 1%) (UNII: S38J6RZN16)	
MENTHOL (UNII: L7T10EIP3A)	

METHYLPARABEN (UNII: A2I8C7HI9T)	
POTASSIUM BENZOATE (UNII: 763YQN2K7K)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY, STRAWBERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0372-0043-04	118 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	03/18/1998	

Labeler - SCOT-TUSSIN Pharmaca Co., Inc. (001203918)