

**VICKS CHILDRENS COUGH CONGESTION- phenylephrine hcl,
dextromethorphan hbr, guaifenesin liquid
The Procter & Gamble Manufacturing Company**

VICKS®

children's Cough Congestion

Drug Facts

Active ingredients (in each 15 mL)

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Purpose

Cough suppressant

Expectorant

Nasal decongestant

Uses

temporarily relieves common cold symptoms:

- nasal congestion
- cough due to minor throat and bronchial irritation
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- a sodium-restricted diet

When using this product, do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not improve within 7 days or occur with a fever
- cough persists for more than 7 days, comes back or occurs with a fever, rash or persistent headache

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

| | |
|---------------------------------|--------------------|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 6 to under 12 yrs | 15 mL every 4 hrs |
| children 4 to under 6 yrs | 7.5 mL every 4 hrs |
| children under 4 yrs | do not use |

Other information

- **each 15 mL contains:**sodium 44 mg
- store at no greater than 25°C.

Inactive ingredients

anhydrous citric acid, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

Questions?

1-800-362-1683

TAMPER EVIDENT: Do not use if printed safety seal under cap is missing or damaged.

DIST. BY: PROCTER & GAMBLE, CINCINNATI, OH 45202

Made in Canada

PRINCIPAL DISPLAY PANEL - 177 ml Bottle Label

VICKS®

children's

Cough

Congestion

Non-Drowsy

Phenylephrine HCl - nasal decongestant

Dextromethorphan HBr - cough suppressant

Guaifenesin - expectorant

FREE of:

Artificial Dyes & Flavors,

High Fructose Corn Syrup & Alcohol

- Cough
- Chest congestion
- Stuffy nose

Ages 4+

6 FL OZ (177 ml)



| Drug Facts | |
|--|--------------------|
| Active ingredients (in each 15 mL) | Purpose |
| Dextromethorphan HBr 10 mg | Cough suppressant |
| Guaifenesin 200 mg | Expectorant |
| Phenylephrine HCl 5 mg | Nasal decongestant |
| Uses | |
| temporarily relieves common cold symptoms: • nasal congestion • cough due to minor throat & bronchial irritation | |
| • helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive. | |
| Warnings | |
| Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. | |
| Ask a doctor before use if you have | |
| • heart disease • high blood pressure • thyroid disease • diabetes | |
| • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) | |
| • persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema | |
| • a sodium-restricted diet | |
| When using this product, do not use more than directed. | |
| Stop use and ask a doctor if | |
| • you get nervous, dizzy or sleepless | |
| • symptoms do not improve within 7 days or occur with a fever | |
| • cough persists for more than 7 days, comes back or occurs with a fever, rash or persistent headache | |
| These could be signs of a serious condition. | |
| If pregnant or breast-feeding, ask a health professional before use. | |
| Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. ▶ | |

| Drug Facts (continued) | |
|---|----------------------------------|
| Directions | |
| • take only as directed | • only use the dose cup provided |
| • do not exceed 4 doses per 24 hrs | |
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 6 to under 12 yrs | 15 mL every 4 hrs |
| children 4 to under 6 yrs | 7.5 mL every 4 hrs |
| children under 4 yrs | do not use |
| Other information | |
| • each 15 mL contains: sodium 44 mg | |
| • Store at no greater than 25°C. | |
| Inactive ingredients | |
| anhydrous citric acid, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose | |
| Questions? 1-800-362-1683 | |



VICKS CHILDRENS COUGH CONGESTION

phenylephrine hcl, dextromethorphan hbr, guaifenesin liquid

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:69423-982 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|----------------------------------|--------------------|
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg in 15 mL |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 5 mg in 15 mL |
| GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) | GUAIFENESIN | 200 mg in 15 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| WATER (UNII: 059QF0KO0R) | |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY) | |
| SODIUM CITRATE (UNII: 1Q73Q2JULR) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |

SUCRALOSE (UNII: 96K6UQ3ZD4)

SORBITOL (UNII: 506T60A25R)

SODIUM CHLORIDE (UNII: 451W47IQ8X)

Product Characteristics

| | | | |
|-----------------|---------------|---------------------|--|
| Color | white (Clear) | Score | |
| Shape | | Size | |
| Flavor | STRAWBERRY | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:69423-982-06 | 177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 04/22/2019 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012 | 04/22/2019 | |

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-----------------|---------|-----------|----------------------------|
| SPERA NEXUS INC | | 097722284 | api manufacture(69423-982) |

Revised: 10/2025

The Procter & Gamble Manufacturing Company