

VICKS NYQUIL KIDS HONEY COLD COUGH CONGESTION- doxylamine succinate, phenylephrine hydrochloride and dextromethorphan hydrobromide liquid
The Procter & Gamble Manufacturing Company

Vicks[®] NyQuil Kids Honey Cold & Cough + Congestion

Drug Facts

Active ingredients (in each 15 mL)

Dextromethorphan HBr 10 mg

Doxylamine succinate 6.25 mg

Phenylephrine HCl 5 mg

Purpose

Cough suppressant

Antihistamine

Nasal decongestant

Uses

temporarily relieves common cold symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- heart disease
- high blood pressure

- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers

When using this product

- **do not use more than directed.**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not improve within 7 days or occur with a fever
- cough persists for more than 7 days, comes back or occurs with a fever, rash or persistent headache

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take only as directed
- only use dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	do not use unless directed by a doctor
children under 4 yrs	do not use

Other information

- **each 15 mL contains:**sodium 14 mg
- store at no greater than 25°C

Inactive ingredients

citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6, flavor (with honey), glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions?**1-800-362-1683****Tamper Evident: Do not use if printed shrinkband seal around the neck is broken or missing.****DIST. BY: PROCTER & GAMBLE, CINCINNATI, OH 45202****PRINCIPAL DISPLAY PANEL - 236 ml Bottle Label****VICKS®**

NyQuil™

Kids

HONEY

COLD & COUGH + CONGESTION

Doxylamine Succinate, Phenylephrine HCl, Dextromethorphan HBr

Sneezing, Runny Nose

Nasal Congestion

Cough

AGES 6+

8 FL OZ (236 mL)

Nighttime Relief

FREE OF ALCOHOL

& ACETAMINOPHEN



Drug Facts (continued)

- cough persists for more than 7 days, comes back or occurs with a fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	do not use unless directed by a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 14 mg
- store at no greater than 25°C

Inactive ingredients citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6, flavor (with honey), glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions? 1-800-362-1683

Drug Facts (continued)

Warnings
Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleep

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- do not use more than directed.
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not improve within 7 days or occur with a fever



VICKS NYQUIL KIDS HONEY COLD COUGH CONGESTION

doxylamine succinate, phenylephrine hydrochloride and dextromethorphan hydrobromide liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69423-983
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII: 7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1W5297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5YI2) (DOXYLAMINE - UNII: 95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg in 15 mL

Inactive Ingredients	
Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

SORBITOL (UNII: 506T60A25R)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	brown	Score	
Shape		Size	
Flavor	HONEY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-983-08	236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/07/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/07/2021	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 10/2024

The Procter & Gamble Manufacturing Company