

TUSSIN COUGH EXPECTORANT COUGH SUPPRESSANT DM- dextromethorphan hydrobromide, guaifenesin liquid
Salado Sales, Inc.

Tussin Cough DM 503

Drug Facts Active ingredients (in each 5 mL tsp)

Dextromethorphan HBr, USP 10 mg

Guaifenesin, USP 100 mg

Purpose

Cough Suppressant
Expectorant

Keep out of reach of children.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use

- in a child under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

Ask A Doctor Before You Use if You Have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop Use And Ask A Doctor If

cough lasts more than 7 days, comes back or is accompanied by fever, rash or

persistent headache. These could be signs of a serious condition.

If Pregnant Or Breast-Feeding

ask a health professional before use.

Directions

- do not take more than 6 doses in any 24-hour period

Age	Dose
adults & children 12 years and over	2 teaspoonfuls every 4 hours
children under 12 years	do not use

Other information

- store at 20-25° C (66-77° F)
- do not refrigerate
- dosage cup provided
- sodium 3 mg per teaspoon

Inactive ingredients

Anhydrous citric acid, dextrose, FD & C red # 40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate

Questions?

Call weekdays from 9:30 AM to 4:30 PM EST at

1-877-798-5944

Tussin Cough DM product label

*** COMPARE TO ROBITUSSIN® PEAK COLD ACTIVE INGREDIENTS**

CVP® CONSUMER VALUE PRODUCTS

TUSSIN COUGH EXPECTORANT COUGH SUPPRESSANT DM

NON-DROWSY COUGH FORMULA

- ADULT
- FOR AGES 12 AND OVER
- CONTROLS COUGH
- HELPS LOOSEN CHEST CONGESTION

4 FL OZ (118 ML)

* This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin® Peak Cold


Box-001 Rev 01

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Distributed by: Salado Sales, Inc. P.O. Box 6115 Temple, TX 76503
www.saladosales.com

LOT.

EXP.



Drug Facts

Active ingredients (in each 5 mL tsp)	Purpose
Dextromethorphan HBr, USP 10 mg...Cough Suppressant	
Guaifenesin, USP 100 mg.....Expectorant	

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use

- in a child under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- do not take more than 6 doses in any 24-hour period

Age	Dose
adults & children 12 years & over	2 teaspoonfuls every 4 hours
children under 12 years	do not use

Other information

- store at 20-25°C (68-77°F)
- do not refrigerate
- dosage cup provided
- sodium 3 mg per teaspoonful

Inactive ingredients Anhydrous citric acid, dextrose, FD&C red # 40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate.

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944.


* This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin® Peak Cold.
BX-001-Rev 01

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Distributed by: Salado Sales, Inc
P.O. Box 6115 Temple, TX 76503
www.saladosales.com



* COMPARE TO ROBITUSSIN® PEAK COLD ACTIVE INGREDIENTS



**EXPECTORANT
COUGH SUPPRESSANT
DM**

**NON-DROWSY
COUGH FORMULA**

- ADULT
- FOR AGES 12 & OVER
- CONTROLS COUGH
- HELPS LOOSEN CHEST CONGESTION

4 FL OZ (118mL)

* COMPARE TO ROBITUSSIN® PEAK COLD ACTIVE INGREDIENTS

CVP

TUSSIN COUGH

EXPECTORANT
COUGH SUPPRESSANT
DM

**NON-DROWSY
COUGH FORMULA**

- ADULT
- FOR AGES 12 & OVER
- CONTROLS COUGH
- HELPS LOOSEN CHEST CONGESTION

4 FL OZ (118mL)

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Drug Facts

Active ingredients (in each 5 mL tsp)
Dextromethorphan HBr, USP 10 mg...Cough Suppressant
Guaifenesin, USP 100 mg.....Expectorant

Purpose
Cough Suppressant
Expectorant

Uses
temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold

Warnings
Do not use • in a child under 12 years of age • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have • cough that occurs with too much phlegm (mucus) such as occurs with smoking, asthma, chronic bronchitis or emphysema
Stop use and ask a doctor if cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
• do not take more than 6 doses in any 24-hour period

Age	Dose
adults & children 12 years & over	2 teaspoonfuls every 4 hours
children under 12 years	do not use

Other information
• store at 20-25°C (68-77°F) • do not refrigerate
• dosage cup provided • sodium 3 mg per teaspoonful

Inactive ingredients Anhydrous citric acid, dextrose, FD&C red # 40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate.

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944.

• This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin® Peak Cold.

LR-001 REV01

LOT: _____

EXP: _____

7 61706 16133 3

Distributed by: Salado Sales, Inc
P.O. Box 6115 Temple, TX 76503
www.saladosales.com

res

CVP

TUSSIN COUGH

EXPECTORANT
COUGH SUPPRESSANT
DM

Drug Facts

Active ingredients (in each 5 mL tsp)
Dextromethorphan HBr, USP 10 mg...Cough Suppressant
Guaifenesin, USP 100 mg.....Expectorant

Purpose
Cough Suppressant
Expectorant

Uses
temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings
Do not use • in a child under 12 years of age • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have • cough that occurs with too much phlegm (mucus) • cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema
Stop use and ask a doctor if cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions
do not take more than 6 doses in any 24-hour period

Age	Dose
adults & children 12 years & over	2 teaspoonfuls every 4 hours
children under 12 years	do not use

Other information
• store at 20-25°C (68-77°F) • do not refrigerate
• dosage cup provided • sodium 3 mg per teaspoonful

Inactive ingredients Anhydrous citric acid, dextrose, FD&C red # 40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate.

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944.

* This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin® Peak Cold.

BX-001-Rev 01

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Distributed by: Salado Sales, Inc
P.O. Box 6115 Temple, TX 76503
www.saladosales.com

7 61706 16133 3

* COMPARE TO ROBITUSSIN® PEAK COLD ACTIVE INGREDIENTS

CVP

TUSSIN COUGH

EXPECTORANT
COUGH SUPPRESSANT
DM

**NON-DROWSY
COUGH FORMULA**

- ADULT
- FOR AGES 12 & OVER
- CONTROLS COUGH
- HELPS LOOSEN CHEST CONGESTION

4 FL OZ (118mL)

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/01/2012	

Labeler - Salado Sales, Inc. (009830555)

Registrant - AptaPharma Inc. (790523323)

Establishment

Name	Address	ID/FEI	Business Operations
AptaPharma Inc.		790523323	manufacture(57243-503)

Revised: 12/2023

Salado Sales, Inc.