

**NON-DROWSY DAYTIME COLD AND FLU MAX SOFTGELS MAXIMUM STRENGTH- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled**  
**TOPCO ASSOCIATES LLC**

-----

**NON-DROWSY DAYTIME**

**Cold and Flu Max**

**Softgels**

**MAXIMUM STRENGTH**

***Active ingredients (in each softgel)***

Acetaminophen 325 mg

Dextromethorphan hydrobromide 10 mg

Phenylephrine hydrochloride 5 mg

***Purposes***

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

***Uses***

- temporarily relieves these symptoms due to a cold or flu:
- minor aches and pains
- headache
- cough
- sore throat
- nasal and sinus congestion
- temporarily reduces fever

***Warnings***

**Liver warning:**

This product contains acetaminophen.  
Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

### **Allergy alert:**

Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash n hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

### **Sore throat warning:**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

### **Ask a doctor before use if you have**

- liver disease n heart disease n high blood pressure
- thyroid disease n diabetes
- cough with excessive phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

### **Stop use and ask a doctor if**

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

- nervousness, dizziness, or sleeplessness occurs
- If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions***

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 softgels with water every 4 hours. Do not exceed 10 softgels in 24 hours or as directed by a doctor.
- children under 12 years: do not use

***Other information***

- store at room temperature. Avoid temperatures above 25°C (77°F).

***Inactive ingredients***

FD&C yellow #6, gelatin, glycerin, pearlin silver, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, titanium dioxide

***Questions or comments?***

**1-888-423-0139**

**Carton**

TopCare health

NDC 36800-720-24

COMPARE TO ALKA-SELTZER® PLUS®  
MAXIMUM STRENGTH DAYTIME COLD & FLU  
POWERMAX™ GELS ACTIVE INGREDIENTS\*

NON-DROWSY DAYTIME

Cold & Flu Max Softgels  
MAXIMUM STRENGTH

PAIN RELIEVER-FEVER REDUCER - ACETAMINOPHEN  
COUGH SUPPRESSANT - DEXTROMETHORPHAN HBr  
NASAL DECONGESTANT - PHENYLEPHRINE HCl

RELIEVES:

- Nasal Congestion
  - Cough
  - Headache & Body Ache
  - Sore Throat
  - Sinus Pressure
- Smaller Capsule •Same Strength

24SOFTGELS\*\*

\*\*Liquid-Filled Capsules

minis

**Drug Facts**  
**Active Ingredients (in each softgel)**  
 Acetaminophen 325 mg...Pain reliever/fever reducer  
 Dextromethorphan...Cough suppressant  
 Phenylephrine hydrochloride 5 mg...Nasal decongestant

**Uses**  
 temporarily relieves these symptoms due to a cold or flu: minor aches and pains, headache, cough, sore throat, nasal and sinus congestion, temporarily reduces fever.

**Warnings**  
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen or 3 or more alcoholic drinks every day while using this product. Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: skin redness, hives, rash, itches, facial swelling, asthma (wheezing), shock. If a skin or general allergic reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. If you have ever had an allergic reaction to this product or any of its ingredients, do not use in children under 12 years of age.

**Directions**  
 do not take more than the recommended dose adults and children 12 years and over: take 2 softgels with water every 4 hours. Do not exceed 10 softgels in 24 hours or as directed by a doctor. children under 12 years: do not use

**Other information**  
 store at room temperature. Avoid temperatures above 25°C (77°F).

**Inactive Ingredients** FD&C yellow #8, gelatin, glycerin, pearlin silver, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, titanium dioxide

**Questions or comments?** 1-888-423-0130

**Drug Facts (continued)**  
 Ask a doctor before use if you have liver disease, heart disease, high blood pressure, thyroid disease, diabetes, cough with excessive phlegm (mucus), difficulty in urination due to enlargement of the prostate gland, persistent or chronic cough such as occurs with smoking, asthma, or emphysema. Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. When using this product do not exceed recommended dosage. Stop use and ask a doctor if pain, cough, or nasal congestion gets worse or lasts more than 7 days, fever gets worse or lasts more than 3 days, redness or swelling is present, new symptoms occur, cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition, nervousness, dizziness, or sleeplessness occurs. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**THIS PRODUCT IS PACKAGED IN A CHILD-RESISTANT AND TAMPER-EVIDENT PACKAGE. USE ONLY IF BLISTERS ARE INTACT.**

**TopCare health**  
 NON-DROWSY DAYTIME  
**Cold & Flu Max Softgels**  
 MAXIMUM STRENGTH  
 PAIN RELIEVER- FEVER REDUCER- ACETAMINOPHEN  
 COUGH SUPPRESSANT - DEXTROMETHORPHAN HBr  
 NASAL DECONGESTANT - PHENYLEPHRINE HCl  
**RELIEVES:**  
 • Nasal Congestion  
 • Cough  
 • Headache & Body Ache  
 • Sore Throat  
 • Sinus Pressure  
 •Smaller Capsule •Same Strength actual size  
**24 SOFTGELS\*\***  
 \*\*Liquid-Filled Capsules  
**minis**  
 COMPARE TO ALKA-SALTZER PLUS® MAXIMUM STRENGTH DAYTIME COLD & FLU POWERMAX™ GELS ACTIVE INGREDIENTS?  
 DISTRIBUTED BY TOPCO ASSOCIATES LLC ELK GROVE VILLAGE, IL 80007 © TOPCO SPL0322 QUESTIONS? 1-888-423-0130 topcare@topco.com www.topcarebrand.com MADE IN INDIA  
 Scan here for more information or call 1-888-423-0130  
 QUALITY GUARANTEED  
 This TopCare® product is laboratory tested to guarantee its highest quality. Your total satisfaction is guaranteed.  
 \*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Alka-Seltzer PLUS® POWERMAX™ GELS.  
 NON-DROWSY DAYTIME  
**Cold & Flu Max Softgels**  
 MAXIMUM STRENGTH  
 •Smaller Capsule •Same Strength  
**minis**  
 (HOW TO USE) (OVERDOSE) (ACETAMINOPHEN) (COUGH SUPPRESSANT - DEXTROMETHORPHAN HBr) (NASAL DECONGESTANT - PHENYLEPHRINE HCl)

## NON-DROWSY DAYTIME COLD AND FLU MAX SOFTGELS MAXIMUM STRENGTH

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:76162-720
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1VS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POTASSIUM ALUMINUM DISILICATE</b> (UNII: SRB14JRX6C)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	OVAL ((oblong))	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	105
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76162-720-24	1 in 1 CARTON	04/20/2022	
1		24 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/20/2022	

**Labeler** - TOPCO ASSOCIATES LLC (006935977)