

**NITETIME MULTI-SYMP TOM COLD/FLU RELIEF- acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate capsule, liquid filled
Greenbrier International**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

NiteTime Multi-Symptom Liquid Capsules Cold/Flu Relief

Drug Facts

Active Ingredient

Active ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Purpose

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves common cold/flu symptoms

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever
- runny nose & sneezing

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Sore throat warning

If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleep

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, & tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see Overdose warning
- do not exceed 4 doses per 24 hrs

Adults & children 12 years and over	2 softgels with water every 6 hrs
Children 4 to under 12 years	ask a doctor
Children under 4 years	do not use

Other information

- store at room temperature

Inactive ingredients

D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, methylparaben, polyethylene glycol, povidone, propylene glycol, propylparaben, purified water, sorbitol, titanium dioxide

Questions?

1-866-467-2748

This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® Nyquil® Cold & Flu Liquicaps®

PRINCIPAL DISPLAY PANEL - 10 Softgel Carton

Assured

Multi-Symptom

NiteTime

Acetaminophen, Doxylamine, Dextromethorphan HBr

Cold/Flu Relief

- Aches & Sore Throat
- Sneezing, Runny Nose
- Cough

10 Softgels

Multi-Symptom
NiteTime
 Liquid Capsules

Cold/Flu Relief

10 SOFTGELS



NDC 33992-0621-0
 *Compare to the active ingredients in Vicks Nyquil®

Multi-Symptom
NiteTime
 Liquid Capsules

Acetaminophen, Doxylamine, Dextromethorphan HBr

Cold/Flu Relief

- Aches, Fever & Sore Throat
- Sneezing, Runny Nose
- Cough



10 SOFTGELS

Multi-Symptom
NiteTime
 Liquid Capsules

Cold/Flu Relief

- Aches, Fever & Sore Throat
- Sneezing, Runny Nose • Cough

10 SOFTGELS

06210DTRFC



6 39277 06948 1

Lot:

Exp:

TAMPER EVIDENT: This package is safety sealed & child resistant. Use only if blisters are intact. If difficult to open, use scissors.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Distributed by:
 Greenbrier International, Inc.
 Chesapeake, VA 23320

Warnings This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses in 24 hours, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product ■ Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly. ■ Do not use ■ with any other products containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ if you are now taking a prescription monoamine oxidase inhibitor or (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drugs. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product ■ to make a child sleep ■ Ask a doctor before use if you have ■ liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ breathing problems or chronic cough that lasts or as occurs ■ with smoking, asthma, chronic bronchitis, or emphysema ■ trouble urinating due to enlarged prostate gland ■ Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

Directions ■ take only as directed - see Overdose warning ■ do not exceed 4 doses per 24 hours

Adults and children	2 softgels with water every 6 hours
Children 4 to 12 years and over	ask a doctor
Children under 4 years	do not use

Other information ■ store at room temperature

Inactive ingredients D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, methylparaben, polyethylene glycol, polydioxane, propylene glycol, povidone, purified water, sorbitol, titanium dioxide

Questions or comments? 1-866-487-2748

* This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks Nyquil® Cold & Flu LiquidCaps®

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:33992-0621-0	1 in 1 CARTON		
1		10 in 1 BLISTER PACK		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	01/04/2013	

Labeler - Greenbrier International (610322518)

Revised: 1/2014

Greenbrier International