CHILDRENS MULTI-SYMPTOM COLD - guaifenesin, dextromethorphan hydrobromide, phenylephrine hydrochloride liquid Chain Drug Consortium, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active ingredients

Drug Facts

Active ingredients

(in each 5 mL teaspoon)

Guaifenesin 100 mg Dextromethorphan HBr 5 mg Phenylephrine HCL 2.5 mg

Purpose

Purpose

Cough suppressant Expectorant Nasal decongestant

Keep out of reach of children

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control center right away.

Uses

Uses

- helps to loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passage ways of bothersome mucus and make coughs more productive
- temporarily relieves these symptoms occurring with a cold:
- cough due to minor throat and bronchial irritation
- nasal congestion
- stuffy nose

Warnings

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for two weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or a pharmacist before taking this product.

Ask a doctor before use

Ask a doctor before use if the child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

When using this product

When using this product

• do not use more than directed

Stop use and ask a doctor

Stop use and ask a doctor if

- your child gets nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache.

These could be signs of a serious condition.

Directions

Directions

• do not take more than 6 doses in any 24-hour period

Age Children 6 years to under 12 years Children 4 years to under 6 years Children under 4 years Dose

2 teaspoons every 4 hours 1 teaspoon every 4 hours Do not use

Other information

Other information

- each 5 mL teaspoon contains: sodium 3 mg
- store between 15-30 ° C (59-86° F)
- do not refrigerate
- dosing cup provided

Inactive ingredients

Inactive ingredients

citric acid anhydrous, D and C red #33,dextrose, FD and C blue #1, FD and C red #40, flavors, glycerin, maltitol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sucralose, xanthan gum

Questions?

Questions?

Call weekdays from 9:30 AM to 4:30 PM EST at

1-877-798-5944

Product Label

NDC 68016-134-00

*COMPARE TO THE ACTIVE INGREDIENTS IN CHILDREN'S MUCINEX®Multisymptom Cold

Premier Value®

CHILDREN'S MULTI- SYMPTOM COLD

EXPECTORANT COUGH SUPPRESSANT NASAL DECONGESTANT

Guaifenesin 100 mg Dextromethorphan HBr 5 mg Phenylephrine HCL 2.5 mg

- * Stuffy Nose
- * Cough
- * Chest Congestion
- * Breaks up Mucus

Alcohol free

For ages 4 to 12

Very Berry Flavor 4 FL OZ (118mL)

INDEPENDENTLY TESTED SATISFACTION GUARANTEED PV DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

*This product is not manufactured or distributed by Reckitt Benckiser Inc. distributor of Children's Mucinex® Multisymptom Cold

If for Any reason you are not satisfied with this product, lease return it to the store where purchased for a full refund.

DISTRIBUTED BY CHAIN DRUG CONSORTIUM 3301 NW BOCA RATON BLVD SUITE 101, BOCA RATON, FL 33431 BX-010

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Drug Facts

Active ingredients Purpose (in each 5 mL teaspoonful)

Dextromethorphan HBr, USP 5 mg.....Cough suppressant Guaifenesin, USP 100 mg......Expectorant Phenylephrine HCl, USP 2.5 mg......Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves these symptoms occurring with a cold:
 - cough due to minor throat and bronchial irritation
 - nasal congestion
 - stuffy nose

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

When using this product

do not use more than directed

Stop use and ask a doctor if

- your child gets nervous, dizzy or sleepless
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache.
 These could be signs of a serious condition.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

do not take more than 6 doses in any 24-hour period

age	dose
Children 6 years	2 teaspoonfuls every
to under 12 years	4 hours.
Children 4 years	1 teaspoonful every
to under 6 years	4 hours
Children under 4 years	do not use

Other information

- each 5 mL teaspoonful contains: sodium 3 mg
- store between 15-30°C (59-86°F)
- do not refrigerate
- dosing cup provided

Inactive ingredients

anhydrous citric acid, D&C red #33, dextrose, FD&C blue #1, FD&C red #40, flavors, glycerin, maltilol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sucralose, xanthan gum

Questions? Call weekdays from 9:30 am to 4:30 pm EST **1-877-798-5944**


*This product is not manufactured or distributed by Reckitt Benckiser Inc., distributor of Children's Mucinex® Multisymptom Cold



If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

DISTRIBUTED BY CHAIN DRUG CONSORTIUM 3301 NW BOCA RATON BLVD SUITE 101, BOCA RATON, FL 33431

BX-010



CHILDREN'S MULTI-SYMPTOM COLD

EXPECTORANT COUGH SUPPRESSANT NASAL DECONGESTANT

Guaifenesin 100 mg Dextromethorphan HBr 5 mg Phenylephrine HCl 2.5 mg

- Stuffy Nose
- Cough
- Chest Congestion
- Breaks up Mucus

Alcohol free For ages 4 to 12

Very Berry Flavor



CHILDRENS MULTI-SYMPTOM COLD

guaifenesin, dextromethorphan hydrobromide, phenylephrine hydrochloride liquid

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:68016-134

Route of Administration ORAL

Active Ingredient/Active Moiety **Ingredient Name Basis of Strength** Strength 100 mg GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) **GUAIFENES IN** in 5 mL DEXTRO METHO RPHAN HYDRO BRO MIDE (UNII: 9 D2RTI9 KYH) **DEXTROMETHORPHAN** 5 mg (DEXTROMETHORPHAN - UNII:7355X3ROTS) **HYDROBROMIDE** in 5 mLPHENYLEPHRINE HYDRO CHLO RIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE -PHENYLEPHRINE 2.5 mg UNII:1WS297W6MV) HYDROCHLORIDE in 5 mL

Ingredient Name	Strength	
ANHYDRO US CITRIC ACID (UNII: XF417D3PSL)		
D&C RED NO. 33 (UNII: 9DBA0SBB0L)		
DEXTROSE (UNII: IY9 XDZ35W2)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
GLYCERIN (UNII: PDC6 A3C0 OX)		
MALTITOL (UNII: D65DG142WK)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		
SACCHARIN SO DIUM (UNII: SB8 ZUX40 TY)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SUCRALOSE (UNII: 96K6UQ3ZD4)		
XANTHAN GUM (UNII: TTV12P4NEE)		

Product Characteristics			
Color		Score	
Shape		Size	
Flavor	BERRY (VERY BERRY)	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-134-00	118 mL in 1 BOTTLE		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/27/2012	

Labeler - Chain Drug Consortium, LLC (101668460)

Registrant - AptaPharma Inc. (790523323)

Establishment				
Name	Address	ID/FEI	Business Operations	
AptaPharma Inc.		790523323	manufacture(68016-134)	