# ADVIL LIQUIGELS- ibuprofen capsule, liquid filled Navajo Manufacturing Company Inc.

-----

#### **Advil Liquigels**

#### **Drug Facts**

## Active ingredient (in each capsule)

Solubilized Ibuprofen equal to 200 mg ibuprofen (NSAID)\* (present as the free acid and potassium salt)
\*nonsteroidal anti-inflammatory drug

#### **Purpose**

Pain Reliever/Fever Reducer

#### Uses

- temporarily relieves minor aches and pains due to:
- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

#### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause

severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

#### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

#### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma
- you are taking a diuretic

#### Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

### When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

### Stop use and ask a doctor if

- you expereince any of the following signs of stomach bleeding:
- feel faint vomit blood
- have bloody or black stools
   have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

### If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

#### Other information

- each capsule contains: potassium 20 mg
- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

#### Inactive ingredients

coconut oil, FD&C green no. 3, gelatinm lecithin (soyabean), pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol

#### Questions or comments?

Call toll free 1-800-88-ADVIL

#### Package Labeling:



## ibuprofen capsule, liquid filled

## **Product Information**

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:67751-148(NDC:0573-0169)
--------------	----------------	--------------------	------------------------------

**Route of Administration** ORAL

## **Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients			
Ingredient Name	Strength		
COCONUT OIL (UNII: Q9L0O73W7L)			
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)			
GELATIN (UNII: 2G86QN327L)			
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDWIA)			
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)			
WATER (UNII: 059QF0KO0R)			
SORBITAN (UNII: 6O92ICV9RU)			
SORBITOL (UNII: 506T60A25R)			

Product Characteristics			
Color	green (Transparent green to blue green)	Score	no score
Shape	CAPSULE (oblong softgel)	Size	20mm
Flavor		Imprint Code	Advil
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:67751-148- 01	1 in 1 CARTON	09/20/2016	09/20/2016	
1		4 in 1 POUCH; Type 0: Not a Combination Product			
2	NDC:67751-148- 02	1 in 1 CARTON; Type 0: Not a Combination Product	06/13/2018		
3	NDC:67751-148- 03	2 in 1 CARTON; Type 0: Not a Combination Product	09/20/2016		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020402	09/20/2016	

## Labeler - Navajo Manufacturing Company Inc. (091917799)

Establishment			
Name	Address	ID/FEI	<b>Business Operations</b>
Navajo Manufacturing Company Inc.		136941411	relabel(67751-148), repack(67751-148)

Revised: 3/2023 Navajo Manufacturing Company Inc.