

SOUND BODY COLD AND FLU RELIEF- acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

Big Lots Stores, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Big Lots, Inc Cold & Flu Relief Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed

by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you

do not notice any signs or symptoms.

Directions

- take only as directed – see overdose warning
- do not exceed 4 doses per 24 hours

adults & children 12 yrs & over	2 softgels with water every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

D&C yellow no. 10, edible ink*, FD&C blue no. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution *may contain this ingredient

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to the Active Ingredients in Vicks[®] NyQuil[®] Cold & Flu

NIGHTTIME

Cold & Flu Relief

Pain Reliever / Fever Reducer – Acetaminophen

Cough Suppressant – Dextromethorphan HBr

Antihistamine – Doxylamine Succinate

24 SOFTGELS

Actual Size

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Distributed By: Perrigo Company
515 Eastern Avenue
Allegan, Michigan 49010
MADE IN CANADA
V#5003185 ITEM#05662



Compare to the Active
Ingredients in Vicks®
NyQuil® Cold & Flu*

NIGHTTIME

Cold & Flu Relief

Pain Reliever / Fever Reducer - Acetaminophen
Cough Suppressant - Dextromethorphan HBr
Antihistamine - Doxylamine Succinate

24 SOFTGELS



Actual Size

CONVENIENT RECLOSING TAB

This product is not manufactured or distributed by Procter & Gamble,
distributor of Vicks® NyQuil® Cold & Flu.

Drug Facts

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

Active ingredients (in each softgel)	Purpose
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 15 mg.....	Cough suppressant
Doxylamine succinate 6.25 mg.....	Antihistamine

Uses temporarily relieves common cold/flu symptoms:
 cough due to minor throat and bronchial irritation minor aches and pains
 runny nose and sneezing sore throat headache fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
 more than 4,000 mg of acetaminophen in 24 hours
 with other drugs containing acetaminophen
 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 skin reddening blisters rash
 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed
 by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use with any other drug containing acetaminophen (prescription or nonprescription).
 If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are
 now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression,
 psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI
 drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before
 taking this product. if you have ever had an allergic reaction to this product or any of its ingredients.
Ask a doctor before use if you have cough that occurs with too much phlegm (mucus)
 a breathing problem such as emphysema or chronic bronchitis liver disease glaucoma
 persistent or chronic cough as occurs with smoking, asthma, or emphysema
 trouble urinating due to an enlarged prostate gland
Ask a doctor or pharmacist before use if you are

8-77F)
dible ink; FD&C blue no. 1, gelatin, glycerin, polyethylene glycol, povidone, polyethylene
*may contain this ingredient

<p>■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin</p> <p>When using this product ■ excitability may occur, especially in children</p> <p>■ be careful when driving a motor vehicle or operating machinery ■ avoid alcoholic drinks</p> <p>■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ marked drowsiness may occur</p> <p>Stop use and ask a doctor if ■ pain or cough gets worse or lasts more than 7 days</p> <p>■ redness or swelling is present ■ fever gets worse or lasts more than 3 days ■ new symptoms occur</p> <p>■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>							
<p>Directions ■ take only as directed - see overdose warning ■ do not exceed 4 doses per 24 hours</p> <table border="1"> <tr> <td>adults & children 12 yrs & over</td> <td>2 softgels with water every 6 hrs</td> </tr> <tr> <td>children 4 to under 12 yrs</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 yrs</td> <td>do not use</td> </tr> </table>		adults & children 12 yrs & over	2 softgels with water every 6 hrs	children 4 to under 12 yrs	ask a doctor	children under 4 yrs	do not use
adults & children 12 yrs & over	2 softgels with water every 6 hrs						
children 4 to under 12 yrs	ask a doctor						
children under 4 yrs	do not use						

Drug Facts (continued)

Other information ■ store at 20-25°C (68-77°F)

Inactive ingredients D&C yellow no. 10, ethyl alcohol, purified water, sorbitol solution

Questions or comments? 1-800-771-9421

GLUTEN FREE

OPEN OTHER END

05662 TM C3

SOUND BODY COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50594-027
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	GREEN (clear)	Score	no score
--------------	---------------	--------------	----------

Shape	OVAL	Size	20mm
Flavor		Imprint Code	056
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50594-027-62	12 in 1 CARTON	03/30/2017	
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	03/30/2017	

Labeler - Big Lots Stores, Inc. (017885351)

Revised: 11/2019

Big Lots Stores, Inc.