TOP QUALITY MFG. TOPICAL ANESTHETIC- benzocaine gel Top Quality Manufacturing, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active Ingredient

Benzocaine 200mg (in each g)

Purpose

Oral Anesthetic

Use

For oral mucosa use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a physician before use. Keep out of reach of children.

Directions

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

Other Information

Store at 59° - 86°F (15°-30°C). Protect from freezing.

Inactive Ingredients

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

Questions or comments?

(800) 483-8559



f pregnant or breast feeding rellow #5, (tartrazine), yellow #6 swelling, rash or fever develops (eep out of reach of children nouth symptoms do not improv in children under 2 years of age Oo not exceed recommended Poison Control Center right away contact occurs, flush with water. Apply only amount needed to contain blue #1, green #3, greeı children under 2 years of age: Questions or comments? Josage. If more than used for lirected by a physician. If sore redness persists or worsens; or pain is accidentally swallowed Store at 59° - 86° F (15 - 30°C). he oral mucosa to prevent or sodium saccharin, water. May get medical help or contact a lavoring, PEG 3350, PEG 400 ee your physician promptly. n 7 days; irritation, pain, or When using this product #5, red #3, red #28, red #40, Woid contact with eyes. If ssk a physician before use. nactive ingredients Other information Protect from freezing. 99291651 as a color additive. · for teething **Directions** do not use

may cause methemoglobinemia a serious condition that must be attention if you or a child in your Benzocaine 200 mg *(in each g)* occur even if you have used this pale, gray or blue colored skir warning: Use of this product reduces the amount of oxyger dizziness or lightheadedness product before. Stop use and Ilergy Alert: Do not use on for more than 7 days unless carried in the blood. This can emporary relief of pain due For oral mucosal use only, as treated promptly because it to minor dental procedures. llergies to local anesthetic directed by dentist. For the such as procaine, butacaine penzocaine or other "caine fatigue or lack of energy patients with a history of Methemoglobinemia seek immediate medical Active ingredients shortness of breath rapid heart rate **Drug Facts** Oral Anesthetic care develops: Do not use headache anesthetics Jurpose cyanosis) ş

TOP QUALITY MFG. TOPICAL ANESTHETIC

benzocaine gel

Product	Inform	ation
Product		ation

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69301-006
--------------	----------------	--------------------	---------------

Route of Administration DENTAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength

BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5) BENZOCAINE 200 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
FD&C YELLOW NO. 5 (UNII: I753WB2F1M)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
D&C GREEN NO. 5 (UNII: 8J6RDU8L9X)	

Product Characteristics

Color	purple	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

# Item Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:69301-006-	30 g in 1 JAR; Type 0: Not a Combination Product	11/01/2018	

Marketing Information

ria keting information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	05/01/2014	

Labeler - Top Quality Manufacturing, Inc. (113630649)

Revised: 1/2022 Top Quality Manufacturing, Inc.