

RYSHI SPF 50 SUNSCREEN- avobenzone 3%, homosalate 15%, octisalate 5%, octocrylene 10% stick

Rite Aid

Sunscreen.

Avobenzone 3%, Homosalate 15%, Octisalate 5%, Octocrylene 10%.

Helps prevent sunburn. If used as directed with other sun protection measures (see **Directions**) decreases the risk of skin cancer and early skin aging caused by the sun.

For external use only. Do not use on damaged or broken skin. **When using this product** keep out of eyes. Rinse with water to remove. **Stop use and ask a doctor** if rash occurs.

If swallowed, get medical help or contact a Poison Control Center right away.

Apply liberally 15 minutes before sun exposure. Reapply after 80 minutes of swimming or sweating, immediately after towel drying, at least every 2 hours.

Sun Protection Measures. Spending time in the sun increased your risk of skin cancer and early skin aging. To decrease this risk, regularly use sunscreen with a Broad Spectrum SPF of 15 or higher and other sun protection measures including: limit time in the sun, especially from 10 am - 2 pm, wear long-sleeved shirts, pants, hats, and sunglasses. Children under 6 months: ask a doctor.

Aloe Barbadensis Leaf Extract, BHT, Caprylic/Capric Triglyceride, Cetyl Alcohol, C12-15 Alkyl Benzoate, C15-19 Alkane, Ethylhexyl Palmitate, Fragrance, Helianthus Annuus (Sunflower) Seed Oil, Ozokerite, Paraffin, Polyester-8, Theobroma Cacao (Cocoa) Seed Butter, VP/Eicosene Copolymer.

ULTIMATE SHEER

SUNSCREEN STICK

50

BROAD SPECTRUM SPF 50

WATER RESISTANT

RYSHI™

WATER RESISTANT (80 MINUTES)

NET WT 1.5 OZ (43 g)

DRUG FACTS

Active ingredients

Avobenzone 3%, Homosalate 15%,
Octisalate 5%, Octocrylene 10%, }

Purpose

Sunscreen

DISTRIBUTED BY:
RITE AID,
30 HUNTER LANE,
CAMP HILL, PA 17011
www.riteaid.com
MADE IN THE USA
WITH DOMESTIC &
FOREIGN
COMPONENTS

**TIME TO SHINE
GUARANTEE**

IF YOU ARE NOT SATISFIED, WE
WILL HAPPILY REFUND YOUR MONEY



PEEL HERE ▼

Drug Facts (continued)

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see **Directions**) decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away

Directions

- apply liberally 15 minutes before sun exposure
- reapply:
 - after 80 minutes of swimming or sweating
 - immediately after towel drying
 - at least every 2 hours

Drug Facts (continued)

Directions (continued)

• Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m.-2 p.m.
- wear long-sleeved shirts, pants, hats and sunglasses
- children under 6 months: ask a doctor

Other information

- may stain some fabrics and surfaces
- protect this product from excessive heat and direct sun

Inactive ingredients

aloe barbadensis leaf extract, BHT, caprylic/capric triglyceride, cetyl alcohol, C12-15 alkyl benzoate, C15-19 alkane, ethylhexyl palmitate, fragrance, helianthus annuus (sunflower) seed oil, ozokerite, paraffin, polyester-8, theobroma cacao (cocoa) seed butter, VP/eicosene copolymer

Questions or comments?

Visit www.riteaid.com

RYSHI SPF 50 SUNSCREEN

avobenzone 3%, homosalate 15%, octisalate 5%, octocrylene 10% stick

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-3115
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AVOBENZONE (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX)	AVOBENZONE	3 g in 100 g
HOMOSALATE (UNII: V06SV4M95S) (HOMOSALATE - UNII:V06SV4M95S)	HOMOSALATE	15 g in 100 g
OCTISALATE (UNII: 4X49Y0596W) (OCTISALATE - UNII:4X49Y0596W)	OCTISALATE	5 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
EICOSYL POVIDONE (UNII: XQQ9MKE2BJ)	
CERESIN (UNII: Q1LS2UJO3A)	
POLYESTER-8 (1400 MW, CYANODIPHENYLPROPENOYL CAPPED) (UNII: T9296U138P)	
ALKYL (C12-15) BENZOATE (UNII: A9EJ3J61HQ)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
C15-19 ALKANE (UNII: CI87N1IM01)	
OCTOCRYLENE (UNII: 5A68WGF6WM)	10 g in 100 g
ETHYLHEXYL PALMITATE (UNII: 2865993309)	
PARAFFIN (UNII: I9O0E3H2ZE)	
ALOE VERA LEAF (UNII: ZY81Z83H0X)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
SUNFLOWER OIL (UNII: 3W1JG795YI)	
COCOA BUTTER (UNII: 512OYT1CRR)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-3115-1	43 g in 1 VIAL, PLASTIC; Type 0: Not a Combination Product	07/05/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	07/05/2022	

Labeler - Rite Aid (014578892)

Registrant - Derma Care Research Labs, LLC (116817470)

Establishment

Name	Address	ID/FEI	Business Operations
Derma Care Research Labs, LLC		116817470	manufacture(11822-3115)

Revised: 12/2024

Rite Aid