RITE AID ANTI-ITCH- diphenhydramine hcl 2% gel Rite Aid

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Rite Aid Anti-Itch Gel, Diphenhydramine HCl 2%

Diphenhydramine HCI 2%

Topical Analgesic.

For the temporary relief of pain and itching associated with insect bites, minor burns, sunburn, minor cuts, scrapes, minor skin irritations, and rashes due to poison ivy, oak, and sumac.

For external use only. Do not use on large areas of the body, with any other products containing diphenhydramine, even one taken by mouth. Ask a doctor before use on chicken pox or measles. When using this product avoid contact with eyes. Stop use and ask a doctor if condition worsens, if symptoms persist for more than 7 days or clear up and occur again within a few days.

If swallowed, get medical help or contact a Poison Control Center right away.

Do not use more than directed. Adults and children 12 years of age and older, apply to the affected area not more than 3 to 4 times daily. Children under 12 years of age: ask a doctor.

Camphor, Citric Acid, Ethylhexylglycerin, Glycerin, Hydroxypropyl Methylcellulose, Phenoxyethanol, Propylene Glycol, SD Alcohol 38-B, Sodium Citrate, Water.



Drug Facts

Active ingredient

Purpose

Diphenhydramine HCI 2%...

Topical Analgesic

Uses

For the temporary relief of pain and itching associated with:

- insect bites
- minor burnş scrapes
- sunburn · minor skin irritations
- minor cuts rashes due to poison ivy, poison oak and poison sumac

Warnings

For external use only

Do not use • on large areas of the body

 with any other products containing diphenhydramine, even one taken by mouth

Ask a doctor before use • on chicken pox • on measles

When using this product avoid contact with eyes

Stop use and ask a doctor if • condition worsens • symptoms persist for more than 7 days or clear up and occur again within a

Keep out of the reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away

Directions

- · do not use more than directed
- · adults and children 2 years or older: apply to affected area not more than 3 to 4 times daily |
 • children under 2 years of age: ask a doctor

Other Information store at room temperature

Inactive ingredients

camphor, citric acid, ethylhexylglycerin, glycerin, hydroxypropyl methylcellulose, phenoxyethanol, propylene glycol, SD alcohol 38-B, sodium citrate, water.



DISTRIBUTED BY: RITE AID, 30 HUNTER LANE, CAMP HILL, PA 17011 www.riteaid.com

SATISFACTION GUARANTEE:

If you're not satisfied, we'll happily refund your money.

RITE AID ANTI-ITCH

diphenhydramine hcl 2% gel

Product Information

HUMAN OTC DRUG NDC:11822-1070 **Product Type** Item Code (Source)

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) DIPHENHYDRAMINE

Inactive Ingredients		
Ingredient Name	Strength	
CAMPHOR (SYNTHETIC) (UNII: 5TJD82A1ET)		
HYPROMELLOSES (UNII: 3NXW29V3WO)		
ALCOHOL (UNII: 3K9958V90M)		
ETHYLHEXYLGLYCERIN (UNII: 147D247K3P)		
GLYCERIN (UNII: PDC6A3C0OX)		
WATER (UNII: 059QF0KO0R)		
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)		
PHENOXYETHANOL (UNII: HIE492ZZ3T)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
SODIUM CITRATE (UNII: 1Q73Q2JULR)		

l	Packaging					
	# Item Code	Package Description	Marketing Start Date	Marketing End Date		
	NDC:11822- 1070-3	103 mL in 1 TUBE; Type 0: Not a Combination Product	03/31/2021			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part348	03/31/2021	
	part348	03/31/2021	

Labeler - Rite Aid (014578892)

Registrant - Derma Care Research Labs, LLC (116817470)

Revised: 7/2022 Rite Aid