

**CORICIDIN HBP DAY AND NIGHT- dextromethorphan hydrobromide, guaifenesin, acetaminophen, chlorpheniramine maleate  
Bayer HealthCare LLC.**

-----  
**Coricidin HBP**

**Day and Night Multi-Symptom Cold**

Coricidin HBP Day

***Drug Facts***

<b><i>Active ingredients (in each softgel)</i></b>	<b><i>Purpose</i></b>
Dextromethorphan hydrobromide 10 mg	Cough Suppressant
Guaifenesin 200 mg	Expectorant

**Uses**

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

**Warnings**

**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- cough that occurs with excessive phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Stop use and ask a doctor if** cough lasts more than 7 days, reoccurs, or occurs with fever, rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Abuse of this product can lead to serious injury.**

**Directions**

- do not exceed recommended dose
- do not take the Day and Night products at the same time; wait 4 hours after the last Night dose before starting Day product
- adults and children 12 years and over: 1 or 2 softgels every 4 hours, not more than 6 softgels in 12 hours
- children under 12 years of age: ask a doctor

### Other Information

- store between 20° to 25°C (68° to 77°F)
- protect from freezing
- protect from excessive moisture

### Inactive Ingredients

FD&C blue no. 1, FD&C red no. 40, gelatin, glycerin, pharmaceutical ink, polyethylene glycol 400, povidone, propylene glycol, sorbitol

Coricidin HBP Night

### Drug Facts

<b>Active ingredients (in each tablet)</b>	<b>Purpose</b>
Acetaminophen 500 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg	Antihistamine
Dextromethorphan hydrobromide 15 mg	Cough Suppressant

### Uses

- temporarily relieves
  - minor aches and pains
  - headache
  - cough
  - runny nose
  - sneezing
- temporarily reduces fever

### Warnings

#### Liver Warning

This product contains acetaminophen.

Sever liver damage may occur if you take

- more than 4 tablets in 12 hours
- with other drugs containing acetaminophen

- 3 or more alcoholic drinks every day while using this product

### **Do not use**

- with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- cough that occurs with excessive phlegm (mucus)
- a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with a rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children. Abuse of this product can lead to serious injury.**

### **Overdose Warning**

Taking more than the recommended dose may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

## Directions

- do not use more than directed (see overdose warning)
- do not take the Day and Night products at the same time; wait 4 hours after the last Day dose before starting Night product
- adults and children 12 years and over: 2 tablets at bedtime and every 6 hours if needed, while symptoms persist, not more than 4 tablets in a 12 hour period
- children under 12 years of age: ask a doctor

## Other Information

- store between 20° to 25°C (68° to 77°F)
- protect from excessive moisture
- this product meets the requirements of USP Dissolution Test 2

## Inactive Ingredients

carnauba wax, FD&C red No. 40 aluminum lake, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, stearic acid, titanium dioxide

## Questions or comments?

Call 1-800-317-2165 (MoN-Fri 9AM-5PM EST)

## PRINCIPAL DISPLAY PANEL - Kit Carton

DUAL FORMULA PACK

**Coricidin**®

### HBP

Decongestant-free **COLD RELIEF** for people

with **HIGH BLOOD PRESSURE**

### DAY

### MULTI-SYMPTOM COLD

Guaifenesin - Expectorant

Dextromethorphan HBr - Cough Suppressant

### Day Relieves:

- Chest Congestion
- Cough

16 DAY SOFTGELS

### NIGHT

Chlorpheniramine Maleate - Antihistamine,  
 Dextromethorphan HBr - Cough Suppressant,  
**Acetaminophen - Pain Reliever/Fever Reducer**  
**See New Warnings Information**

**Night Relieves:**

- Runny Nose & Sneezing
- Cough • Body Aches
- Fever

8 NIGHT TABLETS

		<p>DECONGESTANT-FREE COLD SYMPTOM RELIEF for people with <b>HIGH BLOOD PRESSURE</b></p> <p>No Copy/Color/Varnish LOT /EXPIRY AREA</p> <p><b>SAFETY SEALED:</b> This product is sealed in blister units. If Coricidin® HBP backing has been torn or removed do not use product.</p> 	
<p><b>Coricidin® HBP Day</b></p> <p><b>Drug Facts (continued)</b></p> <p><b>Stop use and ask a doctor</b> if cough lasts more than 7 days, reoccurs, or occurs with fever, rash or persistent headache. These could be signs of a serious condition.</p> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use.</p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>do not exceed recommended dose</li> <li>do not take the Day and Night products at the same time; wait 4 hours after the last Night dose before starting Day product</li> <li>adults and children 12 years and over: 1 or 2 softgels every 4 hours, not more than 6 softgels in 12 hours</li> <li>children under 12 years of age: ask a doctor</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>store between 20° to 25°C (68° to 77°F)</li> <li>protect from freezing</li> <li>protect from excessive moisture</li> </ul> <p><b>Inactive ingredients</b></p> <p>FD&amp;C blue no. 1, FD&amp;C red no. 40, gelatin, glycerin, pharmaceutical ink, polyethylene glycol 400, povidone, propylene glycol, sorbitol</p> <p><b>PARENTS:</b> Learn about teen medicine abuse <a href="http://www.StopMedicineAbuse.org">www.StopMedicineAbuse.org</a></p> <p>Bayer, the Bayer Cross and Coricidin HBP are registered trademarks of Bayer. © 2015 Bayer. Dist. by: Bayer HealthCare LLC Whippany, NJ 07981 For more information, visit <a href="http://www.coricidinhbp.com">www.coricidinhbp.com</a> 38096-02 Bayer</p>	<p><b>Coricidin® HBP Night</b></p> <p><b>Drug Facts (continued)</b></p> <p><b>Ask a doctor or pharmacist before use if you are</b></p> <ul style="list-style-type: none"> <li>taking the blood thinning drug warfarin</li> <li>taking sedatives or tranquilizers</li> </ul> <p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>excitability may occur, especially in children</li> <li>marked drowsiness may occur</li> <li>avoid alcoholic beverages</li> <li>alcohol, sedatives and tranquilizers may increase drowsiness</li> <li>use caution when driving a motor vehicle or operating machinery</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>pain or cough gets worse or lasts more than 7 days</li> <li>fever gets worse or lasts more than 3 days</li> <li>redness or swelling is present</li> <li>new symptoms occur</li> <li>cough comes back or occurs with a rash or headache that lasts. These could be signs of a serious condition.</li> </ul> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use.</p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>do not use more than directed</li> <li>do not take the Day and Night products at the same time; wait 4 hours after the last Day dose before starting Night product</li> <li>adults and children 12 years and over: 2 tablets at bedtime and every 6 hours if needed, while symptoms persist, not more than 4 tablets in a 12 hour period</li> <li>children under 12 years of age: ask a doctor</li> </ul>	<p><b>Coricidin® HBP Day</b></p> <p><b>Drug Facts</b></p> <p><b>Active ingredients Purpose (in each softgel)</b></p> <p>Dextromethorphan hydrobromide 10 mg..... Cough Suppressant    Guaifenesin 200 mg..... Expectorant</p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold</li> <li>helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive</li> </ul> <p><b>Warnings</b></p> <p><b>Do not use</b> if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>cough that occurs with excessive phlegm (mucus)</li> <li>persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema</li> </ul>	
 <p>03809602</p>		<p><b>Coricidin® HBP Night</b></p> <p><b>Drug Facts (continued)</b></p> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>store between 20° to 25°C (68° to 77°F)</li> <li>protect from excessive moisture</li> <li>this product meets the requirements of USP Dissolution Test 2</li> </ul> <p><b>Inactive ingredients</b></p> <p>cambeba wax, FD&amp;C red No. 40 aluminum lake, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, stearic acid, titanium dioxide</p> <p><b>Questions or comments?</b> 1-800-317-2165 (Mon-Fri 9AM - 5PM EST)</p>	

**CORICIDIN HBP DAY AND NIGHT**

dextromethorphan hydrobromide, guaifenesin, acetaminophen, chlorpheniramine maleate kit

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11523-4765
---------------------	----------------	---------------------------	----------------

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11523-4765-1	1 in 1 CARTON; Type 0: Not a Combination Product	08/28/2009	

## Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	2 BLISTER PACK	16
Part 2	2 BLISTER PACK	8

## Part 1 of 2

### CORICIDIN HBP DAY

dextromethorphan hydrobromide and guaifenesin capsule, gelatin coated

## Product Information

Route of Administration ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg

## Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN (UNII: 2G86QN327L)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SORBITOL (UNII: 506T60A25R)	

## Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	C;DAY
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11523-4221-1	2 in 1 CARTON		
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/28/2009	

## Part 2 of 2

### CORICIDIN HBP NIGHT

acetaminophen, chlorpheniramine maleate and dextromethorphan hydrobromide tablet

## Product Information

Route of Administration	ORAL
-------------------------	------

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>ALUMINUM OXIDE</b> (UNII: LMI26O6933)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>LACTOSE</b> (UNII: J2B2A4N98G)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	18mm
<b>Flavor</b>		<b>Imprint Code</b>	C;NIGHT
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11523-4111-1	2 in 1 CARTON		
1		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/28/2009	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/28/2009	

**Labeler** - Bayer HealthCare LLC. (112117283)

Revised: 12/2025

Bayer HealthCare LLC.