

COLD AND FLU NIGHTTIME/MULTI-SYMPTOM- acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

Walgreen Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Walgreens 44-660

Active ingredients (in each liquid-filled capsule)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

- temporarily relieves common cold and flu symptoms:
 - sore throat
 - minor aches and pains
 - fever
 - headache
 - runny nose and sneezing
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients
- to make a child sleepy
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

- liver disease
- glaucoma
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic beverages
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- do not take more than 8 capsules per 24 hours
- adults and children 12 years and over: take 2 capsules with water every 6 hours
- children under 12 years : ask a doctor

Other information

- protect from heat, humidity and light
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS**

TORN OR BROKEN

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

D&C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol

Questions or comments?

1-800-426-9391

Principal Display Panel

Walgreens

**Compare to Vicks® NyQuil® Cold & Flu
LiquiCaps® active ingredients††**

NDC 0363-6600-08

NIGHTTIME

Cold & Flu

**ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr - COUGH SUPPRESSANT
DOXYLAMINE SUCCINATE/ ANTIHISTAMINE**

MULTI-SYMPTOM

- Relieves aches, fever, sore throat, sneezing, runny nose & cough

24

LIQUID CAPS ACTUAL SIZE

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS
TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

Walgreens Pharmacist Recommended

Walgreens Pharmacist Survey

††This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® NyQuil® Cold & Flu LiquiCaps®.

50844 REV0418B66008

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015

Walgreens
100% SATISFACTION
GUARANTEED

walgreens.com ©2017 Walgreen Co.
PRODUCT OF CHINA. PACKAGED AND
QUALITY ASSURED IN THE U.S.A.

ORF00817-F
REV0719

Drug Facts (continued)

Directions ■ do not take more than directed
 ■ do not take more than 8 capsules per 24 hours
 ■ adults and children 12 years and over: take 2 capsules with water every 6 hours
 ■ children under 12 years: ask a doctor

Other information ■ protect from heat, humidity and light
 ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

Inactive ingredients D&C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol

(Questions or comments? 1-800-426-9391)

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

NO PRINT/NO VARNISH LOT & EXP DATE

Drug Facts (continued)

Warnings ■ This product contains acetaminophen. Severe liver damage may occur if you take
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
 ■ Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include ■ skin redness ■ blisters ■ rash
 ■ If a skin reaction occurs, stop use and seek medical help right away.
 ■ Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
 ■ Do not use
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients
 ■ to make a child sleepy

Uses ■ temporarily relieves common cold and flu symptoms;
 ■ sore throat ■ minor aches and pains
 ■ headache ■ fever ■ runny nose and sneezing
 ■ cough due to minor throat and bronchial irritation

Active ingredients (in each liquid-filled capsule)
 Acetaminophen 325 mg Pain reliever/fever reducer
 Dextromethorphan HBr 15 mg Cough suppressant
 Doxylamine succinate 6.25 mg Antihistamine

Purpose

Drug Facts (continued)

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ Ask a doctor before use if you have
 ■ liver disease ■ glaucoma
 ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
 ■ cough that occurs with too much phlegm (mucus)
 ■ difficulty in urination due to enlargement of the prostate gland
 ■ Ask a doctor or pharmacist before use if you are
 ■ taking the blood thinning drug warfarin
 ■ taking sedatives or tranquilizers
 ■ When using this product, ■ do not exceed recommended dosage
 ■ avoid alcoholic beverages
 ■ marked drowsiness may occur
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness
 ■ Stop use and ask a doctor if
 ■ pain or cough gets worse or lasts more than 7 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ new symptoms occur
 ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
 ■ If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

ITEM 384888 W000000-0000-0
 1917117433
 4

DISTRIBUTED BY: WALGREEN CO.
 200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED

walgreens.com ©2017 Walgreen Co.
 PRODUCT OF CHINA, PACKAGED AND QUALITY ASSURED IN THE U.S.A.

Walgreens Pharmacist Recommended
 Walgreens Pharmacy Survey
 If This Product Is Not Manufactured or Distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® NyQuil® Cold & Flu LiquiCaps®,
 50844 REV0418B66008

NIGHTTIME
Cold & Flu



Walgreens

NIGHTTIME
Cold & Flu

Compare to Vicks® NyQuil® Cold & Flu LiquiCaps® active ingredients!¹

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
 DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
 DOXYLAMINE SUCCINATE / ANTIHISTAMINE

MULTI-SYMPOM

- Relieves aches, fever, sore throat, sneezing, runny nose & cough

24
 LIQUID CAPS



ACTUAL SIZE

COLD AND FLU NIGHTTIME/MULTI-SYMP TOM

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-6600
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	GREEN (clear)	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	660
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-6600-08	2 in 1 CARTON	03/01/2015	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0363-6600-21	2 in 1 CARTON	03/01/2015	
2		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	03/01/2015	

Labeler - Walgreen Company (008965063)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	PACK(0363-6600)

Revised: 8/2019

Walgreen Company