

**MAGNESIUM CITRATE- magnesium citrate oral solution - lemon liquid**  
**Shield Pharmaceuticals Corp**

---

**Magnesium Citrate Oral Solution - Lemon**

**Active Ingredient**

Magnesium citrate 1.745g per fl oz

**Purpose**

Saline Laxative

**Use**

- for relief of occasional constipation (irregularity).
- Generally produces bowel movement in 1/2 to 6 hours

**Warnings**

**Ask a doctor before use if you have**

- kidney disease
- a magnesium restricted diet
- abdominal pain, nausea, or vomiting
- noticed a sudden change in bowel habits that persists over a period of 2 weeks
- already used a laxative for a period longer than 1 week

**Ask a doctor or pharmacist before use if you are**

- taking any other drug.
- take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

**Stop use and ask a doctor if you have rectal**

bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222).

**Directions**

- shake well before using \_ drink a full glass (8 ounces) of liquid with each dose
- may be taken as a single daily dose or in divided doses

---

adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours

children 6 to under 3 to 12 years of age	7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age	2 to 3 fl oz in 24 hours maximum 3 fl oz in 24 hours
children under 2 years of age	ask a doctor

## Other Information

- Store at temperatures between 46° and 86°F (8° and 30° C)
- each fl oz contains: magnesium 290 mg
- each fl oz contains: sodium 1 mg

## Inactive Ingredients

benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

## Questions?

1-800-373-6981

**Drug facts (continued)**

**Directions** ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over	6.5 to 10 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age	3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age	2 to 3 fl oz in 24 hours maximum 3 fl oz in 24 hours
children under 2 years of age	ask a doctor

**Other information** ■ Store at temperatures between 46° and 86°F (8° and 30° C)  
■ each fl oz contains: magnesium 290 mg  
■ each fl oz contains: sodium 1 mg

**Inactive ingredients** benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

**Questions or comments?** 1-800-373-6981

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS BROKEN OR MISSING

Distributed by:  
Your Company name  
Address



0 00000 00000 0

**ValuRx**

# Magnesium Citrate

**Saline Laxative / Oral Solution**

**Sugar Free** **Dye Free**

**10 fl oz (296 mL)** **Lemon Flavor**



**Drug facts**

**Active ingredients (in fl. oz)** Purpose  
Magnesium citrate 1.745g per fl oz .....Saline Laxative

**Uses**  
■ for relief of occasional constipation (irregularity).  
■ Generally produces bowel movement in 12 to 6 hours

**Warnings**  
**Ask a doctor before use if you have**  
■ kidney disease  
■ a magnesium restricted diet  
■ abdominal pain, nausea, or vomiting  
■ noticed a sudden change in bowel habits that persists over a period of 2 weeks  
■ already used a laxative for a period longer than 1 week

**Ask a doctor or pharmacist before use if you are**  
■ taking any other drug.  
■ take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

**Stop use and ask a doctor if** you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222). ►

## MAGNESIUM CITRATE

magnesium citrate oral solution - lemon liquid

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83059-0127
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>MAGNESIUM CITRATE</b> (UNII: RHO26O1T9V) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM CITRATE	1.745 g in 296 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>BENZOIC ACID</b> (UNII: 8SKNOB0MIM)	
<b>CITRIC ACID</b> (UNII: 2968PHW8QP)	
<b>DISODIUM EDTA-COPPER</b> (UNII: 6V475AX06U)	1.745 g in 296 mL
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>WATER</b> (UNII: 059QF0KOOR)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83059-0127-1	296 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/26/2026	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	01/26/2026	

**Labeler** - Shield Pharmaceuticals Corp (118724924)

Revised: 1/2026

Shield Pharmaceuticals Corp