#### MECLIZINE HYDROCHLORIDE- meclizine hcl tablet NORTHEAST PHARMA

-----

## hpc 778L (378)

#### Active ingredient (in each tablet)

Meclizine HCI 25mg

#### Purpose

Antiemetic

#### Uses

For the prevention and treatment of these symptoms associated with motion sickness • nausea • vomiting • dizziness

#### Warnings

**Do not use** in children under 12 years of age unless directed by a doctor.

### Ask a doctor before use if you have

- breathing problems, such as emphysema or chronic bronchitis
- glaucoma
- difficulty urinating due to an enlarged prostate gland

# A sk a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

#### When using this product

- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery
- I f pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

- take the first dose 1/2 hour to 1 hour before starting activity
- Adults and children 12 years of age and over:

Take 1 to 2 tablets once daily, or as directed by a doctor Do not exceed 2 tablets in 24 hours.

## Other information

- **TAMPER EVIDENT:**Do not use if imprinted seal under cap is broken or missing.
- store at 20-25°C (68-77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from humidity

## **Inactive ingredients**

colloidal silicon dioxide, D&C yellow #10 lake, lactose monohydrate, magnesium stearate, pregelatinized starch.

## **Questions or comments?**

1-800-540-3765

## package label

NDC 80136-378-01	Purpose Antiemetic	<ul> <li>dizziness</li> </ul>	unless	glaucoma rchronic bronchitis I prostate gland if you are taking	Icoholic beverages y increase drowsiness or operating machinery t health professional	In case of overdose, get Control Center right away.	r before starting activity age and over: directed by a doctor hours.	ER EVIDENT: Do not s broken or missing. cursions permitted protect from humidity	con dioxide, ydrate, starch.	distributed by the DRAMAMINE®.	REV 1023L
Meclizine Hydrochloride Antiemetic • Antinausea	each tablet)	Uses For the prevention and treatment of these symptoms with motion sickness • nausea • vomiting • di	12 years of age unless	a or ged	When using this product • drowslness may occur • avoid alcoholic beverages • alcohol, sedatives and tranquilizers may increase drowsiness • alcohol, sedatives and tranquilizers may increase drowsiness • use caution when drwing a motor vehicle or operating machinery • use caution when drwing a motor vehicle or operating machinery for pregnant or breast-feeding, ask a health professional proferments	n. In case of o	s of un 24	on • TAMPER EVIDENT: Do not al under cap is broken or missing. (68-77°F); excursions permitted (59°-86°F) • protect from humid	dal silic monoh inized s		
FOR PREVENTION AND TREATMENT OF MOTION SICKNESS for up to 24 hours Ages 12 and Over		and treatment ress • nause	children under 1 octor.	Ask a doctor before use if you have • breathing problems, such as emphysem • difficulty urinating due to an enlar Ask a doctor or pharmacist before u sachtives or tranouilizers	The number of the second secon	reach of children. or contact a Poison	dose 1/2 hour to 1 children 12 year blets once daily, c exceed 2 tablets	information • T/ mprinted seal under at 20-25°C (68-77° in 15°-30°C (59°-86	<i>ingredients</i> w #10 lake, k m stearate, p	uct is not manufacture. Inthe registered tradema I By: Northeast Pharma indom VT 05403 1.54	
Compare to active ingredient in DRAMAMINE® LESS DROWSY*	Drug Facts Active ingredient (in Medizine HCl 25ma	s the prevention motion sickr	<i>Warnings</i> Do not use for childr directed by a doctor.	Ask a doctor before use • breathing problems, such • difficulty urinating due Ask a doctor or pharma	When using this p when using this p a drowsiness may a alcohol, sedatives • use caution when d ff pregnant or bre	ut of re d help or	ctions e the first ults and 1 to 2 tal Do not	Other informati use if imprinted set e store at 20-25°C between 15°-30°C	Inactive in D&C yellow magnesium	*This product is r owner of the rec Distributed By: N Couth Burlington	8 www.heart
100 Tablets • 25 mg Each	Drug I Active Meclizir	Uses For the with n	Do n dire	Ask • • bras	When • drow • alcoh • use co If preg	Kee	Dire • tak • Ad				

## **MECLIZINE HYDROCHLORIDE**

meclizine hcl tablet								
Product Information								
Product Type	HUMAN OTC DRUG	Item Code (So	ource)	NDC:80136-378				
Route of Administration	ORAL							
Active Ingredient/Active Moiety								
Ingre	<b>Basis of Strength</b>		Strength					
MECLIZINE HYDROCHLORIDE (UN	MECLIZ INE		25					

Inactive Ingredients								
	Strength							
SILICON DIOXIDE (UNII: ETJ7								
STARCH, CORN (UNII: 08232								
MAGNESIUM STEARATE (UN								
D&C YELLOW NO. 10 ALUM	D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)							
Product Characteristics								
Color	yellow	Score	no score					
Shape	ROUND	Size	9mm					
Flavor		Imprint Code	44;403					
Contains	ontains							
Packaging								
# Item Code	Package Descri	iption	Marketing Start Date	Marketing End Date				
<b>1</b> NDC:80136-378- 01100 in 1 BOTTLE; Type 0: Not a Combination Product05/01/2022								
Marketing Information								
Marketing App Category	lication Number Citation		Marketing Start Date	Marketing End Date				
OTC Monograph Drug M009			05/01/2022					

Labeler - NORTHEAST PHARMA (081232935)

Registrant - Geri-Care Pharmaceutical Corp (611196254)

Revised: 11/2023

NORTHEAST PHARMA