

POVIDONE IODINE- povidone iodine 10% liquid
Pharma Nobis, LLC

Private Label Povidone Iodine

Drug Facts

Active Ingredient

Povidone-Iodine 10%
(1% available iodine)

Purpose

First Aid Antiseptic

Use

First aid antiseptic to prevent infection in minor cuts and burns.

Warnings

For External Use Only

Ask a doctor before use if you have

- deep puncture wounds
- animal bites
- serious burns

Stop use and consult a doctor if

- the condition persists or gets worse
- irritation and redness develop and persists for more than 72 hours

When using this product do not

- use in eyes
- use on individuals who are allergic or sensitive to iodine or use for longer than 1 week unless directed by a doctor
- apply over large areas of the body

Keep out of reach of children.

In case of accidental ingestion, seek professional assistance or contact a Poison Control

Center immediately.

Directions

Clean the affected area. Apply a small amount on the area 1 to 3 times daily. May be covered with sterile bandage. If badaged, let it dry first.

Other information

store at room temperature, 20-25C (68-77F)

Inactive Ingredients

Citric Acid, Disodium Phosphate, Glycerin, nonoxynol-9. Sodium Hydroxide and Purified Water.

Sunmark Label

8 fl oz

The image shows a rectangular label for Sunmark Povidone-Iodine 10% Topical Solution. The label has a blue background with a large red circular graphic on the left side. The text is primarily white and yellow. At the top left is the Sunmark logo. To its right, it says 'COMPARE TO BETADINE® ACTIVE INGREDIENT* NDC 49348-622-37'. The main product name 'povidone-iodine' is written in large white letters, with '10% Topical Solution' below it in yellow. At the bottom left, it says 'FIRST AID ANTISEPTIC' and '8 FL OZ (237 mL)'. On the right side, there is a white box containing 'Drug Facts' information, including Active Ingredient (Povidone Iodine 10%), Purpose (Antiseptic), Uses (First aid antiseptic), Warnings (For external use only), Do not (use in eyes, longer than 1 week, on allergic individuals, over large areas), Stop use and consult a doctor if (condition persists, irritation/redness), Ask a doctor in case of (deep/puncture wounds, animal bites, serious burns), Keep out of reach of children, Directions (Clean affected area, apply 1-3 times daily), Other Information (store at room temperature), and Inactive Ingredients (Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide, Purified Water). A barcode is on the right edge with the number 0 10939 14233 7. At the bottom right, it says 'MEKESON Empowering Healthcare Distributed By McKesson One Post Street, San Francisco, CA 94104 Money Back Guarantee'. A small disclaimer at the very bottom reads '*This product is not manufactured or distributed by Purdue Frederick, owner of the registered trademark Betadine®'.

Drug Facts	
Active Ingredient Povidone Iodine 10%.....Antiseptic (*Equivalent to 1% titratable iodine)	Purpose
Uses First aid antiseptic to prevent infection in minor cuts, burns.	
Warnings For external use only. Do not: ■ use in the eyes. ■ use longer than 1 week unless directed by a doctor. ■ use on individuals who are allergic or sensitive to iodine. ■ apply over large areas of the body.	
Stop use and consult a doctor if ■ condition persists or gets worse. ■ irritation and redness develop and persist for more than 72 hours.	
Ask a doctor in case of: ■ deep or puncture wounds ■ animal bites ■ serious burns.	
Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.	
Directions ■ Clean the affected area. ■ Apply a small amount on the area 1 to 3 times daily. ■ May be covered with a sterile bandage. ■ If bandaged, let dry first.	
Other Information store at room temperature, 20-25° C (68-77° F).	
Inactive Ingredients: Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water.	

16 fl oz

sunmark™

COMPARE TO BETADINE®
ACTIVE INGREDIENT*
NDC 49348-622-38

povidone-iodine

10% Topical Solution

FIRST AID ANTISEPTIC

16 FL OZ (1 PT) 473 mL

Drug Facts

Active Ingredient Povidone Iodine 10%
Purpose Antiseptic
(*Equivalent to 1% titratable iodine)

Uses
First aid antiseptic to prevent infection in minor cuts, burns.

Warnings For external use only.

Do not:
■ use in the eyes.
■ use longer than 1 week unless directed by a doctor.
■ use on individuals who are allergic or sensitive to iodine.
■ apply over large areas of the body.

Stop use and consult a doctor if
■ condition persists or gets worse.
■ irritation and redness develop and persist for more than 72 hours.

Ask a doctor in case of:
■ deep or puncture wounds ■ animal bites ■ serious burns.

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Directions
■ Clean the affected area. ■ Apply a small amount on the area 1 to 3 times daily. ■ May be covered with a sterile bandage. ■ If bandaged, let dry first.

Other Information store at room temperature, 20-25° C (68-77° F).

Inactive Ingredients: Citric Acid, Disodium Phosphate, Glycerin, Nonoxonyl-9, Sodium Hydroxide and Purified Water.

*This product is not manufactured or distributed by Purdue Frederick, owner of the registered trademark Betadine®



MEKESON
Distributed By McKesson
One Post Street, San Francisco, CA 94104
Money Back Guarantee

Good Neighbor Label

NDC 24385-053-55



Compare to
active ingredient
in Betadine®

Povidone-Iodine

10% Topical Solution

Topical Antiseptic

Helps prevent infection in minor cuts, scrapes, and burns

- Use for first aid & general hygiene
- Helps reduce bacteria that can potentially cause skin infection
- Water soluble

8 FL OZ (236 mL)

Drug Facts	Drug Facts (continued)
Active ingredient Povidone-Iodine USP, 10% w/v (equivalent to 1% titratable iodine) Purpose Antiseptic	Directions ■ Clean the affected area ■ Apply a small amount of this product to the area 1 – 3 times daily ■ May be covered with a sterile bandage ■ If bandaged, let dry first
Use First aid antiseptic to help prevent the risk of infection in minor cuts, scrapes and burns	Other information Store at room temperature: 15° – 30° C (59° – 86° F)
Warnings For external use only. Do not ■ use in the eyes ■ use on individuals who are allergic or sensitive to iodine ■ apply over large areas of the body	Inactive ingredients citric acid, disodium phosphate, nonoxonyl-9, glycerin, purified water, sodium hydroxide.
Discontinue use and ask a doctor ■ if condition persists or gets worse ■ for use longer than 1 week	*This product is not manufactured or distributed by Purdue Frederick, owner of the registered trademark, Betadine®.
Ask a doctor in case of ■ deep or puncture wounds ■ animal bites ■ serious burns	A46492 04 2022
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	



ABC# 127-092

R102513RLG

Distributed by: AmerisourceBergen
1300 Morris Drive
Chesterbrook, PA 19087
Made in U.S.A.

Questions or Comments?
1-800-662-3435

www.goodneighborpharmacy.com

NDC 24385-053-55



povidone iodine, 10% solution

*Compare to active ingredient in Betadine®

Topical antiseptic
Helps prevent infection in minor cuts, scrapes, and burns

- Use for first aid & general hygiene
- Helps reduce bacteria that can potentially cause skin infection
- Water soluble

8
fl. oz.
(236 mL)

LOT

EXP

Drug Facts

Active ingredient	Purpose
Povidone Iodine USP, 10% w/v (equivalent to 1% titratable iodine)	Antiseptic

Use First aid antiseptic to help prevent the risk of infection in minor cuts, scrapes and burns

Warnings For external use only.

Do not

- use in the eyes
- use on individuals who are allergic or sensitive to iodine
- apply over large areas of the body

Discontinue use and ask a doctor

- if condition persists or gets worse
- for use longer than 1 week

Ask a doctor in case of

- deep or puncture wounds
- animal bites
- serious burns

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- Clean the affected area
- Apply a small amount of this product to the area 1 – 3 times daily
- May be covered with a sterile bandage
- If bandaged, let dry first

Other information

Store at room temperature: 15° – 30° C (59° – 86° F)

Inactive ingredients

citric acid, glycerin, Makon 8, purified water, sodium hydroxide

*This product is not manufactured or distributed by Purdue Frederick, owner of the registered trademark, Betadine®.

Rev. A




Distributed By
AmerisourceBergen
Valley Forge, PA 19482-0959

Visit us at
www.myGNP.com



Leader Label

Drug Facts	
Active ingredient	Purpose
Povidone-iodine 10%.....First Aid Antiseptic (1% available iodine)	
Uses	
First aid antiseptic to help prevent infection in minor cuts, scrapes and burns	
Warnings For external use only.	
Ask a doctor before use if you have	
<ul style="list-style-type: none"> ■ deep or puncture wounds ■ animal bites ■ serious burns 	
Stop use and consult a doctor if	
<ul style="list-style-type: none"> ■ the condition persists or gets worse ■ you need to use this product for more than 1 week 	
When using this product do not	
<ul style="list-style-type: none"> ■ use in the eyes ■ use if you are allergic to povidone-iodine or any other ingredients in this preparation ■ apply over large areas of the body 	
Keep out of reach of children. If swallowed, seek medical help or contact a Poison Control Center (1-800-222-1222) immediately.	
Continued ▶	



NDC 70900-0402-1

Povidone-Iodine

10% Topical Solution

First Aid Antiseptic

Kills Germs Promptly in Minor Cuts, Scrapes and Burns

COMPARE TO BETADINE®
active ingredient**

100% Money Back Guarantee

8 FL OZ (237 mL)

† SAFETY SEALED

***Warning:** Do not use if Tamper Evident Seal is broken or missing. This product is sealed with a foil seal under cap.

Drug Facts (continued)
Directions
<ul style="list-style-type: none"> ■ clean the affected area ■ apply a small amount on the area 1 to 3 times daily ■ may be covered with sterile bandage ■ if bandaged let dry first
Other information
<ul style="list-style-type: none"> ■ store at room temperature, 20-25° C (68-77° F)
Inactive ingredients
Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water
Questions or comments? 1-800-662-3435

**This product is not manufactured or distributed by Purdue Products L.P., owner of the registered trademark Betadine®.

CIN 5455472
REV. 1/19



© 2019 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.

DISTRIBUTED BY CARDINAL HEALTH
DUBLIN, OHIO 43017
www.myleader.com
1-800-200-6313


100% Money Back Guarantee

Return to place of purchase.

A47502
05 2022

NDC 37205-186-34

*SAFETY SEALED



Compare to Betadine® active ingredient**

Povidone-Iodine Solution

10% Topical Solution

Antiseptic / Germicide

Promptly Kills Germs
In Cuts, Burns, Scrapes
and Minor Skin Wounds

*WARNING: Do not use if Tamper Evident Seal is broken or missing This product is sealed with a foil seal under cap

8 FL OZ (237mL)

Drug Facts	
Active ingredient	Purpose
Povidone-iodine, 10% (Equivalent to 1% titratable iodine).....	Antiseptic
Uses	
<ul style="list-style-type: none"> ■ Health-care antiseptic for preparation of the skin prior to surgery ■ First aid antiseptic to help prevent infection in minor cuts, scrapes and burns. 	
Warnings	
For external use only.	
Do not use	
<ul style="list-style-type: none"> ■ in the eyes ■ as a first aid antiseptic longer than one week ■ on individuals who are allergic or sensitive to iodine ■ over large areas of the body 	
Stop use and ask a doctor	
<ul style="list-style-type: none"> ■ if condition persists or gets worse ■ if irritation and redness develop and persist for over 72 hours. 	
Ask a doctor in case of	
<ul style="list-style-type: none"> ■ deep puncture wounds ■ animal bites ■ serious burns 	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions:	
For preparation of the skin prior to surgery:	
<ul style="list-style-type: none"> ■ apply to the operative site prior to surgery. 	
As a first aid antiseptic:	
<ul style="list-style-type: none"> ■ clean the affected area ■ apply a small amount of this product to the area 1 – 3 times daily ■ may be covered with a sterile bandage ■ if bandaged, let dry first. 	
Other information Store at room temperature: 15° – 30° C (59° – 86° F)	
Inactive ingredients citric acid, disodium phosphate, glycerin, nonoxynol-9, sodium hydroxide, and purified water.	

* This product is not manufactured or distributed by Purdue Frederick, owner of the registered trademark BETADINE®.

R052311RLG

DISTRIBUTED BY:
CARDINAL HEALTH DUBLIN, OHIO 43017
CIN 2443554 www.myleader.com 1-800-200-6313




0 96295 10673 2

All Leader® Brand Products are 100% satisfaction guaranteed or return to place of purchase for a full refund.

Quality Choice Label

Drug Facts	
Active ingredient	Purpose
Povidone-iodine 10%.....First Aid Antiseptic (*Equivalent to 1% titratable iodine)	
Indications	
First aid antiseptic to prevent infection in minor cuts, scrapes and burns	
Warnings For external use only.	
Ask a doctor before use if you have	
<ul style="list-style-type: none"> ■ deep or puncture wounds ■ animal bites ■ serious burns 	
Stop use and consult a doctor if	
<ul style="list-style-type: none"> ■ the condition persists or gets worse ■ irritation and redness develop and persists for more than 72 hours 	
When using this product do not	
<ul style="list-style-type: none"> ■ use in the eyes ■ use on individuals who are allergic or sensitive to iodine or use for longer than 1 week unless directed by a doctor ■ apply over large areas of the body 	
Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.	
<i>Continued ▶</i>	

NDC 63868-230-08



***Compare to active ingredient in Betadine®**



Povidone-Iodine

Topical Antiseptic Germicide

10% Topical Solution

Helps Prevent Infection In Cuts, Burns, Scrapes and Minor Skin Wounds

For external use only

8 fl OZ (237mL)

Drug Facts (continued)

Directions

- clean the affected area ■ apply a small amount on the area 1 to 3 times daily ■ may be covered with sterile bandage ■ if bandaged let dry first

Other information

- store at room temperature, 20-25° C (68-77° F)

Inactive ingredients
Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water

**This product is not manufactured or distributed by Purdue Products L.P., owner of the registered trademark Betadine®.

A46494
04 2022




L-QC-PW-8 R02/14

6 35515 99230 6

Distributed by C.D.M.A., Inc.®
43157 W. Nine Mile
Novi, MI 48376-0995
www.qualitychoice.com
Questions: 248-449-9300

MADE IN USA



*Compare to the active ingredients of BETADINE®

NDC# 63868-230-08

Povidone Iodine

10% Topical Solution

TOPICAL ANTISEPTIC GERMICIDE


Helps Prevent Infections in:

- Burns • Cuts
- Minor Skin Wounds
- Scrapes

FOR EXTERNAL USE ONLY


8 fl oz (237 mL)

Drug Facts	
Active Ingredient	Purpose
Povidone-iodine, 10% (equivalent 1% titratable iodine)	Antimicrobial
Uses	
<ul style="list-style-type: none"> • health-care antiseptic for preparation of the skin prior to surgery • first aid antiseptic to help prevent infection in minor cuts, scrapes and burns 	
Warnings For external use only	
Do not use	
<ul style="list-style-type: none"> • in the eyes or apply over large areas of the body • as a first aid antiseptic longer than one week unless directed by a doctor • on individuals who are allergic or sensitive to iodine 	
Stop use and ask a doctor	
<ul style="list-style-type: none"> • if condition persists or gets worse • if irritation and redness develop and persist for over 72 hours 	
Ask a doctor in case of	
<ul style="list-style-type: none"> • deep or puncture wounds • animal bites • serious burns 	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions	
For preparation of the skin prior to surgery	
<ul style="list-style-type: none"> • apply to the operative site prior to surgery 	
As a first aid antiseptic	
<ul style="list-style-type: none"> • clean the affected area • apply a small amount of this product to the area 1-3 times daily • may be covered with a sterile bandage • if bandaged, let dry first 	
Inactive ingredients	
citric acid, disodium phosphate, glycerin, nonoxynol-9, sodium hydroxide, purified water USP.	



6 35515 99230 6

© DISTRIBUTED BY QUALITY CHOICE
NOVI, MI 48376-0995
(800) 935-2362
MADE IN U.S.A.
QC 98515
www.chalndrug.com



L-QC-PW-8 R02/07

Select Brand Label

NDC 15127-052-55

select brand
the lower price name brand

Povidone Iodine 10% Topical Solution

TOPICAL ANTISEPTIC GERMICIDE

Helps Prevent Infections in:

- Burns • Cuts
- Minor Skin Wounds
- Scrapes

FOR EXTERNAL USE ONLY



Compare to the active ingredients of BETADINE®*

8 fl oz (237 mL)

Drug Facts

Active Ingredient	Purpose
Povidone-Iodine, 10% (equivalent 1% titratable iodine)	Antimicrobial

Uses

- health-care antiseptic for preparation of the skin prior to surgery
- first aid antiseptic to help prevent infection in minor cuts, scrapes and burns

Warnings For external use only

Do not use

- in the eyes or apply over large areas of the body
- as a first aid antiseptic longer than one week unless directed by a doctor
- on individuals who are allergic or sensitive to iodine

Stop use and ask a doctor

- if condition persists or gets worse
- if irritation and redness develop and persist for over 72 hours

Ask a doctor in case of

- deep or puncture wounds • animal bites • serious burns

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

For preparation of the skin prior to surgery

- apply to the operative site prior to surgery

As a first aid antiseptic

- clean the affected area
- apply a small amount of this product to the area 1-3 times daily
- may be covered with a sterile bandage
- if bandaged, let dry first

Inactive ingredients citric acid, disodium phosphate, glycerin, nonoxynol-9, sodium hydroxide, purified water

Distributed by:
SELECT BRAND® DISTRIBUTORS
Pine Bluff, AR 71603 USA
AC (870) 535-3635
L-SB-PW-8 R08/06



Publix Label

Publix

povidone-iodine

10% TOPICAL SOLUTION

first aid antiseptic

Compare to Betadine®*

8 FL OZ (236mL)



*This Product is not manufactured or distributed by Purdue Frederick Company, owner of the registered trademark Betadine®

Drug Facts

Active Ingredient	Purpose
Povidone Iodine 10% (*Equivalent to 1% titratable iodine)	Antiseptic

Uses
First aid antiseptic to prevent infection in minor cuts, burns.

Warnings For external use only.

Do not:

- use in the eyes.
- use longer than 1 week unless directed by a doctor.
- use on individuals who are allergic or sensitive to iodine.
- apply over large areas of the body.

Stop use and consult a doctor if

- condition persists or gets worse.
- irritation and redness develop and persist for more than 72 hours.

Ask a doctor in case of:

- deep or puncture wounds ■ animal bites ■ serious burns.

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Directions

- Clean the affected area. ■ Apply a small amount on the area 1 to 3 times daily. ■ May be covered with a sterile bandage. ■ If bandaged, let dry first.

Other Information store at room temperature, 20-25° C (68-77° F).

Inactive Ingredients: Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water.

DISTRIBUTED BY PUBLIX SUPER MARKETS, INC. • 3300 PUBLIX CORPORATE PARKWAY, LAKELAND, FL 33811
800-242-1227 • publix.com • PUBLIX GUARANTEE: COMPLETE SATISFACTION OR YOUR MONEY BACK

Premier Value Label

Drug Facts

Active ingredient Purpose
Povidone-iodine 10%.....Antiseptic
(*Equivalent to 1% titratable iodine)

Use

First aid antiseptic to prevent infection in minor cuts and burns

Warnings For external use only.

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and consult a doctor if

- the condition persists or gets worse
- irritation and redness develop and persists for more than 72 hours

When using this product do not

- do not use in the eyes
- use on individuals who are allergic or sensitive to iodine or use for longer than 1 week unless directed by a doctor
- do not apply over large areas of the body

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Continued ►



NDC 68016-415-00
**Compare to Active Ingredient in Betadine®

Povidone-Iodine

10%* TOPICAL SOLUTION USP

Topical Antiseptic Microbicide

NET WT. 8 FL OZ (237 mL)



Drug Facts (continued)

Directions

- clean the affected area
- apply a small amount on the area 1 to 3 times daily
- may be covered with sterile bandage
- if bandaged let dry first

Other information

- store at room temperature, 20-25° C (68-77° F)

Inactive ingredients

Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water

**This product is not manufactured or distributed by Purdue Products L.P., owner of the registered trademark Betadine®.

R110113.RLG



8 40986 01294 5

If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

DISTRIBUTED BY: CHAIN DRUG CONSORTIUM
3301 NW BOCA RATON BLVD
SUITE 101, BOCA RATON, FL 33431
MADE IN U.S.A.

Fred's Label

FRED'S

First Aid Antiseptic Solution

10% Povidone-Iodine Topical Solution

Kills Germs And Helps Prevent The Risk Of Infection In:

- Minor Cuts
- Scrapes
- Burns



Compare to Active Ingredients In:

BETADINE®*
Povidone-Iodine

8 FL OZ (236 mL)

Area left blank for lot and expiration date.

Drug Facts

Active ingredient Purpose
Povidone- Iodine USP,
10% w/v (equivalent to
1% titratable iodine)Antiseptic

Use First aid antiseptic to help prevent the risk of infection in minor cuts, scrapes and burns

Warnings For external use only.

- Do not**
- use in the eyes
 - use on individuals who are allergic or sensitive to iodine
 - apply over large areas of the body

Discontinue use and ask a doctor

- if condition persists or gets worse
- for use longer than 1 week

Ask a doctor in case of

- deep or puncture wounds
- animal bites
- serious burns

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. ►

Drug Facts (continued)

Directions

- Clean the affected area
- Apply a small amount of this product to the area 1 – 3 times daily
- May be covered with a sterile bandage
- If bandaged, let dry first

Other information

Store at room temperature:
15° – 30° C (59° – 86° F)

Inactive ingredients

citric acid, disodium phosphate, glycerin, nonoxynol-9, sodium hydroxide, purified water.

*This product is not manufactured or distributed by Purdue Frederick, owner of the registered trademark, Betadine®.

R062511RLG



0 84579 10500 7

FRED'S Promise

Not Happy With This Product?
Just Bring It Back For a Refund!

Distributed By: Fred's, Inc.
4300 New Getwell Rd, Memphis, TN 38118
www.fredsinc.com

DDM Label



POVIDONE- IODINE

10% Topical Solution

**Topical Antiseptic
Germicide**
Helps Prevent Infections in:
• Burns • Cuts • Scrapes
For External Use Only

8 FL OZ (237mL)

Drug Facts

Active Ingredient	Purpose
Povidone-iodine, 10% (equivalent 1% titratable iodine)	Antimicrobial

Uses • health-care antiseptic for preparation of the skin prior to surgery
• first aid antiseptic to help prevent infection in minor cuts, scrapes and burns

Warnings For external use only

Do not use

- in the eyes or apply over large areas of the body
- as a first aid antiseptic longer than one week unless directed by a doctor
- on individuals who are allergic or sensitive to iodine

Stop use and ask a doctor

- if condition persists or gets worse
- if irritation and redness develop and persist for over 72 hours

Ask a doctor in case of

- deep or puncture wounds • animal bites • serious burns

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

For preparation of the skin prior to surgery

- apply to the operative site prior to surgery

As a first aid antiseptic

- clean the affected area
- apply a small amount of this product to the area 1-3 times daily
- may be covered with a sterile bandage
- if bandaged, let dry first

Inactive ingredients: citric acid, disodium phosphate, glycerin, nonoxynol-9, sodium hydroxide, purified water

Satisfaction guaranteed - if dissatisfied, return unused portion and package to store where purchased. If unable to return to store, send reason for dissatisfaction, name address and empty package to: discount drug mart, 211 commerce drive, medina ohio 44256

DISTRIBUTED BY:
DRUG MART-FOOD FAIR
211 Commerce Dr.
MEDINA, OHIO Zip 44256
Questions or Comments? 1-800-833-6278
www.DISCOUNT-DRUGMART.COM



Rite Aid Label

Drug Facts

Active ingredient	Purpose
Povidone-iodine 10% (1% available iodine)	First Aid Antiseptic

Uses
First aid to help prevent infection in minor ■ cuts ■ scrapes ■ burns

Warnings For external use only
Do not use ■ in the eyes ■ over large areas of the body ■ if you are allergic to povidone-iodine or any other ingredients in this preparation

Ask a doctor before use if you have

- deep or puncture wounds ■ serious burns ■ animal bites

Stop use and ask a doctor if ■ the condition persists or gets worse ■ you need to use this product for more than 1 week

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- clean the affected area ■ apply a small amount of product to the area 1 to 3 times daily ■ may be covered with sterile bandage ■ if bandaged let dry first

Continued ▶

Safety Sealed*

ANTISEPTIC solution

POVIDONE-IODINE USP 10%

FIRST AID ANTISEPTIC
GERMICIDE

kills germs promptly and
helps prevent infections in:
minor cuts, burns, scrapes
and skin wounds

FOR EXTERNAL USE ONLY



8 FL OZ
(237 mL)

*Warning: Do not use if Tamper Evident Seal is broken or missing. This product is sealed with a foil seal under cap.

Drug Facts (continued)

Other information

- store at room temperature, 20-25° C (68-77° F)

Inactive ingredients
Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Purified Water and Sodium Hydroxide

Questions or Comments? 1-800-862-3435

DISTRIBUTED BY:
RITE AID, 30 HUNTER LANE,
CAMP HILL, PA 17011
www.riteaid.com

SATISFACTION GUARANTEE

If you're not satisfied, we'll
happily refund your money.



A47197
05 2022



first aid antiseptic solution

povidone-iodine 10%

topical antiseptic germicide

helps prevent infections in:

- burns
- cuts
- scrapes
- minor skin wounds

FOR EXTERNAL USE ONLY

8 FL OZ (237 mL)

Drug Facts

Active ingredient	Purpose
Povidone-iodine, 10% (equivalent 1% titratable iodine)	Antiseptic

Uses • first aid antiseptic to prevent infection in minor cuts and burns

Warnings For external use only

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and consult a doctor if

- the condition persists or gets worse
- irritation and redness develop and persists for more than 72 hours

When using this product do not use

- in the eyes
- on individuals who are allergic or sensitive to iodine or use for longer than 1 week unless directed by a doctor
- apply over large areas of the body

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Directions

- clean the affected area
- apply a small amount on the area 1 to 3 times daily
- may be covered with sterile bandage
- if bandaged, let dry first

Other Information

- store at room temperature, 20-25°C (68-77°F)

Inactive ingredients Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water


DISTRIBUTED BY: RITE AID
30 HUNTER LANE, CAMP HILL, PA 17011

IF YOU'RE NOT SATISFIED, WE'LL HAPPILY REFUND YOUR MONEY.

100% SATISFACTION GUARANTEE

ITEM 0037757

R092016RLG



0 11822 37757 7

CVS Label



Compare to the active ingredients in Betadine**

Povidone-Iodine SOLUTION

FIRST AID ANTISEPTIC

Kills germs promptly in minor cuts, scrapes & burns

10%

8 FL OZ (237 mL)

Drug Facts

Active ingredient	Purpose
Povidone-iodine 10% (1% available iodine)	First aid antiseptic

Uses first aid to help prevent infection in minor cuts scrapes burns

Warnings For external use only

Do not use

- in the eyes
- over large areas of the body

Ask a doctor before use in case of deep or puncture wounds, animal bites, or serious burns

Stop use and ask a doctor if

- the condition persists or gets worse
- you need to use this product for more than 1 week

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- clean the affected area
- apply a small amount of product to the area 1 to 3 times daily
- may be covered with a sterile bandage
- if bandaged, let dry first

Other information ■ store at 20°-25°C (68°-77°F)

Inactive ingredients citric acid, dibasic sodium phosphate, glycerin, nonoxynol-9, purified water, sodium hydroxide

Questions or comments? 1-800-719-9268

#955398

UPC# 050428410189

0 50428 48136 3

*This product is not manufactured or distributed by Purdue Products L.P., owner of the registered trademark Betadine®

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2015 CVS/pharmacy CVS.com® 1-800-SHOP CVS
Made in the U.S.A. V-17624

Vida Miaa Label

Drug Facts

Active ingredient **Purpose**
Povidone-Iodine 10%.....Antiseptic
(*Equivalent to 1% titratable iodine)

Use

First aid antiseptic to prevent infection in minor cuts and burns

Warnings

For external use only.

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and consult a doctor if

- the condition persists or gets worse
- irritation and redness develop and persists for more than 72 hours

When using this product do not

- do not use in the eyes
- use on individuals who are allergic or sensitive to iodine or use for longer than 1 week unless directed by a doctor
- do not apply over large areas of the body

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately. *Continued ▶*

Compare to
Betadine[®]*



Compare con
Betadine[®]*

mi salud

Povidone-Iodine
POVIDONA-YODADA

10%* TOPICAL SOLUTION USP
SOLUCIÓN TÓPICA AL 10%* USP

FIRST AID ANTISEPTIC

- First aid antiseptic to prevent infection in minor cuts and burns.

ANTISÉPTICO PARA PRIMEROS AUXILIOS

- Antiséptico de primeros auxilios para prevenir infecciones en cortaduras y quemaduras leves.

8 FL. OZ. (237 mL) • 8 ONZAS FL. (237 mL)

Drug Facts (continued)

Directions

- clean the affected area
- apply a small amount on the area 1 to 3 times daily
- may be covered with sterile bandage
- if bandaged let dry first

Other information

- store at room temperature, 20-25° C (68-77° F)

Inactive ingredients

Citric Acid, Disodium Phosphate, Glycerin, Nonoxonyl-9, Sodium Hydroxide and Purified Water

*This product is not manufactured or distributed by Purdue Products L.P., owner of the registered trademark Betadine[®].

Distributed by/ Distribuido por: R082413.RLG

MAGELLAN
DISTRIBUTION SOLUTIONS

Medley, Florida 33178

Questions or Comments? ☎1-800-862-3435



8 47717 06074 9

Levante etiquetas para información

Información sobre el Medicamento

Ingrediente activo **Propósito**
Povidona Yodada al 10%..... Antiséptico
(*Equivalente al 1% de yodo titulable)

Use

Antiséptico de primeros auxilios para prevenir infecciones en cortaduras y quemaduras leves.

Advertencias:

Para el uso externo solamente.

Consulte un médico antes de usar si usted tiene

- heridas o punciones profundas
- mordeduras de animales
- quemaduras graves

Suspenda el uso de este producto y consulte un médico si

- la condición persiste o empeora
- aparecen irritación y enrojecimiento y persisten por más de 72 horas.

Al utilizar este producto no

- No lo usa en los ojos
- NO use en personas alérgicas o sensibles al yodo. NO use por más de 1 semana a menos que sea ordenado por un médico
- No aplique sobre áreas extensas del cuerpo

Continuación ▶



Compare con
Betadine[®]*

mi salud

POVIDONA-YODADA
SOLUCIÓN TÓPICA AL 10%* USP

ANTISÉPTICO PARA PRIMEROS AUXILIOS

- Antiséptico de primeros auxilios para prevenir infecciones en cortaduras y quemaduras leves.

8 ONZAS FL. (237 mL)



Información sobre el Medicamento (continuación)

Mantener fuera del alcance de los niños. En caso de ingestión accidental, busque asistencia profesional o contacte un Centro de Control de Envenenamiento inmediatamente.

Instrucciones

- Limpie el área afectada
- aplique una cantidad pequeña en el área de 1 a 3 veces al día
- puede cubrirse con un vendaje estéril, pero deje secar antes de cubrir.

Otra información

- Almacene a temperatura ambiente, 20-25 ° C (68-77 ° F)

Ingredientes inactivos

Ácido Cítrico, Fosfato Disódico, Glicerina, Nonoxinol-9, Hidróxido de Sodio y Agua Purificada

Este producto no es fabricado ni distribuido por Purdue Products L.P., propietario de la marca registrada Betadine[®].

Distribuido por/ Distribuido por:

MAGELLAN
DISTRIBUTION SOLUTIONS

Medley, Florida 33178

¿Preguntas o comentarios? ☎1-800-862-3435

TIEDOWN

HEB Label

**This product is not manufactured or distributed by Purdue Products L.P., owner of the registered trademark Betadine®.



GUARANTEE
We believe the high quality of this H-E-B product makes it an outstanding value. We hope you'll agree. If not, we'll cheerfully refund your money. Thanks for shopping with us.

MADE WITH PRIDE AND CARE FOR H-E-B, SAN ANTONIO, TX 78204

Questions or Comments? Call 1-800-662-3435

0 1104



8

Compare to Betadine® Active Ingredient**



Povidone-Iodine 10%*

TOPICAL SOLUTION USP

First Aid Antiseptic

Use to Prevent Infection in Minor Cuts, Scrapes and Burns

8 FL OZ (237mL)

Drug Facts

Active ingredient	Purpose
Povidone-Iodine 10% (*Equivalent to 1% titratable iodine)	Antiseptic

Uses
First aid antiseptic to prevent infection in minor cuts and burns

Warnings For external use only
Ask a doctor before use if you have
 ■ deep or puncture wounds ■ animal bites ■ serious burns

Stop use and consult a doctor if
 ■ the condition persists or gets worse ■ irritation and redness develop and persists for more than 72 hours

When using this product do not
 ■ do not use in the eyes ■ use on individuals who are allergic or sensitive to iodine or use for longer than 1 week unless directed by a doctor ■ do not apply over large areas of the body

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Directions
 ■ clean the affected area ■ apply a small amount on the area 1 to 3 times daily ■ may be covered with sterile bandage ■ if bandaged let dry first

Other Information
 ■ store at room temperature, 20-25° C (68-77° F)

Inactive Ingredients
 citric acid, disodium phosphate, glycerin, nonoxynol-9, sodium hydroxide and purified water

Health Mart Label

Drug Facts	
Active ingredient	Purpose
Povidone-Iodine 10% (*Equivalent to 1% titratable iodine)	First Aid Antiseptic
Indications	
First aid antiseptic to help prevent infection in minor cuts, scrapes, and burns	
Warnings	
For external use only	
Ask a doctor before use if you have	
■ deep or puncture wounds	
■ animal bites	
■ serious burns	
Stop use and consult a doctor if	
■ the condition persists or gets worse	
■ irritation and redness develop and persists for more than 72 hours	
When using this product do not	
■ use in the eyes	
■ use on individuals who are allergic or sensitive to iodine or use for longer than 1 week unless directed by a doctor	
■ apply over large areas of the body	
Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.	
Continued ▶	

NDC 62011-0118-1

Compare to Betadine® Active Ingredient**

Povidone-Iodine

Topical Solution 10% USP

Antiseptic

• Kills germs and helps prevent the risk of infection in minor cuts, scrapes & burns

8 Fl. oz (236 mL)

Drug Facts (continued)
Directions
■ clean the affected area ■ apply a small amount on the area 1 to 3 times daily ■ may be covered with sterile bandage ■ if bandaged, let dry first
Other information
■ store at room temperature, 20-25° C (68-77° F)
Inactive ingredients
Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water

**This product is not manufactured or distributed by Purdue Products L.P., owner of the registered trademark Betadine®.

A44102
08 2021

McKESSON

Another Quality Product
Distributed by McKesson
One Post Street, San Francisco, CA 94104
Money Back Guarantee
healthmart.com





Health Mart
PHARMACY

NDC 62011-0118-1

compare to Betadine® Solution
active ingredient*

povidone-iodine

10% topical solution usp

kills germs and helps prevent the
risk of infection in minor cuts,
scrapes and burns

first aid antiseptic

8 fl oz (236 mL)

Drug Facts

Active Ingredient	Purpose
Povidone-Iodine 10%.....	Antiseptic
(*Equivalent to 1% titratable iodine)	

Uses
First aid antiseptic to prevent infection in minor cuts, burns.

Warnings For external use only

Do not:

- use in the eyes
- use longer than 1 week unless directed by a doctor
- use on individuals who are allergic or sensitive to iodine
- apply over large areas of the body

Stop use and consult a doctor if

- condition persists or gets worse
- irritation and redness develop and persist for more than 72 hours

Ask a doctor in case of:

- deep or puncture wounds
- animal bites
- serious burns

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Directions

- Clean the affected area
- Apply a small amount on the area 1 to 3 times daily
- May be covered with a sterile bandage
- If bandaged, let dry first

Other Information store at room temperature, 20-25° C (68-77° F)

Inactive Ingredients: Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water.

*This product is not manufactured or distributed by Purdue Frederick, owner of the registered trademark Betadine®

McKesson

Distributed by McKesson
One First Street, San Francisco, CA 94104
Money Back Guarantee
www.healthmart.com/healthmartbrand



Purdue Betadine Label

BETADINE
ANTISEPTIC



FIRST AID ANTISEPTIC
10% POVIDONE-IODINE

Solution

NO-STING PROMISE*

Kills Germs Promptly In
Minor Cuts, Scrapes & Burns

6 fl oz (177ml)

Do not use if seal under cap
is missing or damaged.

*Our promise or your money back.
Visit www.Betadine.com for details.

Drug Facts

Active ingredient	Purpose
Povidone-iodine, 10% (1% available iodine).....	First aid antiseptic

Uses first aid to help prevent infection in minor cuts scrapes burns

Warnings For external use only

Do not use in the eyes over large areas of the body if you are allergic to povidone-iodine or any other ingredients in this preparation

Ask a doctor before use if you have deep or puncture wounds serious burns animal bites

Stop use and ask a doctor if the condition persists or gets worse you need to use this product for more than 1 week

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away

Directions

- clean the affected area
- apply a small amount of product to the area 1 to 3 times daily
- may be covered with a sterile bandage
- if bandaged, let dry first

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients citric acid, disodium phosphate, glycerin, nonoxynol-9, purified water, sodium hydroxide

Questions? 1-888-827-0624 (8am-5pm, EST, Mon.-Fri.).

**FIRST AID
FOR THE WHOLE
FAMILY**

Helps protect against
skin infection

Golden brown color
indicates area treated



Betadine and Avrio Health are registered trademarks of Avrio Health L.P.
©2022 Avrio Health L.P.
Dist. by: Avrio Health L.P. Stamford, CT 06901-3431

ARIO
health®

305768-0A



BETADINE
ANTISEPTIC

50 YEARS
TRUSTED IN HOSPITALS FOR OVER

10% POVIDONE-IODINE
FIRST AID ANTISEPTIC

Solution

NO-STING PROMISE*
Kills Germs Promptly In Minor Cuts, Scrapes & Burns

8 fl oz (237ml)

Do not use if imprinted seal under cap is missing or damaged.

*Our promise or your money back. Visit www.Betadine.com for details.

Drug Facts	Purpose
Active ingredient Povidone-iodine, 10% (1% available iodine)	First aid antiseptic
Uses first aid to help prevent infection in minor ■ cuts ■ scrapes ■ burns	
Warnings For external use only Do not use ■ in the eyes ■ over large areas of the body ■ if you are allergic to povidone-iodine or any other ingredients in this preparation	
Ask a doctor before use if you have ■ deep or puncture wounds ■ serious burns ■ animal bites	
Stop use and ask a doctor if ■ the condition persists or gets worse ■ you need to use this product for more than 1 week	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away	
Directions ■ clean the affected area ■ apply a small amount of product to the area 1 to 3 times daily ■ may be covered with a sterile bandage ■ if bandaged, let dry first	
Other information ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)	
Inactive ingredients citric acid, disodium phosphate, glycerin, monoxynol-9, purified water, sodium hydroxide	
Questions? 1-888-827-0624 (8am-5pm, EST, Mon.-Fri.)	

FIRST AID FOR THE WHOLE FAMILY

Helps protect against skin infection

Golden brown color indicates area treated

BETADINE is a registered trademark of Avrio Health L.P. © 2020 Avrio Health L.P. Dist. by Avrio Health L.P. Stamford, CT 06901-3431

ARIO health

67618115008 5

Top Care Label

Drug Facts	
Active ingredient	Purpose
Povidone-Iodine 10%.....	First Aid Antiseptic (1% available iodine)
Uses First aid to help prevent infection in minor ■ cuts ■ scrapes ■ burns	
Warnings For external use only Do not use ■ in the eyes ■ over large areas of the body ■ if you are allergic to povidone-iodine or any other ingredients in this preparation	
Ask a doctor before use if you have ■ deep or puncture wounds ■ serious burns ■ animal bites	
Stop use and ask a doctor if ■ the condition persists or gets worse ■ you need to use this product for more than 1 week	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions ■ clean the affected area ■ apply a small amount of product to the area 1 to 3 times daily ■ may be covered with sterile bandage ■ if bandaged let dry first	
<i>Continued</i> ▶	

✓QUALITY GUARANTEED



*Safety Sealed

COMPARE TO BETADINE® ACTIVE INGREDIENT!

Povidone-Iodine

10% TOPICAL SOLUTION USP
FIRST AID ANTISEPTIC

- Kills Germs in Minor Cuts, Scrapes & Burns
- Used by Hospitals, Doctors & Nurses
- Water Soluble

8 FL OZ (237 mL)

*Warning: Do not use if Tamper Evident Seal is broken or missing. This product is sealed with a foil seal under cap.

Drug Facts (continued)

Other information

■ store at room temperature, 20-25° C (68-77° F)

Inactive ingredients

Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water

Questions or Comments? 1-888-423-0139

† This product is not manufactured or distributed by Purdue Products L.P., distributor of Betadine® First Aid Solution 10% Povidone-Iodine Topical Solution DISTRIBUTED BY TOPCO ASSOCIATES LLC, ELK GROVE VILLAGE, IL 60007 © TOPCO HHGA0521 QUESTIONS? 1-888-423-0139 topcare@topco.com www.topcarebrand.com



Scan here for more information or call 1-888-423-0139

R082021



Drug Facts	
Active ingredient	Purpose
Povidone-Iodine 10%.....	First Aid Antiseptic
(1% available iodine)	
Uses	
First aid to help prevent infection in minor ■ cuts ■ scrapes ■ burns	
Warnings For external use only	
Do not use ■ in the eyes ■ over large areas of the body ■ if you are allergic to povidone-iodine or any other ingredients in this preparation	
Ask a doctor before use if you have	
■ deep or puncture wounds ■ serious burns ■ animal bites	
Stop use and ask a doctor if ■ the condition persists or gets worse ■ you need to use this product for more than 1 week	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions	
■ clean the affected area ■ apply a small amount of product to the area 1 to 3 times daily ■ may be covered with sterile bandage ■ if bandaged let dry first	
<i>Continued ▶</i>	



*Safety Sealed

COMPARE TO BETADINE®
ACTIVE INGREDIENT*

Povidone-Iodine

10% TOPICAL SOLUTION USP
FIRST AID ANTISEPTIC

- Kills Germs in Minor Cuts, Scrapes & Burns
- Used by Hospitals, Doctors & Nurses
- Water Soluble



8 FL OZ (237 mL)

*Warning: Do not use if Tamper Evident Seal is broken or missing. This product is sealed with a foil seal under cap.

Drug Facts (continued)

Other information

■ store at room temperature, 20-25° C (68-77° F)

Inactive ingredients

Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water

Questions or Comments? 1-888-423-0139

† This product is not manufactured or distributed by Purdue Products L.P., distributor of Betadine® First Aid Solution 10% Povidone-Iodine Topical Solution
DISTRIBUTED BY TOPCO ASSOCIATES LLC, ELK GROVE VILLAGE, IL 60007
©2019PCN H4F4ADP10 011F5D1W57 1-888-423-0139
topcare@topco.com www.topcarebrand.com



Scan here
for more
information

A46495
04 2022

✓ QUALITY GUARANTEED



POVIDONE IODINE

povidone iodine 10% liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:82645-907
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POVIDONE-IODINE (UNII: 85H0HZU99M) (IODINE - UNII:9679TC07X4)	IODINE	100 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
GLYCERIN (UNII: PDC6A3C0OX)	
NONOXYNOL-9 (UNII: 48Q180SH9T)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date

1	NDC:82645-907-98	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/14/2017	
2	NDC:82645-907-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/14/2017	
3	NDC:82645-907-96	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/19/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	01/01/2008	

Labeler - Pharma Nobis, LLC (118564114)

Registrant - Pharma Nobis, LLC (118564114)

Establishment

Name	Address	ID/FEI	Business Operations
Pharma Nobis, LLC		118564114	manufacture(82645-907) , analysis(82645-907) , pack(82645-907) , label(82645-907)

Revised: 12/2023

Pharma Nobis, LLC