

**KROGER NIGHTTIME COLD AND FLU SOFTGEL- acetaminophen,
dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled
The Kroger Company**

Kroger NightTime Cold and Flu Softgel

Drug Facts

Active ingredients (in each Softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Purposes

Pain reliever / Fever reducer

Cough suppressant

Antihistamine

Uses:

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever
- runny nose & sneezing

Warnings:

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a

doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleep

Ask a doctor before use if you have • liver disease • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema • trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers • taking the blood thinning drug warfarin

When using this product • do not use more than directed. • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, & tranquilizers may increase drowsiness

Stop use and ask a doctor if • pain or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

Directions:

- take only as directed - see Overdose warning
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	2 Softgels with water every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information:

- store at room temperature between 20-25 °C (68-77 °F)

Inactive ingredients:

D&C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol

400, povidone, propylene glycol, purified water, sorbitol sorbitan solution

Questions or comments:

1-800-632-6900

COMPARE TO the active ingredients of VICKS® NYQUIL®

COLD & FLU NIGHTTIME RELIEF LIQUICAPS® *See bottom panel

POWERFUL NIGHTTIME RELIEF

OUR PHARMACIST RECOMMENDED

Multi-Symptom Relief

TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN OR SHOWS ANY SIGN OF TAMPERING

READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

**DISTRIBUTED BY
THE KROGER CO.,
CINCINNATI, OHIO 45202**

MADE IN UAE

For More Product Information, Scan UPC

Using Your Kroger App or Call 800-632-6900

*This product is not manufactured or distributed by The Proctor & Gamble Company. Vicks®, NyQuil®, and LiquiCaps® are registered trademarks of The Proctor & Gamble Company.

PARENTS:

Learn about teen medicine abuse


www.StopMedicineAbuse.org

Our Brands, Our Guarantee.

Love It or Your Money Back.

www.kroger.com

Packaging



COMPARE TO the active ingredients of VICKS® NYQUIL® COLD & FLU NIGHTTIME RELIEF LIQUICAPS® *See bottom panel

NDC 30142-086-48

Kroger

OUR PHARMACIST RECOMMENDED

Cold & Flu

NightTime

Multi-Symptom Relief

**Acetaminophen
Dextromethorphan HBr
Doxylamine Succinate**

ACTUAL SIZE

Pain Reliever
Fever Reducer
Cough Suppressant
Antihistamine

48 SOFTGELS

Kroger

Cold & Flu

NightTime

Multi-Symptom Relief

**Acetaminophen
Dextromethorphan HBr
Doxylamine Succinate**

Pain Reliever
Fever Reducer
Cough Suppressant
Antihistamine

48 SOFTGELS

Drug Facts

Active ingredients (in each Softgel)

Acetaminophen 325 mg.....Pain reliever / Fever reducer
Dextromethorphan HBr 15 mg.....Cough suppressant
Doxylamine succinate 6.25 mg.....Antihistamine

Uses: temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- minor aches & pains
- sore throat
- fever
- headache
- runny nose & sneezing

Warnings:

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses in 24 hrs, which is the maximum daily amount for this product

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- hives
- itching

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleep

Ask a doctor before use if you have liver disease, glaucoma, cough that occurs with too much phlegm (mucus), a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema, trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers, taking the blood thinning drug warfarin

When using this product do not use more than directed. excitability may occur, especially in children

- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, & tranquilizers may increase drowsiness

Stop use and ask a doctor if pain or cough gets worse or lasts more than 7 days, fever gets worse or lasts more than 3 days, redness or swelling is present, new symptoms occur, cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions:

- adults & children 12 yrs & over 2 softgels with water every 6 hrs
- children 4 to under 12 yrs ask a doctor
- children under 4 yrs do not use

Other information:

- store at room temperature between 20-25 °C (68-77 °F)

Inactive ingredients:

D&C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution

Questions or comments: 1-800-632-6900

Drug Facts (continued)

Directions:

- adults & children 12 yrs & over 2 softgels with water every 6 hrs
- children 4 to under 12 yrs ask a doctor
- children under 4 yrs do not use

Other information:

- store at room temperature between 20-25 °C (68-77 °F)

Inactive ingredients:

D&C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution

Questions or comments: 1-800-632-6900

Drug Facts (continued)

Directions:

- adults & children 12 yrs & over 2 softgels with water every 6 hrs
- children 4 to under 12 yrs ask a doctor
- children under 4 yrs do not use

Other information:

- store at room temperature between 20-25 °C (68-77 °F)

Inactive ingredients:

D&C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution

Questions or comments: 1-800-632-6900

DRUG FACTS TABLE

Drug Facts	
Active ingredients (in each Softgel)	Purposes
Acetaminophen 325 mg.....	Pain reliever / Fever reducer
Dextromethorphan HBr 15 mg.....	Cough suppressant
Doxylamine succinate 6.25 mg.....	Antihistamine
Uses: temporarily relieves common cold/flu symptoms:	
<ul style="list-style-type: none"> cough due to minor throat & bronchial irritation minor aches & pains 	<ul style="list-style-type: none"> sore throat fever headache runny nose & sneezing
Warnings:	
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses in 24 hrs, which is the maximum daily amount for this product	
<ul style="list-style-type: none"> with other drugs containing acetaminophen 3 or more alcoholic drinks daily while using this product 	
Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:	
<ul style="list-style-type: none"> skin reddening hives itching 	
If a skin reaction occurs, stop use and seek medical help right away.	
Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.	
Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.	
<ul style="list-style-type: none"> if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. to make a child sleep 	
Ask a doctor before use if you have liver disease, glaucoma, cough that occurs with too much phlegm (mucus), a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema, trouble urinating due to enlarged prostate gland	
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers, taking the blood thinning drug warfarin	
When using this product do not use more than directed. excitability may occur, especially in children	
<ul style="list-style-type: none"> marked drowsiness may occur avoid alcoholic drinks be careful when driving a motor vehicle or operating machinery alcohol, sedatives, & tranquilizers may increase drowsiness 	
Stop use and ask a doctor if pain or cough gets worse or lasts more than 7 days, fever gets worse or lasts more than 3 days, redness or swelling is present, new symptoms occur, cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children.	
Overdose warning: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.	
Drug Facts (continued)	
Directions:	
adults & children 12 yrs & over	2 softgels with water every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use
Other information:	
store at room temperature between 20-25 °C (68-77 °F)	
Inactive ingredients:	
D&C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution	
Questions or comments: 1-800-632-6900	

KROGER NIGHTTIME COLD AND FLU SOFTGEL

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:30142-086
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	808
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30142-086-48	4 in 1 CARTON	08/31/2022	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/31/2022	

Labeler - The Kroger Company (006999528)

Revised: 12/2024

The Kroger Company