

**BMD SOLUBLE- bacitracin methylene disalicylate powder, for solution
Phibro Animal Health**

BMD® Soluble

BMD® Soluble

(bacitracin methylenedisalicylate soluble powder)

Proportioner Pack

For poultry and livestock drinking water

Broiler and replacement chickens

An aid in prevention and control of necrotic enteritis caused by *Clostridium perfringens* susceptible to bacitracin methylenedisalicylate.

Growing turkeys

An aid in control of transmissible enteritis (blue comb, mud fever) in growing turkeys complicated by organisms susceptible to bacitracin methylenedisalicylate.

Growing quail

For the prevention of ulcerative enteritis due to *Clostridium colinum* susceptible to bacitracin methylenedisalicylate.

Swine

For use in the treatment of swine dysentery associated with *Brachyspira hyodysenteriae*.

Active ingredient

Contains 51.2 g bacitracin activity from bacitracin methylenedisalicylate equivalent to 200 g bacitracin activity per pound or to 18,520 units bacitracin (master standard) per gram.

Ingredients

Granulated sugar, sodium bicarbonate, anti-caking agent.

Caution

**Not for use in humans. Keep out of reach of children.
For oral use in animals only.**

Not for use in humans.

**Keep out of reach of children.
For oral use in animals only.**

Concentrated

For water medicators.

Directions for use

Animal	Condition	Prevention Control	
Broiler and replacement chickens	Necrotic enteritis caused by <i>Clostridium perfringens</i> susceptible to bacitracin methylenedisalicylate.	100 mg / gal	200 - 400 mg / gal
Growing turkeys	Transmissible enteritis (blue comb, mud fever) complicated by organisms susceptible to bacitracin methylenedisalicylate.	-	400 mg / gal
Growing quail	Ulcerative enteritis due to <i>Clostridium colinum</i> susceptible to bacitracin methylenedisalicylate..	400 mg / gal	-
Swine	Swine dysentery (bloody scours) associated with <i>Brachyspira hyodysenteriae</i> .	-	1,000 mg / gal

Store between 20°C -25°C (68°F - 77°F); excursions permitted between 15°C - 37°C (59°F - 99°F).

Restricted Drug (California) - Use only as directed.

Approved by FDA under NADA # 065-470

Phibro, Phibro logo and BMD are trademarks owned by or licensed to Phibro Animal Health Corporation or its affiliates.

Questions/Comments? For technical assistance or to obtain a Safety Data Sheet(s), contact Phibro Animal Health Corporation at 888-403-0074 or www.pahc.com. To report side effects, contact Phibro Animal Health at 1-888-475-7355. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <https://www.fda.gov/reportanimalae>

Prepare a fresh solution daily and use as sole source of drinking water.

Mixing instructions

For Proportioners

Select the treatment dosage. Set the proportioner at the desired delivery rate. To prepare the proportioner's stock solution, place the indicated quantity of BMD Soluble in a 2-gallon container, fill with water and stir until dissolved.

For Tanks

One package of BMD Soluble will medicate approximately 50 gallons of water at a dosage of 1000 mg/gallon.

PROPORTIONER SET AT: PROPORTIONER SET AT:		
Treatment dosage	1 ounce/gallon	2 ounces/gallon
100 mg/gal	1/2 pack	1/4 pack
200 mg/gal*	1 pack	1/2 pack
400 mg/gal	2 packs	1 pack
1,000 mg/gal	5 packs	2-1/2 packs

* 200 mg/gallon is equivalent to about 100 g of feed grade BMD per ton of feed

*200 mg/gallon is equivalent to about 100 g of feed grade BMD per ton of feed

To control an outbreak, start medication at first clinical signs of disease. Consult a diagnostic laboratory or veterinarian to determine the diagnosis and advice regarding the optimal level of drug. For necrotic enteritis in broiler and replacement chickens, administer continuously 5-7 days or as long as clinical signs persist, then reduce medication to prevention levels (100 mg/gal).

For transmissible enteritis in turkeys, administer continuously as long as clinical signs persist. For swine dysentery start medication at first signs of the disease or at time of exposure and administer continuously for 7 days or until signs of dysentery disappear. Treatment not to exceed 14 days.

CAUTION: Not for use in swine weighing more than 250 lb.

Distributed by:

Phibro Animal Health Corporation

Teaneck, NJ 07666

TAKE TIME



**OBSERVE LABEL
DIRECTIONS**

Take Time Statement

Net Weight 4.1 oz (116.2 g)

176-8018-01

PRINCIPAL DISPLAY PANEL - bag label

NO COPY

Prepare a fresh solution daily and use as sole source of drinking water.

Mixing instructions

For Proportioners:
Select the treatment dosage.
Set the proportioner at the desired delivery rate. To prepare the proportioner's stock solution, place the indicated quantity of BMD Soluble in a 2-gallon container, fill with water and stir until dissolved.

For Tanks:
One package of BMD Soluble will medicate approximately 50 gallons of water at a dosage of 1000 mg/gallon.

Treatment dosage	PROPORTIONER SET AT:
100 mg/gal	1 ounce/gallon
200 mg/gal*	1/2 pack
400 mg/gal	1 pack
1,000 mg/gal	2 packs
	5 packs

* 200 mg/gallon is equivalent to about 100 g of feed grade BMD per ton of feed.

To control an outbreak, start medication at first clinical signs of disease. Consult a diagnostic laboratory or veterinarian to determine the diagnosis and advised regarding the optimal level of drug. For necrotic enteritis in broiler and replacement chickens, administer continuously 5-7 days or as long as clinical signs persist, then reduce medication to prevention levels (100 mg/gal).

PROPORTIONER SET AT:
2 ounces/gallon
1/4 pack
1/2 pack
1 pack
2-1/2 packs

For transmissible enteritis in turkeys, administer continuously as long as clinical signs persist. For swine dysentery start medication at first signs of the disease or at time of exposure and administer continuously for 7 days or until signs of dysentery disappear. Treatment not to exceed 14 days.
CAUTION: Not for use in swine weighing more than 250 lb.

CORNER RADIUS



0 66104 67169 0

BMD® Soluble

(bacitracin methylenedisalicylate soluble powder)

NO COPY

Proportioner Pack

For poultry and livestock drinking water

Active ingredient

Contains 51.2 g bacitracin activity from bacitracin methylenedisalicylate equivalent to 200 g bacitracin activity per pound or to 15,200 units bacitracin (mild standard) per gram.

Ingredients

Granulated sugar, sodium bicarbonate, anti-caking agent.

Caution: Not for use in humans.

Keep out of reach of children.

For oral use in animals only.

Concentrated

For water medicators.

Store between 20°C - 25°C (68°F - 77°F); extremes permitted between 15°C - 37°C (59°F - 99°F).

Restricted Drug (Oftentimes - Use only as directed).

Approved by FDA under NADA # 036-470.

Phibro, Phibro logo and BMD are trademarks owned by or licensed to Phibro Animal Health Corporation or its affiliates.

Questions/Comments? For technical assistance or to obtain a Safety Data Sheet(s), contact Phibro Animal Health Corporation at 888-403-0074 or www.phibro.com. To report side effects, contact Phibro Animal Health at 1-888-475-7355. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportmeds>

Broiler and replacement chickens

An aid in prevention and control of necrotic enteritis caused by *Clostridium perfringens* susceptible to bacitracin methylenedisalicylate.

Growing turkeys

An aid in prevention and control of transmissible enteritis (blue comb, mud fever) complicated by organisms susceptible to bacitracin methylenedisalicylate.

Growing quail

For the prevention of ulcerative enteritis due to *Clostridium colinum* susceptible to bacitracin methylenedisalicylate.

Swine

For use in the treatment of swine dysentery associated with *Brachyspira hyodysenteriae*.

Take Time



Observe Label Directions

Net Weight: 4.1 oz (116.2 g)

176-6018-02

Distributed by:
Phibro Animal Health Corporation
Teterboro, NJ 07986

Phibro
ANIMAL HEALTH CORPORATION

CORNER RADIUS

Indications for use

Animal

Condition

Prevention

Control

Broiler and replacement chickens

Necrotic enteritis caused by *Clostridium perfringens* susceptible to bacitracin methylenedisalicylate.

100 mg / gal

200 - 400 mg / gal

Growing turkeys

Transmissible enteritis (blue comb, mud fever) complicated by organisms susceptible to bacitracin methylenedisalicylate.

-

400 mg / gal

Growing quail

Ulcerative enteritis due to *Clostridium colinum* susceptible to bacitracin methylenedisalicylate.

400 mg / gal

-

Swine

Swine dysentery (bloody scours) associated with *Brachyspira hyodysenteriae*.

-

1,000 mg / gal

NO COPY

Z Reference No.	FG Material No.	Description
514508/01	10006716	FOIL, BMD SOLUBLE 4.1OZ
Specification / DWG	Dimensions	Country
Z14-801228	8.011" x 11.625"	US
MPN No.	Date	Additional Information

Colors

BLACK

Distinct

PMS 543C

White

Fluorescent

BMD Soluble Label

BMD SOLUBLE

bacitracin methylene disalicylate powder, for solution

Product Information

Product Type	OTC TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:66104-1320
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BACITRACIN METHYLENEDISALICYLATE (UNII: JGV60J52FT) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN	200 mg in 0.45 kg

Product Characteristics

Color	brown (Tan to light brown)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66104-1320-1	50 in 1 CONTAINER		
1		0.1162 kg in 1 PACKET		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA065470	03/22/2007	

Labeler - Phibro Animal Health (006989008)