

**HEADACHE RELIEF- acetaminophen, aspirin, caffeine tablet, film coated  
L.N.K. International, Inc.**

-----  
**Convenience Valet 44-334B**

***Active ingredients (in each caplet)***

Acetaminophen 250 mg

Aspirin 250 mg (NSAID\*)

Caffeine 65 mg

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

Pain reliever

Pain reliever aid

***Uses***

- temporarily relieves minor aches and pains due to:
  - a cold
  - headache
  - arthritis
  - muscular aches
  - toothache
  - premenstrual and menstrual cramps

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- shock
- facial swelling
- asthma (wheezing)

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- rash
- blisters
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- 3 or more alcoholic drinks every day while using this product
- with other drugs containing acetaminophen

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if**

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - vomit blood
  - have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts for more than 10 days

- fever gets worse or lasts for more than 3 days
- redness or swelling is present
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours. Do not take more than 8 caplets in 24 hours.
- children under 12 years: ask a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

**Inactive ingredients**

corn starch, crospovidone, hypromellose, microcrystalline cellulose, povidone, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

**Questions or comments?**

**1-877-507-6516**

**Principal Display Panel**

**24/7 life**

BY 7-ELEVEN™

Extra Strength

**Headache Relief**

**Acetaminophen,**

Aspirin **(NSAID)**, Caffeine

Pain Reliever/Pain Reliever Aid

compare to **EXCEDRIN®**  
**EXTRA STRENGTH**  
**HEADACHE** active ingredients†

**24**  
**CAPLETS**

Actual Size

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED**  
**SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by Haleon CH SARL,  
owner of the registered trademark Excedrin® Extra Strength Headache.

50844 REV0224C33408 5641C-SE-08-24

DISTRIBUTED BY 7-ELEVEN, INC., IRVING, TX 75063 WWW.7-ELEVEN.COM

**Satisfaction Guaranteed 1-800-255-0711**

no print / no varnish area  
lot no. & exp. date

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING



B-1205-334B-08  
REV0224C33408

**KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

**Drug Facts**

**Active ingredients (in each caplet)**

Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID)	Pain reliever
Caffeine 65 mg	"nonsteroidal anti-inflammatory drug"

**Uses**

- temporarily relieves minor aches and pains due to:
  - a cold
  - toothache
  - muscular aches
  - headache
  - premenstrual and menstrual cramps
  - arthritis

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Aspirin alert:** Aspirin may cause a severe allergic reaction, which may include:
 

- hives
- facial swelling
- shock
- asthma (wheezing)

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
 

- skin redness
- blisters
- skin peeling
- itching

If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take
 

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen

**217life**  
BY 7-ELEVEN™

**Extra Strength Headache Relief**  
Acetaminophen, Aspirin (NSAID), Caffeine  
Pain Reliever/Pain Reliever Aid

compare to **EXCEDRIN® EXTRA STRENGTH HEADACHE** active ingredients!

**24 CAPLETS** | Actual Size

**Drug Facts (continued)**

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

**Inactive ingredients** corn starch, crospovidone, hypromellose, microcrystalline cellulose, povidone, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

**Questions or comments?** 1-877-507-6516

†This product is not manufactured or distributed by Halson CH SARL, owner of the registered trademark Excedrin® Extra Strength Headache. 50844 REV0224C33408 5641C-SE-08-24

DISTRIBUTED BY 7-ELEVEN, INC., IRVING, TX 75063 WWW.7-ELEVEN.COM

**Satisfaction Guaranteed 1-800-255-0711**

**Drug Facts (continued)**

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug or medicine for diabetes, gout, or arthritis
- if you are pregnant or breastfeeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
- if you are taking a blood thinner (anticoagulant) or steroid drug
- if you are taking a diuretic
- if you are taking a heart disease, liver cirrhosis, or kidney disease medicine
- if you have asthma
- if you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- if you have liver disease
- if you have a history of stomach problems, such as heartburn
- if you are taking a diuretic
- if you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- if you have asthma

**Stop use and ask a doctor if you experience any of the following signs of stomach bleeding:**

- black stools
- blood in vomit
- feeling faint
- stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present
- any new symptoms appear

**Directions**

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours. Do not take more than 8 caplets in 24 hours.
- children under 12 years: ask a doctor

**Drug Facts (continued)**

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product
- are age 60 or older
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if**

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

**Convenience Valet 44-334**

<b>HEADACHE RELIEF</b>			
acetaminophen, aspirin, caffeine tablet, film coated			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50844-258

**Route of Administration** ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	44;334
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-258-08	1 in 1 CARTON	07/15/2019	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/15/2019	

**Labeler** - L.N.K. International, Inc. (038154464)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(50844-258)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(50844-258)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(50844-258)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(50844-258)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(50844-258)

Revised: 12/2025

L.N.K. International, Inc.