

# **MEDIQUE AT HOME BACK PAIN RELIEF- magnesium salicylate tablet, film coated**

**Unifirst First Aid Corporation**

-----  
**Medique @ Home Back Pain Relief**

## ***Drug Facts***

### ***Active ingredient (in each caplet)***

Magnesium Salicylate Tetrahydrate (NSAID)\* 580 mg

(equivalent to 467.2 mg of anhydrous magnesium salicylate)

\*nonsteroidal anti-inflammatory drug

### ***Purpose***

Pain reliever

### ***Uses***

For the temporary relief of minor backache pain

### ***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

### ***Allergy alert:***

Magnesium salicylate may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

### ***Stomach bleeding warning:***

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcohol drinks every day while using this product
- take more or for a longer time than directed

**Do not use** if you are allergic to salicylates (including aspirin) or to any other pain reliever/fever reducer.

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
- if ringing in the ears or loss of hearing occurs
- new symptoms occur
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. **It is especially important not to use magnesium salicylate at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

***Directions***

- drink a full glass of water with each dose

**Adults and children: (12 years and over)** Take 2 caplets every 6 hours, not to exceed 8 caplets in 24 hours

**Children under 12 years:** Do not use unless directed by a doctor

***Other information***

- each caplet contains: magnesium 40 mg

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

■ **TAMPER EVIDENT** child resistant packets

■ do not use any opened or torn packets

### Inactive ingredients

hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, stearic acid, titanium dioxide, triacetin

**Questions or comments?** 1-800-634-7680

Medique® @ Home

Back Pain Relief

Magnesium Salicylate Tetrahydrate

(NSAID) 580 mg

Pain Reliever

Pull to Open

Child-Resistant Packets

\*not actual size

50 Packets

(2 Caplets Per Packet)

**Medique® @ Home**  
**Back Pain Relief**  
Magnesium Salicylate Tetrahydrate (NSAID) 580 mg Pain Reliever  
Child-Resistant Packets  
50 Packets (2 Caplets Per Packet)

**Medique® @ Home**  
**Back Pain Relief**  
Magnesium Salicylate Tetrahydrate (NSAID) 580 mg Pain Reliever  
Child-Resistant Packets  
50 Packets (2 Caplets Per Packet)

**Drug Facts**  
**Active ingredient (in each caplet)**  
Magnesium Salicylate Tetrahydrate (NSAID)<sup>1</sup> 580 mg  
(Equivalent to 582 mg of anhydrous magnesium salicylate)  
**Purpose**  
Pain reliever  
**Nonsteroidal anti-inflammatory drug**

**Warnings**  
Reynolds syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Always alert.** Magnesium salicylate may cause a severe allergic reaction which may include:  
● hives ● facial swelling ● asthma (wheezing) ● shock  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
● are age 60 or older ● have had stomach ulcers or bleeding problems  
● take a blood thinner (anticoagulant) or steroid drug  
● take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
● have 3 or more alcohol drinks every day while using this product  
● take more or for a longer time than directed

**Do not use if you are allergic to salicylates (including aspirin) or to any other pain reliever/fever reducer.**  
**Ask a doctor before use if:**  
● stomach bleeding warning applies to you  
● you have a history of stomach problems such as heartburn  
● you have high blood pressure, heart disease, liver disease, or kidney disease  
● you have asthma  
● you are taking aspirin

**Ask a doctor or pharmacist before use if you are taking a prescription drug for:** ● gout ● diabetes ● arthritis

**Stop use and ask a doctor if:**  
● an allergic reaction occurs. Seek medical help right away.  
● you experience any of the following signs of stomach bleeding:  
● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better  
● occurs in the area of loss of hearing occur  
● new symptoms occur  
● pain gets worse or lasts more than 10 days  
● fever gets worse or lasts more than 3 days  
● worsens or swelling is present  
● These could be signs of a serious condition. (continued on back panel)

**Drug Facts (continued)**  
If pregnant or breast-feeding, ask a health care professional before use. It is especially important not to use magnesium salicylate at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**  
● drink a full glass of water with each dose  
**Adults and children (12 years and over)** Take 2 caplets every 6 hours, not to exceed 8 caplets in 24 hours.  
**Children under 12 years:** Do not use unless directed by a doctor.

**Other information**  
● each caplet contains magnesium 40 mg  
● store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)  
● **TAMPER EVIDENT** child-resistant packets  
● do not use any opened or torn packets

**Inactive ingredients**  
hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, stearic acid, titanium dioxide, triacetin

**Questions or comments?** 1-800-634-7680

MDQ074B  
7473380X  
Rev. 7/2012  
Mfg. 24

100 Caplets 50 Packets (2 Tablets Per Packet)  
50 Packets (2 Caplets Per Packet)  
Retain carton for complete product information  
3 47682 74733 6 www.medicqueproducts.com

## MEDIQUE AT HOME BACK PAIN RELIEF

magnesium salicylate tablet, film coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-747
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>MAGNESIUM SALICYLATE</b> (UNII: 41728CY7UX) (SALICYLIC ACID - UNII:O414PZ4LPZ)	MAGNESIUM SALICYLATE	580 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ05DW1A)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (Caplet)	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	44;338
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-747-33	50 in 1 BOX	09/26/2022	
1	NDC:47682-747-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-747-99	2 in 1 PACKET; Type 0: Not a Combination Product	09/26/2022	

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M013	09/26/2022	

**Labeler** - Unifirst First Aid Corporation (832947092)

Revised: 9/2025

Unifirst First Aid Corporation